

### **REMARKS**

Claims 1-37 are pending in the patent application. Claims 38-86 have been cancelled without prejudice as a result of a restriction requirement. These claims are being pursued in other patent applications. New claims 87-189 have been added to the application. Claims 7, 8, 27, and 28 have been amended to eliminate the phrase "the step" to be consistent with the other dependent claims and to make clear that the claims are not "step-plus-function" claims. No new matter has been entered. After entry of these amendments, claims 1-37 and 87-189 are present in the patent application.

#### **I. Restriction Requirement/Election**

The Applicants are confirming the election made by John Gatz on April 24, 2003 of claims 1-37. The remaining claims of the application (claims 38-86) are being cancelled without prejudice as a result of this restriction requirement.

#### **II. 35 U.S.C. § 103(a) Rejections**

##### **A. Embodiments/Methods of the Present Invention**

The Applicants are submitting herewith evidence in the form of a 37 C.F.R. §1.132 declaration by one of the co-inventors Mr. Gary R. DelDuca ("the DelDuca Declaration") (Exhibit A) to assist in explaining the present invention and showing the non-obviousness of the invention.

The present invention is directed to novel methods of manufacturing modified atmosphere packages. For example, independent claims 1, 22, 172, and 181 recite, *inter alia*, "a first package including a non-barrier portion substantially permeable to oxygen", "a second package substantially impermeable to oxygen", and a low oxygen environment that includes from about 0.1 to about 0.8 vol% carbon monoxide (CO). Independent claims 91 and 107 recite, *inter alia*, "a first package that is substantially impermeable to oxygen", "exposing the raw meat to ambient atmosphere by modifying the first package", "after exposing the raw meat to ambient atmosphere, placing the raw meat into a second package that includes a non-barrier portion substantially permeable to oxygen", and a low oxygen environment that includes from about 0.1 to about 0.8 vol% CO. Independent claims 123, 142, and 161 recite, *inter alia*, "a first package

including a non-barrier portion substantially permeable to oxygen", "a second package substantially impermeable to oxygen", and a low oxygen environment that includes less than 0.8 vol% CO or from about 0.3 to about 0.5 vol% CO.

The methods of manufacturing the modified atmosphere packages have several advantages as follows: (a) the "seasoning" period of the raw meat may be reduced or eliminated; (b) the ability to obtain consistent blooming with cuts off pigment-sensitive meats (*e.g.*, round bone) is improved; and (c) the ability to avoid "fixing" the color of the meat pigment to red. *See, e.g.*, page 11, line 29 - page 12, line 15; page 13, lines 11-17 of the application and DelDuca Decl. ¶ 4.

The "seasoning" period is the time period needed to diffuse the oxygen so that the meat has the ability to fully bloom. Page 3, lines 17-19 of the application and DelDuca Decl. ¶ 5. Trays, such as polystyrene foam trays, have a substantial amount of oxygen contained in its cellular structure that results in a time period of as long as about 5 to about 6 days to diffuse the oxygen contained in its cellular structure. Page 3, lines 21-23 of the application and DelDuca Decl. ¶ 5. If a foam tray is not used, the "seasoning" period can be reduced to one or two days. Page 3, lines 24-25 of the application and DelDuca Decl. ¶ 5. The reduction or elimination of the seasoning period "allows the meat to be displayed for retail sale much sooner than in existing low oxygen packaging systems." Page 11, line 32 - page 12, line 2 of the application and DelDuca Decl. ¶ 5. Seasoning periods are not desired by the retailers or packers because of the "need to store and maintain the meat-filled packages for an extended duration before being opened for retail sale." Page 3, lines 27-28 of the application and DelDuca Decl. ¶ 5.

The present invention does not "fix" the color of the meat pigment to red with its use of CO, but rather the meat pigment tends to turn brown in a natural time period. *See* page 12, lines 10-12 of the application and DelDuca Decl. ¶ 6. It is important to prevent the meat color from being "fixed" because it is unsafe (and potentially dangerous) to consume a piece of meat that has a bright red color that consumers associate with freshness, but has an unacceptable amount of bacteria. DelDuca Decl. ¶ 7.

The present invention "surprisingly allows the meat pigment to convert to metmyoglobin in a similar fashion as fresh, raw meat in a retail environment." Page 12, lines 7-10 of the application and DelDuca Decl. ¶ 7. Specifically, the color of the meat after exposure to ambient

temperature degrades in a fashion not beyond the point of microbial soundness as if the CO had never been added to the modified packaging system. DelDuca Decl. ¶ 7.

The meat used in the modified atmosphere packaging of the present invention substantially maintains its color during the shipping process because the package has a modified atmosphere that includes from about 0.1% to about 0.8% carbon monoxide. DelDuca Decl. ¶ 8. In one method, after removal of the package that is substantially permeable to oxygen, the CO is lost to the atmosphere. See page 12, lines 2-6 of the application and DelDuca Decl. ¶ 8. The CO may be lost to the atmosphere through the package that includes a non-barrier portion that is substantially permeable to oxygen. See *id.* and page 13, lines 5-10 of the application. This allows the conversion of the carboxymyoglobin to oxymyoglobin by using the oxygen from the air. Page 12, lines 4-7 of the application and DelDuca Decl. ¶ 8. The “gas mixture used in the modified atmosphere packages of the present invention, after removal, allows the carboxymyoglobin to convert to oxymyoglobin and then to metmyoglobin (brown) in a natural time period.” Page 12, lines 3-5 of the application and DelDuca Decl. ¶ 8. Thus, the color of the meat is not “fixed” in the present invention.

**B. A *Prima Facie* Case Has Not Been Presented with Respect to Independent Claims 1 and 22**

The two pending independent claims (claims 1 and 22) include, *inter alia*, (a) “a first package including a non-barrier portion substantially permeable to oxygen”, (b) “a second package substantially impermeable to oxygen”, and (c) a low oxygen environment that includes from about 0.1 to about 0.8vol.% carbon monoxide (CO).

None of the applied references includes, *inter alia*, such limitations that are recited in independent claims 1 and 22.

As acknowledged in the Office Action, the article entitled “The storage life of beef and pork packaged in an atmosphere with low carbon monoxide and high carbon dioxide” from *Meat Science* to Sorheim (“Sorheim”) does not disclose a packaging system having (a) “a first package including a non-barrier portion substantially permeable to oxygen”, and (b) “a second package substantially impermeable to oxygen”, as recited in independent claims 1 and 22.

Rather, Sorheim discloses several meat packaging systems consisting of trays wrapped in polymeric film. Sorheim at page 158. The Sorheim meat packaging systems include the use of 0.4% CO, 60% CO<sub>2</sub>, and 40% N<sub>2</sub>. *Id.* Sorheim also discloses the use of CO in amounts of approximately 0.3 to 0.5%. *Id.* at page 157.

Unlike the present invention, the Sorheim meat packaging systems do not address the problem of "fixing" the meat color because the CO remains in the Sorheim sealed packaging system until the polymeric film is removed by the customer. In other words, during retail display, the CO remains in contact with the meat in the Sorheim sealed package. *See, e.g., id.* at page 157 (Sorheim discloses that "a low CO/high CO<sub>2</sub> atmosphere is effective for preserving retail-ready meat."). The Sorheim meat packaging systems, therefore, can undesirably fix the color of the meat pigment to red.

The other applied references of U.S. Patent No. 6,112,890 to Columbo ("Columbo") and U.S. Patent No. 5,711,978 to Breen ("Breen") do not disclose the use of CO in their respective packaging system, let alone the claimed amount of CO recited in the independent claims. Colombo discloses the use of solid carbon dioxide in its packaging system. *See, e.g.,* col. 3, lines 19-39; col. 7, lines 10-29; and col. 9, lines 42-45. Breen discloses the use of a "substantially pure carbon dioxide" environment. *See, e.g.,* col. 5, lines 10-14.

There is no teaching or suggestion to combine the references of Sorheim and Columbo, or Sorheim and Breen. The mere fact that references can be combined together or modified does not render the resultant combination obvious unless the prior art also suggests the desirability of the combination. *In re Mills*, 916 F.2d 680 (Fed. Cir. 1990). "When a rejection depends on a combination of prior art references, there must be some teaching, suggestion, or motivation to combine the references." *Rouffet*, 149 F.3d at 1355, 47 U.S.P.Q.2d at 1456, (*citing In re Geiger*, 815 F.2d 686, 688, 2 U.S.P.Q.2d 1276, 1278 (Fed. Cir. 1987)). Evidence of a suggestion, teaching, or motivation to combine "must be clear and particular." *Ex parte Maruyama*, 2001 WL 1918556, \*3 (Bd. Pat. App. & Inter. 2001), (*citing C.R. Bard, Inc. v. M3 Sys. Inc.*, 157 F.3d 1340, 1352, 48 U.S.P.Q.2d 1225, 1232 (Fed. Cir. 1998)). None of the applied references addresses the problem of "fixing" the meat color to a bright red color with CO. Preventing the fixing of the meat color is important because it is unsafe (and potentially dangerous) to consume



a piece of meat that has a bright red color that consumers associate with freshness, but has an unacceptable amount of bacteria. DelDuca Decl. ¶ 7.

Sorheim not only do not address the problem of fixing the color, but rather discloses that its meat packaging systems with a modified atmosphere of "0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> had a bright stable red colour that lasted beyond the time of spoilage." Abstract of Sorheim. This is exactly the problem that the present invention overcomes. Neither Columbo nor Breen mentions CO, let alone recognizes the problem of fixing the color.

Thus, there is no motivation to combine these references to address the problem and consequently the solution proposed in the present invention. Additionally, Sorheim, the only applied reference that proposes a packaging system with CO, believes that its packaging solution is effective. *See generally* abstract of Sorheim. The authors of Sorheim, who are members of the Norwegian Food Research Institute or the Norwegian Meat Cooperative, also state that the Norwegian meat industry "is seeking amendments of current EU food regulations relating to the use of CO in MAP [modified atmosphere packaging] of red meats." Sorheim at pages 157, 158 (noting that the EU as of 1999 does not allow CO in its MAP of meat). Thus, it is clear that Sorheim believes that its meat packaging systems are desirable and there is no teaching or incentive to combine Sorheim with either Columbo, Breen, or the combination thereof.

Obviousness cannot "be established using hindsight or in view of the teachings or suggestions of the invention." *Ex parte Maguire* (Appendix 9), 2002 WL 1801466, \*3 (Bd. Pat. App. & Inter. 2002), (*quoting Para-Ordnance Mfg. Inc. v. SGS Importers Int'l Inc.*, 73 F.3d 1085, 1087, 37 U.S.P.Q.2d 1237, 1239 (Fed. Cir. 1995), *cert. denied*, 519 U.S. 822 (1996)). In other words, the knowledge to combine "can not come from the applicant's invention itself." *In re Oetiker*, 977 F.2d 1443, 1447, 24 U.S.P.Q.2d 1443, 1446 (Fed. Cir. 1992) (emphasis added).

Thus, the Applicants believe that a *prima facie* case has not been presented with Sorheim, Columbo, Breen, or any combination thereof.

### **C. Evidence of Non-Obviousness of Independent Claims 1 and 22**

Assuming, *arguendo*, that a *prima facie* case has been presented (which Applicants believe is not the case), the Applicants are submitting evidence of non-obviousness in the form of

a 37 C.F.R. §1.132 declaration by one of the co-inventors Mr. Gary R. DelDuca ("the DelDuca Declaration")(Exhibit A) to assist in showing the non-obviousness of the invention.

**i. CO Not Allowed with Fresh Meat in the United States Since At Least 1962**

Carbon monoxide (CO) has not been allowed to be used with fresh meat in the United States for about 40 years.<sup>1,2</sup> The Food and Drug Administration ("FDA") regulation that currently prevents using CO with meat packaging systems in the United States is 21 C.F.R. § 173.350.

The food additive combustion product gas may be safely used in the processing and packaging of the foods designated in paragraph (c) of this section for the purpose of removing and displacing oxygen...(b) The food additive meets the following specifications: (1) Carbon monoxide content not to exceed 4.5 percent by volume...(c) It [carbon monoxide] is used or intended for use to displace or remove oxygen in the processing, storage, or packaging of beverage products and other food, except fresh meats.

Exhibit D (emphasis added); *see also* DelDuca Decl. ¶ 9.

The concern of the FDA is believed to be that CO fixes the fresh meat color to a degree that allows the retailer to sell meat that looks good (a bright red color), but is unsafe and potentially dangerous to consume because it has unacceptable levels of bacteria. DelDuca Decl. ¶ 10.<sup>3</sup> This act of fixing the meat color to a bright red color is referred to as "economic adulteration." DelDuca Decl. ¶ 10.

**ii. CO Now Allowed In Pactiv's ActiveTech™ Meat Packaging System**

After about 40 years of not allowing CO to be used with fresh meats in the United States, the Applicants came up with novel approaches of using CO in modified atmosphere packaging (MAP) systems that avoided the concerns of "fixing" the meat color. DelDuca Decl. ¶ 11. The

---

<sup>1</sup> 21 U.S.C. § 121.1060 was first promulgated on August 2, 1961 (Exhibit B) and permitted the use of combustion product gas containing up to 4.5% CO for use "to displace or remove oxygen or remove oxygen in the processing, storage, or packaging of citrus products, vegetable fats and vegetable oils, coffee, and wine." In December 14, 1962, 21 U.S.C. § 121.1060 (Exhibit C) was amended to exclude fresh meats. In March of 1977, 21 U.S.C. § 121.1060 was re-designated as 21 C.F.R. § 173.350.

<sup>2</sup> *See also* DelDuca Decl. ¶ 9.

<sup>3</sup> *See, e.g.,* Exhibit E (In a 1962 letter, the FDA told a Whirlpool representative that it might need additional data "to establish that the treatment of meat would not serve to cause the meat to retain its fresh red color longer than meat not so treated" and that the FDA has a question "concerning possible deception of the consumer where treatment of the meat leads to longer retention of the fresh red color.")

assignee of the patent application (Pactiv Corporation) gave notice to the FDA of a specific embodiment and process and evidence supporting Pactiv's conclusion that CO as used is GRAS (generally recognized as safe). *See* Exhibit F; DelDuca Decl. ¶ 12. The MAP system in Pactiv's GRAS notice may be used for packaging meats such as fresh cuts of case-ready muscle meat and ground case-ready meat to maintain wholesomeness, provide flexibility in distribution, and losses due to spoilage at retail sale. Exhibit F at page 5; DelDuca Decl. ¶ 12.

The specific MAP system that was presented in the GRAS notice used 0.4 % CO in a meat packaging system and was referred to as Pactiv's ActiveTech™ meat packaging system. Exhibit F at page 5; DelDuca Decl. ¶ 13. The ActiveTech™ meat packaging system traditionally includes meats being placed in polystyrene trays and covered with oxygen-permeable, polyvinyl chloride ("PVC") overwraps. Exhibit F at page 6; DelDuca Decl. ¶ 13. The wrapped trays of meat are then placed in an outer barrier bag. Ambient air is removed and replaced with a blend of 0.4% CO, 30% carbon dioxide, and the balance being nitrogen. Exhibit F at page 8; DelDuca Decl. ¶ 13. The myoglobin of the meat converts from oxymyoglobin to carboxymyoglobin (red). Exhibit F at page 9; DelDuca Decl. ¶ 13. The meat maintains its red color while in storage until the package is opened for retail display by removing the outer barrier bag. Exhibit F at page 9; DelDuca Decl. ¶ 13.

The package will lose CO to the atmosphere and, thus, at retail display will not have CO. DelDuca Decl. ¶ 12. Once in retail display, the meat's myoglobin begins its natural conversion to metmyoglobin (brown). Exhibit F at page 9; DelDuca Decl. ¶ 13. The CO used in the MAP system did not mask the spoilage or extend the color life beyond the point of wholesomeness (*i.e.*, the point of microbial soundness). Exhibit F at pages 9-10; DelDuca Decl. ¶ 13.

The FDA stated that it had no questions regarding Pactiv's conclusion about Pactiv's ActiveTech™ meat packaging system using 0.4% CO being GRAS because of the evidence presented by Pactiv in its notice. Exhibit G at page 1; DelDuca Decl. ¶ 14. Specifically, the FDA stated the following: "Based on the data and information reviewed, Pactiv's GRAS panel conclude[d] that CO, when produced in accordance with current good manufacturing practice and meeting appropriate food grade specifications, is GRAS, though scientific procedures under the conditions of its intended use." Exhibit G at page 1. This FDA review allows Pactiv to use

CO with fresh meat in its application. DelDuca Decl. ¶ 14. It is believed to be the first system to overcome the prohibition of CO with fresh meat in the United States in the last 40 years. *Id.*

Thus, a problem of fixing meat color with CO that was recognized for at least the last 40 years was overcome by Pactiv's meat packaging system and process of the same. *See, e.g.*, Exhibits E, F, and G. The Pactiv process in the GRAS notice is an example of one process that would be covered by claims 1 and 22 of the present invention. Specifically, the Pactiv process included (a) a first package that included a PVC overwrap, (b) a retail cut of raw meat within the first package that was sealed, (c) a second outer barrier bag that covered the first package, (d) a gas mixture that comprised 0.4 vol.% that formed a low oxygen environment so as to form carboxymyoglobin on a surface of the raw meat, and (e) sealing the second package.

Therefore, in addition to the applied references not presenting a *prima facie* case, the Applicants also believe that the present invention is allowable because of the compelling evidence of non-obviousness. Therefore, independent claims 1 and 22 are believed to be allowable over the applied references of Sorheim, Columbo, Breen, or any combination thereof.

**D. Dependent Claims 2-21, 23-37 and 87-90**

In dependent claims 12, 19, 31, 50 and 53, further references were applied in combination with either Sorheim and Columbo, or Sorheim and Breen. Neither DE 1935566 to Verbruggen ("Verbruggen") nor U.S. Patent 4,522,835 to Woodruff ("Woodruff") addresses the deficiencies in the previously applied references of Sorheim, Columbo, Breen, or any combination thereof. Dependent claims 2-21, 23-37 and 87-90, which depend directly or indirectly on independent claim 1 or 22, are believed to be allowable over Sorheim, Columbo, Breen, Woodruff, Verbruggen, or any combination thereof for at least the same reasons as discussed with respect to claims 1 and 22.

**E. Independent Claims 91, 107, 123, 142, 161, 172, and 181**

Independent claims 91 and 107 recite, *inter alia*, "a first package that is substantially impermeable to oxygen", "exposing the raw meat to ambient atmosphere by modifying the first package", "after exposing the raw meat to ambient atmosphere, placing the raw meat into a second package that includes a non-barrier portion substantially permeable to oxygen", and a low oxygen environment that includes from about 0.1 to about 0.8 vol% CO. Independent claims 123, 142, and 161 recite, *inter alia*, "a first package including a non-barrier portion substantially


permeable to oxygen", "a second package substantially impermeable to oxygen", and a low oxygen environment that includes less than 0.8 vol% CO or from about 0.3 to about 0.5 vol% CO. Independent claims 172 and 181 recite, *inter alia*, "a first package including a non-barrier portion substantially permeable to oxygen", "a second package substantially impermeable to oxygen", and a low oxygen environment that includes from about 0.1 to about 0.8 vol% carbon monoxide (CO). Independent claims 91, 107, 123, 142, 161, 172, and 181 are believed to be allowable over the references of Sorheim, Columbo, Breen, or any combination thereof for at least the same reasons as discussed above with respect to claims 1 and 22. Dependent claims 92-106, 108-122, 124-141, 143-160, 162-171, 173-180, and 182-189, which depend directly or indirectly on independent claim 91, 107, 123, 142, 161, 172, or 181, are believed to be allowable over Sorheim, Columbo, Breen, Woodruff, Verbruggen, or any combination thereof for at least the same reasons as discussed with respect to claims 1 and 22.

**F. Conclusion**

The Applicants submit that the claims are in a condition for allowance and action toward that end is earnestly solicited. The Applicants have enclosed a check in the amount of \$1,400.00, which includes a fee of \$110.00 for a one month extension of time. It is believed that no further fees are due; however, should any additional fees be required (except for payment of the issue fee), the Commissioner is authorized to deduct the fees from Jenkins & Gilchrist, P.C. Deposit Account No. 10-0447, Order No. 47097-01080.

Respectfully submitted,

September 8, 2003  
Date

  
\_\_\_\_\_  
John C. Gatz  
Reg. No. 41,774  
Jenkins & Gilchrist, P.C.  
225 West Washington Street, Suite 2600  
Chicago, IL 60606-3418  
(312) 425-3900

Attorneys for Applicants

**PATENT**

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

Appl. No. : 09/915,150  
Applicant : Gary R. DelDuca *et al.*  
Filed : July 25, 2001  
Title : Modified Atmospheric Packages and Methods for Making the Same  
  
TC/A.U. : 1761  
Examiner : Robert A. Madsen  
  
Docket No. : 47097-01080

**DECLARATION OF GARY R. DELDUCA  
UNDER 37 C.F.R. § 1.132**

Mail Stop Amendments  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313

**CERTIFICATE OF MAILING  
37 C.F.R. 1.8**

I hereby certify that this correspondence is being deposited with the U.S. Postal Service as First Class Mail in an envelope addressed to: Mail Stop Amendments, Commissioner for Patents, P.O. Box 1450, Alexandria, Virginia 22313 on the date indicated below:

September 8, 2003  
Date

  
Adrienne White

Dear Commissioner:

I, Gary R. DelDuca, declare that:

1. I hold a degree of B.S. in Mechanical Engineering From Rochester Institute of Technology in Rochester, New York that was obtained in 1980.

2. From 1980-1995, I worked as a developmental and senior engineer for Mobil Chemical Company, Plastics Division. As a developmental engineer, I worked in process and product development in the area of foam products. As a senior engineer, some of my responsibilities included designing specialized machinery that included machinery directed to stacking trays for meat processes. Mobil Chemical Company, Plastics Division was purchased by Tenneco Inc. in 1995. From 1995 to the present, I have been a Technical Manager for Tenneco Packaging Inc. in the area of modified atmosphere packaging (MAP) for meats. My responsibilities have included designing, developing, and implementing such modified

atmosphere packaging for meat and processes using the same. In 1999, Tenneco Packaging Inc. was renamed Pactiv Corporation.

4. The present invention is directed to methods of manufacturing a modified atmosphere package that includes carbon monoxide (CO). The invention has several advantages: (a) the "seasoning" period of the raw meat may be reduced or eliminated; (b) the ability to obtain consistent blooming with cuts off pigment-sensitive meats (*e.g.*, round bone) is improved; and (c) the ability to avoid "fixing" the color of the meat pigment to red. *See, e.g.*, page 11, line 29 - page 12, line 15; page 13, lines 11-17 of the application.

5. The "seasoning" period is the time period needed to diffuse the oxygen so that the meat has the ability to fully bloom. Page 3, lines 17-19 of the application. Trays, such as polystyrene foam trays, have a substantial amount of oxygen contained in its cellular structure that results in a time period of as long as about 5 to about 6 days to diffuse the oxygen contained in its cellular structure. Page 3, lines 21-23 of the application. If a foam tray is not used, the "seasoning" period can be reduced to one or two days. Page 3, lines 24-25 of the application. The reduction or elimination of the seasoning period "allows the meat to be displayed for retail sale much sooner than in existing low oxygen packaging systems." Page 11, line 32 - page 12, line 2 of the application. Seasoning periods are not desired by the retailers or packers because of the "need to store and maintain the meat-filled packages for an extended duration before being opened for retail sale." Page 3, lines 27-28 of the application.

6. Importantly, the present invention does not "fix" the color of the meat pigment to red with its use of CO, but rather the meat pigment tends to turn brown in a natural time period. *See* page 12, lines 10-12 of the application.

7. It is important to prevent the meat color from being "fixed" because it is unsafe (and potentially dangerous) to consume a piece of meat that has a bright red color that consumers associate with freshness, but has an unacceptable amount of bacteria. The present invention "surprisingly allows the meat pigment to convert to metmyoglobin in a similar fashion as fresh, raw meat in a retail environment." Page 12, lines 7-10 of the application. Specifically, the color of the meat after exposure to the ambient atmosphere degrades in a fashion not beyond the point of microbial soundness as if the CO had never been added to the modified packaging system.

8. The meat used in the modified atmosphere packaging of the present invention substantially maintains its color during the shipping process because the package has a modified atmosphere that includes from about 0.1% to about 0.8% CO. In one method, after removal of the package that is substantially permeable to oxygen, the CO is lost to the atmosphere. See page 12, lines 2-6 of the application. The CO may be lost to the atmosphere through the package that includes a non-barrier portion that is substantially permeable to oxygen. See *id.* and page 13, lines 5-10 of the application. This allows the conversion of the carboxymyoglobin to oxymyoglobin by using the oxygen from the air. Page 12, lines 4-7 of the application. The "gas mixture used in the modified atmosphere packages of the present invention, after removal, allows the carboxymyoglobin to convert to oxymyoglobin and then to metmyoglobin (brown) in a natural time period." Thus, the present invention does not "fix" the color. Page 12, lines 3-5 of the application.

9. Carbon monoxide (CO) has not been allowed to be used with fresh meat in the United States for about 40 years. The Food and Drug Administration ("FDA") regulation that currently prevents using CO with meat packaging systems in the United States is 21 C.F.R. § 173.350.

10. The concern of the FDA is believed to be that CO fixes the fresh meat color to a degree that allows the retailer to sell meat that looks good (a bright red color), but is unsafe and potentially dangerous to consume because it has unacceptable levels of bacteria. This act of fixing the meat color to a bright red color is referred to as "economic adulteration."

11. After about 40 years of not allowing CO to be used with fresh meats in the United States, the Applicants came up with novel approaches of using CO in modified atmosphere packaging (MAP) systems that avoided the concerns of "fixing" the meat color.

12. Pactiv Corporation, the assignee of the present invention, then gave notice to the FDA of a specific embodiment and process and evidence supporting Pactiv's conclusion that CO as used is GRAS (generally recognized as safe). The MAP system in Pactiv's GRAS notice may be used for packaging meats such as fresh cuts of case-ready muscle meat and ground case-ready meat to maintain wholesomeness, provide flexibility in distribution, and losses due to spoilage at retail sale.




13. The specific MAP system that was presented in the GRAS notice used 0.4% CO in a meat packaging system and was referred to as Pactiv's ActiveTech™ meat packaging system. The ActiveTech™ meat packaging system traditionally includes meats being placed in polystyrene trays and covered with oxygen-permeable, polyvinyl chloride ("PVC") overwraps. The wrapped trays of meat are then placed in an outer barrier bag. Ambient air is removed and replaced with a blend of 0.4% CO, 30% carbon dioxide, and the balance being nitrogen. The myoglobin of the meat converts from oxymyoglobin to carboxymyoglobin (red). The meat maintains its red color while in storage until the package is opened for retail display by removing the outer barrier bag. The package will lose CO to the atmosphere and, thus, the retail display will not have CO. Once in retail display, the meat's myoglobin begins its natural conversion to metmyoglobin (brown). The CO used in the Pactiv MAP system did not mask the spoilage or extend the color life beyond the point of wholesomeness (*i.e.*, the point of microbial soundness).

14. The FDA stated that it had no questions regarding Pactiv's conclusion about Pactiv's ActiveTech™ meat packaging system using 0.4% CO being GRAS because of the evidence presented by Pactiv in its notice. This FDA review allows Pactiv to use CO with fresh meat in its application. It is believed to be the first system to overcome the prohibition of CO with fresh meat in the U.S. in the last 40 years.

15. I hereby declare that all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true; and, further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date: \_\_\_\_\_

9/8/03

  
Gary R. DelDuca

and S-2-(ethylthio)ethyl phosphorothioates) on the same raw agricultural commodity the total amount of such pesticides shall not yield more residue than that permitted by the larger of the two tolerances, calculated as demeton.

Section 120.105 is amended by adding thereto tolerances for residues of demeton in or on sugar beet tops and sugar beets. As amended § 120.105 reads as follows:

**§ 120.105 Tolerances for residues of demeton.**

Tolerances for residues of demeton (a mixture of O,O-diethyl O-(and S)-2-(ethylthio)ethyl phosphorothioates) are established as follows:

12 parts per million in or on alfalfa hay, clover hay.

5 parts per million in or on almond hulls, fresh alfalfa, fresh clover, sugar beet tops.

1.25 parts per million in or on grapes, hops.

0.75 part per million in or on almonds, apples, apricots, broccoli, brussels sprouts, cabbage, cauliflower, celery, cottonseed, grapefruit, lemons, lettuce, muskmelons, oranges, peaches, pears, peas, pecans, peppers, plums (fresh prunes), potatoes, strawberries, tomatoes, walnuts.

0.5 part per million in or on sugar beets.

0.3 part per million in or on beans.

B. The Commissioner of Food and Drugs, having evaluated the data submitted in a petition filed by Chemagro Corporation, P.O. Box 4913, Kansas City 20, Missouri, and other relevant material, has concluded that the following regulation should issue with respect to residues of the food additive demeton present in dehydrated sugar beet pulp. Such residues have been shown to occur from application of the pesticide to sugar beets under agricultural uses provided for by a concurrent regulation under section 408 of the act. Therefore, pursuant to the provisions of the Federal Food, Drug, and Cosmetic Act (sec. 409(c)(4), 72 Stat. 1786; 21 U.S.C. 348(c)(4)), and under the authority delegated to the Commissioner by the Secretary of Health, Education, and Welfare (25 F.R. 8625), the food additive regulations (21 CFR Part 121) are amended by adding to Subpart C the following new section:

**§ 121.221 Demeton.**

A tolerance of 5 parts per million is established for residues of demeton (a mixture of O,O-diethyl O-(and S)-2-(ethylthio)ethyl phosphorothioates) in dehydrated sugar beet pulp for livestock feed when present therein as a result of the application of the pesticide in the production of sugar beets, provided that if residues of O,O-diethyl S-2-(ethylthio)ethyl phosphorodithioate are also present, the total of both residues shall not exceed 5 parts per million.

Any person who will be adversely affected by the foregoing order may at any time prior to the thirtieth day from the date of its publication in the FEDERAL REGISTER, Clerk, De-

on, and Wel-

(are Room 5440, 330 Independence Avenue, S.W., Washington 25, D.C. written objections thereto. Objections shall show wherein the person filing will be adversely affected by the order and specify with particularity the provisions of the order deemed objectionable and the grounds for the objections. If a hearing is requested, the objections must state the issues for the hearing. A hearing will be granted if the objections are supported by grounds legally sufficient to justify the relief sought. Objections may be accompanied by a memorandum or brief in support thereof. All documents shall be filed in triplicate.

**Effective date.** This order shall be effective on the date of its publication in the FEDERAL REGISTER.

(Secs. 408(d)(2), 409(c)(4); 68 Stat. 512, 72 Stat. 1786; 21 U.S.C. 346a(d)(2), 348(c)(4))

Dated: July 26, 1961.

[SEAL] GEO. P. LARRICK,  
Commissioner of Food and Drugs.

[F.R. Doc. 61-7270; Filed, Aug. 1, 1961; 8:50 a.m.]

**PART 121—FOOD ADDITIVES**

**Subpart C—Food Additives Permitted in Animal Feed and Animal Feed Supplements**

**O,O-DIETHYL S-2-(ETHYLTHIO)ETHYL PHOSPHORODITHIOATE**

Pursuant to sections 409 and 701 of the Federal Food, Drug, and Cosmetic Act and under the authority delegated to the Commissioner of Food and Drugs by the Secretary of Health, Education, and Welfare (25 F.R. 8625), § 121.215 of the food additive regulations (26 F.R. 2595) is revised to read as follows:

**§ 121.215 O,O-Diethyl S-2-(ethylthio)ethyl phosphorodithioate.**

A tolerance of 5 parts per million is established for residues of O,O-diethyl S-2-(ethylthio)ethyl phosphorodithioate, calculated as demeton, in dehydrated sugar beet pulp for livestock feed when present therein as a result of the application of the pesticide to the growing agricultural crop, provided that, if residues of demeton are also present, the total of both residues shall not exceed 5 parts per million.

This amendment does not require notice and public procedure since it is made for the purpose of bringing § 121.215 into conformity with the pesticide regulations.

**Effective date.** This order shall be effective on the date of its publication in the FEDERAL REGISTER.

(Secs. 409, 701; 52 Stat. 1055, 72 Stat. 1785; 21 U.S.C. 348, 371)

Dated: July 26, 1961.

[SEAL] GEO. P. LARRICK,  
Commissioner of Food and Drugs.

[F.R. Doc. 61-7272; Filed, Aug. 1, 1961; 8:50 a.m.]

**PART 121—FOOD ADDITIVES**

**Subpart D—Food Additives Permitted in Food for Human Consumption**

**COMBUSTION PRODUCT GAS**

The Commissioner of Food and Drugs, having evaluated the data submitted by the Vitagen Corporation, 354 South Spring Street, Los Angeles 13, California, and other relevant material, has concluded that the following food additive regulation should issue with respect to the food additive combustion product gas used for the displacing and removal of oxygen in processing and packing of food. Therefore, pursuant to the provisions of the Federal Food, Drug, and Cosmetic Act (sec. 409(c)(1), 72 Stat. 1786; 21 U.S.C. 348(c)(1)), and under the authority delegated to the Commissioner by the Secretary of Health, Education, and Welfare (25 F.R. 8625), the food additive regulations (21 CFR 121) are amended by adding to Subpart D the following new section:

**§ 121.1060 Combustion product gas.**

The food additive combustion product gas may be safely used in the processing and packaging of the foods designated in paragraph (c) of this section for the purpose of removing and displacing oxygen in accordance with the following prescribed conditions:

(a) The food additive is manufactured by the controlled combustion in air of butane, propane, or natural gas. The combustion equipment shall be provided with an absorption-type filter capable of removing possible toxic impurities through which all gas used in the treatment of food shall pass; and with suitable controls to insure that any combustion products failing to meet the specifications provided in this section will be prevented from reaching the food being treated.

(b) The food additive meets the following specifications:

(1) Carbon monoxide content not to exceed 4.5 percent by volume.

(2) The ultraviolet absorbance in iso-octane solution in the range 255 millimicrons to 310 millimicrons not to exceed one-third of the standard reference absorbance when tested as described in paragraph (e) of this section.

(c) It is used or intended for use to displace or remove oxygen in the processing, storage, or packaging of citrus products, vegetable fats and vegetable oils, coffee, and wine.

(d) To assure safe use of the additive in addition to the other information required by the act, the label or labeling of the combustion device shall bear adequate directions for use to provide a combustion product gas that complies with the limitations prescribed in paragraph (b) of this section, including instructions to assure proper filtration.

(e) The food additive is tested for compliance with paragraph (b)(2) by the following empirical method:

**Spectrophotometric measurements.** All measurements are made in an ultraviolet spectrophotometer in optical cells of 5 centimeters in length, and in the range of 255 millimicrons to 310 millimicrons, under the same instrumental conditions. The standard reference absorbance is the absorbance at

Aug. 2, 1961  
26 FR 6918-9

O (and S)-2-(ethylthio)ethyl phosphorothioates) on the same raw agricultural commodity, the total amount of such pesticides shall not yield more residue than that permitted by the larger of the two tolerances calculated as demeton.

Section 120.105 is amended by adding thereto tolerances for residues of demeton in or on sugar beet tops and sugar beets. As amended § 120.105 reads as follows:

**§ 120.105 Tolerances for residues of demeton.**

Tolerances for residues of demeton (a mixture of O,O-diethyl O (and S)-2-(ethylthio) ethyl phosphorothioates) are established as follows:

12 parts per million in or on alfalfa hay, clover hay.

5 parts per million in or on almond hulls, fresh alfalfa, fresh clover, sugar beet tops.

1.25 parts per million in or on grapes, hops.

0.75 part per million in or on almonds, apples, apricots, broccoli, brussels sprouts, cabbage, cauliflower, celery, cottonseed, grapefruit, lemons, lettuce, muskmelons, oranges, peaches, pears, peas, pecans, peppers, plums (fresh prunes), potatoes, strawberries, tomatoes, walnuts.

0.5 part per million in or on sugar beets.

0.3 part per million in or on beans.

B. The Commissioner of Food and Drugs, having evaluated the data submitted in a petition filed by Chemagro Corporation, P.O. Box 4913, Kansas City 20, Missouri, and other relevant material, has concluded that the following regulation should issue with respect to residues of the food additive demeton present in dehydrated sugar beet pulp. Such residues have been shown to occur from application of the pesticide to sugar beets under agricultural uses provided for by a concurrent regulation under section 408 of the act. Therefore, pursuant to the provisions of the Federal Food, Drug, and Cosmetic Act (sec. 409(c)(4), 72 Stat. 1786; 21 U.S.C. 348(c)(4)), and under the authority delegated to the Commissioner by the Secretary of Health, Education, and Welfare (25 F.R. 8625), the food additive regulations (21 CFR Part 121) are amended by adding to Subpart C the following new section:

**§ 121.221 Demeton.**

A tolerance of 5 parts per million is established for residues of demeton (a mixture of O,O-diethyl O (and S)-2-(ethylthio)ethyl phosphorothioates) in dehydrated sugar beet pulp for livestock feed when present therein as a result of the application of the pesticide in the production of sugar beets, provided that if residues of O,O-diethyl S-2-(ethylthio) ethyl phosphorodithioate are also present, the total of both residues shall not exceed 5 parts per million.

Any person who will be adversely affected by the foregoing order may at any time prior to the thirtieth day from the date of its publication in the FEDERAL REGISTER file with the Hearing Clerk, Department of Health, Education, and Wel-

fare Room 5440 380 Independence Avenue SW., Washington 25, D.C., written objections thereto. Objections shall show wherein the person filing will be adversely affected by the order and specify with particularity the provisions of the order deemed objectionable and the grounds for the objections. If a hearing is requested, the objections must state the issues for the hearing. A hearing will be granted if the objections are supported by grounds legally sufficient to justify the relief sought. Objections may be accompanied by a memorandum or brief in support thereof. All documents shall be filed in triplicate.

**Effective date.** This order shall be effective on the date of its publication in the FEDERAL REGISTER.

(Secs. 408(d)(2), 409(c)(4); 68 Stat. 512, 72 Stat. 1786; 21 U.S.C. 346a(d)(2), 348(c)(4))

Dated: July 26, 1961.

[SEAL] GEO. P. LARRICK,  
Commissioner of Food and Drugs.

[F.R. Doc. 61-7270; Filed, Aug. 1, 1961; 8:50 a.m.]

**PART 121—FOOD ADDITIVES**

**Subpart C—Food Additives Permitted in Animal Feed and Animal Feed Supplements**

**O,O-DIETHYL S-2-(ETHYLTHIO)ETHYL PHOSPHORODITHIOATE**

Pursuant to sections 409 and 701 of the Federal Food, Drug, and Cosmetic Act and under the authority delegated to the Commissioner of Food and Drugs by the Secretary of Health, Education, and Welfare (25 F.R. 8625), § 121.215 of the food additive regulations (26 F.R. 2595) is revised to read as follows:

**§ 121.215 O,O-Diethyl S-2-(ethylthio) ethylphosphorodithioate.**

A tolerance of 5 parts per million is established for residues of O,O-diethyl S-2-(ethylthio)ethyl phosphorodithioate, calculated as demeton, in dehydrated sugar beet pulp for livestock feed when present therein as a result of the application of the pesticide to the growing agricultural crop, provided that, if residues of demeton are also present, the total of both residues shall not exceed 5 parts per million.

This amendment does not require notice and public procedure since it is made for the purpose of bringing § 121.215 into conformity with the pesticide regulations.

**Effective date.** This order shall be effective on the date of its publication in the FEDERAL REGISTER.

(Secs. 409, 701; 52 Stat. 1055, 72 Stat. 1785; 21 U.S.C. 348, 371)

Dated: July 26, 1961.

[SEAL] GEO. P. LARRICK,  
Commissioner of Food and Drugs.

[F.R. Doc. 61-7272; Filed, Aug. 1, 1961; 8:50 a.m.]

**PART 121—FOOD ADDITIVES**

**Subpart D—Food Additives Permitted in Food for Human Consumption**

**COMBUSTION PRODUCT GAS**

The Commissioner of Food and Drugs, having evaluated the data submitted by the Vitagen Corporation, 354 South Spring Street, Los Angeles 13, California, and other relevant material, has concluded that the following food additive regulation should issue with respect to the food additive combustion product gas used for the displacing and removal of oxygen in processing and packing of food. Therefore, pursuant to the provisions of the Federal Food, Drug, and Cosmetic Act (sec. 409(c)(1), 72 Stat. 1786; 21 U.S.C. 348(c)(1)), and under the authority delegated to the Commissioner by the Secretary of Health, Education, and Welfare (25 F.R. 8625), the food additive regulations (21 CFR 121) are amended by adding to Subpart D the following new section:

**§ 121.1060 Combustion product gas.**

The food additive combustion product gas may be safely used in the processing and packaging of the foods designated in paragraph (c) of this section for the purpose of removing and displacing oxygen in accordance with the following prescribed conditions:

(a) The food additive is manufactured by the controlled combustion in air of butane, propane, or natural gas. The combustion equipment shall be provided with an absorption-type filter capable of removing possible toxic impurities through which all gas used in the treatment of food shall pass; and with suitable controls to insure that any combustion products failing to meet the specifications provided in this section will be prevented from reaching the food being treated.

(b) The food additive meets the following specifications:

(1) Carbon monoxide content not to exceed 4.5 percent by volume.

(2) The ultraviolet absorbance in iso-octane solution in the range 255 millimicrons to 310 millimicrons not to exceed one-third of the standard reference absorbance when tested as described in paragraph (e) of this section.

(c) It is used or intended for use to displace or remove oxygen in the processing, storage, or packaging of citrus products, vegetable fats and vegetable oils, coffee, and wine.

(d) To assure safe use of the additive in addition to the other information required by the act, the label or labeling of the combustion device shall bear adequate directions for use to provide combustion product gas that complies with the limitations prescribed in paragraph (b) of this section, including instructions to assure proper filtration.

(e) The food additive is tested for compliance with paragraph (b)(2) by the following empirical method:

**Spectrophotometric measurements.** All measurements are made in an ultraviolet spectrophotometer in optical cells of 5 centimeters in length, and in the range of 255 millimicrons to 310 millimicrons, under the same instrumental conditions. The standard reference absorbance is the absorbance



with particularity the grounds of the order deemed objectionable and the grounds for the objections. If a hearing is requested, the objections must state the issues for the hearing. A hearing will be granted if the objections are supported by grounds legally sufficient to justify the relief sought. Objections may be accompanied by a memorandum or brief in support thereof. All documents shall be filed in quintuplicate.

**Effective date.** This order shall be effective on the date of its publication in the *FEDERAL REGISTER*.

(Sec. 409(c) (1), 72 Stat. 1786; 21 U.S.C. 348(c) (1))

Dated: December 7, 1962.

GEO. P. LARRICK,  
*Commissioner of Food and Drugs.*

[F.R. Doc. 62-12361; Filed, Dec. 13, 1962; 8:46 a.m.]

## PART 121—FOOD ADDITIVES

### Subpart D—Food Additives Permitted in Food for Human Consumption

#### COMBUSTION PRODUCT GAS

The Commissioner of Food and Drugs, having evaluated the data submitted in petitions filed by the Whirlpool Corporation, Benton Harbor, Michigan, and the Vitagen Corporation, 1263 Westwood Boulevard, Los Angeles, California, and other relevant material, has concluded that the food additive regulation with respect to combustion product gas should be amended as set forth below. Therefore, pursuant to the provisions of the Federal Food, Drug, and Cosmetic Act (sec. 409(c) (1), 72 Stat. 1786; 21 U.S.C. 348(c) (1)), and under the authority delegated to the Commissioner by the Secretary of Health, Education, and Welfare (25 F.R. 8625), § 121.1060(c) (21 CFR 121.1060; 27 F.R. 4014) is amended to read as follows:

#### § 121.1060 Combustion product gas.

(c) It is used or intended for use to displace or remove oxygen in the processing, storage, or packaging of beverage products and other food, except fresh meats.

Any person who will be adversely affected by the foregoing order may at any time within 30 days from the date of its publication in the *FEDERAL REGISTER* file with the Hearing Clerk, Department of Health, Education, and Welfare, Room 5440, 330 Independence Avenue SW, Washington 25, D.C., written objections thereto. Objections shall show wherein the person filing will be adversely affected by the order and specify with particularity the provisions of the order deemed objectionable and the grounds for the objections. If a hearing is requested, the objections must state the issues for the hearing. A hearing will be granted if the objections are supported by grounds legally sufficient to justify the relief sought. Objections may be accompanied by a memorandum or brief in support thereof. All documents shall be filed in quintuplicate.

**Effective date.** This order shall be effective on the date of its publication in the *FEDERAL REGISTER*.

(Sec. 409(c) (1), 72 Stat. 1786; 21 U.S.C. 348(c) (1))

Dated: December 7, 1962.

GEO. P. LARRICK,  
*Commissioner of Food and Drugs.*  
[F.R. Doc. 62-12360; Filed, Dec. 13, 1962; 8:46 a.m.]

## Title 39—POSTAL SERVICE

### Chapter I—Post Office Department

#### PART 168—DIRECTORY OF INTERNATIONAL MAIL

##### Individual Country Amendments

The regulations of the Post Office Department in § 168.5 *Individual country regulations* are amended as follows:

I. In country "Bolivia", under Parcel Post, amend the item "Prohibitions" to read as follows:

**Prohibitions.** Firearms, daggers, black-jacks, brass knuckles, sidearms and concealable weapons.

Cigarette lighters.

Gambling devices.

Pharmaceutical and medicinal products, unless approved by the Bolivian health authorities. In case of doubt, senders should ascertain from the addressees in advance of mailing whether the medicine they desire to send will be admitted.

Articles which violate the Bolivian trademark laws.

Counterfeit or illegal currency; advertisements imitating currency or postage stamps, except for philatelic or numismatic catalogs.

Adulterated or harmful beverages or foodstuffs.

II. In country "Canada", as amended by 27 F.R. 404, 27 F.R. 10369, under Parcel Post, the item "Prohibitions" is amended by revising the sixth paragraph to include "Plumage and skins of wild birds" and by adding a new paragraph at the end thereof to prescribe regulations for importing meat. As so amended, paragraph six and the new paragraph read as follows:

**Prohibitions.** . . .

Commercial tags of metal. Prison-made goods being sold or intended for sale by a person or firm. Plumage and skins of wild birds.

Meat and meat food products, unless federally inspected and passed and marked accordingly. If intended for sale, export certification by the United States Department of Agriculture is also required. Meat or meat food product for personal use is exempt from export certification, but the addressee is required to certify to the Canadian authorities that it will not be offered for sale in Canada.

III. In country "Japan", under Parcel Post, the item "Prohibitions" is amended by revising the second paragraph to in-

clude wool samples among animal products. As so amended, the second paragraph reads as follows:

**Prohibitions.** . . .

The following must be accompanied by official inspection certificates showing that they are free from domestic animals' infectious disease: Meat, bones, skin, hair, feathers, horns or hoofs of hoofed animals, rabbits, or poultry, wool samples, poultry eggs for hatching; honey bees.

IV. In country "Kenya and Uganda", as amended by 27 F.R. 3738, 27 F.R. 5659, under Parcel Post, amend the tabular information immediately following the item "Air parcel rates" by striking out "Weight limit: 11 pounds", and inserting in lieu thereof "Weight limit: 22 pounds."

V. In country "Laos", as amended by 27 F.R. 8592, amend the item "Observations" where it appears both under Postal Union Mail and Parcel Post, to respectively read as follows:

**Observations.** The following are the only post offices in operation:

Vientiane.	Paksé.
Honeisal.	Paksong.
Luangprabang.	Khongsedoné.
Sayaboury.	Champassak.
Paksane.	Muong Kong.
Khammouane.	Saravane.
Savannakhet.	Attopeu.

**Observations.** See the item "Observations" under Postal Union Mail for post offices which are in operation.

VI. In country "Tanganyika Territory" under Parcel Post, amend the tabular information immediately following the item "Air parcel rates" by striking out "Weight limits: 11 pounds" and inserting in lieu thereof "Weight limits: 22 pounds".

VII. In country "Thailand", as amended by 27 F.R. 7022, under Parcel Post, make the following changes to show that insured parcel post service is available.

A. Amend the tabular information immediately following the item "Air parcel rates" to read as follows:

Weight limit: 22 pounds
Sealing: Insured parcels must, and ordinary parcels may be sealed
Registration: No
Insurance: Yes
Postal forms required:
1 Form 2922
1 Form 2966

B. Strike out the item "Indemnity, No provision." and insert in lieu thereof the following:

**Insurance.** The following insurance fees and limits of indemnity apply:

Limit of indemnity:	Fee, cents
Not over \$10.....	20
From \$10.01 to \$25.....	25
From \$25.01 to \$50.....	35
From \$50.01 to \$100.....	55

Insured parcels may only be addressed to Bangkok or Dhonburi.

Print on the wrapper, near the "INSURED" endorsement and number, the amount for which the parcel is insured. This amount shall be shown in United

§ 173.350

other information required by the act, the following:

- (i) The name of the additive, chloropentafluoroethane.
  - (ii) The percentage of the additive present in the case of a mixture.
  - (iii) The designation "food grade".
- (2) The label or labeling of the food additive container shall bear adequate directions for use.

[42 FR 14526, Mar. 15, 1977, as amended at 43 FR 11317, Mar. 17, 1978; 43 FR 14644, Apr. 7, 1978]

§ 173.350 Combustion product gas.

The food additive combustion product gas may be safely used in the processing and packaging of the foods designated in paragraph (c) of this section for the purpose of removing and displacing oxygen in accordance with the following prescribed conditions:

(a) The food additive is manufactured by the controlled combustion in air of butane, propane, or natural gas. The combustion equipment shall be provided with an absorption-type filter capable of removing possible toxic impurities, through which all gas used in the treatment of food shall pass; and with suitable controls to insure that any combustion products failing to meet the specifications provided in this section will be prevented from reaching the food being treated.

(b) The food additive meets the following specifications:

(1) Carbon monoxide content not to exceed 4.5 percent by volume.

(2) The ultraviolet absorbance in isooctane solution in the range 255 millimicrons to 310 millimicrons not to exceed one-third of the standard reference absorbance when tested as described in paragraph (e) of this section.

(c) It is used or intended for use to displace or remove oxygen in the processing, storage, or packaging of beverage products and other food, except fresh meats.

(d) To assure safe use of the additive in addition to the other information required by the act, the label or labeling of the combustion device shall bear adequate directions for use to provide a combustion product gas that complies with the limitations prescribed in paragraph (b) of this section, including instructions to assure proper filtration.

21 CFR Ch. I (4-1-03 Edition)

(e) The food additive is tested for compliance with paragraph (b)(2) by the following empirical method:

*Spectrophotometric measurements.* All measurements are made in an ultraviolet spectrophotometer in optical cells of 5 centimeters in length, and in the range of 255 millimicrons to 310 millimicrons, under the same instrumental conditions. The standard reference absorbance is the absorbance at 275 millimicrons of a standard reference solution of naphthalene (National Bureau of Standards Material No. 577 or equivalent in purity) containing a concentration of 1.4 milligrams per liter in purified isooctane, measured against isooctane of the same spectral purity in 5-centimeter cells. (This absorbance will be approximately 0.30.)

*Solvent.* The solvent used is pure grade isooctane having an ultraviolet absorbance not to exceed 0.05 measured against distilled water as a reference. Upon passage of purified inert gas through some isooctane under the identical conditions of the test, a lowering of the absorbance value has been observed. The absorbance of isooctane to be used in this procedure shall not be more than 0.02 lower in the range 255 millimicrons to 310 millimicrons, inclusive, than that of the untreated solvent as measured in a 5-centimeter cell. If necessary to obtain the prescribed purities, the isooctane may be passed through activated silica gel.

*Apparatus.* To assure reproducible results, the additive is passed into the isooctane solution through a gas-absorption train consisting of the following components and necessary connections:

1. A gas flow meter with a range up to 30 liters per hour provided with a constant differential relay or other device to maintain a constant flow rate independent of the input pressure.

2. An absorption apparatus consisting of an inlet gas dispersion tube inserted to the bottom of a covered cylindrical vessel with a suitable outlet on the vessel for effluent gas. The dimensions and arrangement of tube and vessel are such that the inlet tube introduces the gas at a point not above 5/4 inches below the surface of the solvent through a sintered glass outlet. The dimensions of the vessel are such, and both inlet and vessel are so designed, that the gas can be bubbled through 60 milliliters of isooctane solvent at a rate up to 30 liters per hour without mechanical loss of solvent. The level corresponding to 60 milliliters should be marked on the vessel.

3. A cooling bath containing crushed ice and water to permit immersion of the absorption vessel at least to the solvent level mark.

*Caution.* The various parts of the absorption train must be connected by gas-tight tubing and joints composed of materials which will neither remove components from

nor add components to the gas stream. The gas source is connected in series to the flow-rate device, the flow meter, and the absorption apparatus in that order. Ventilation should be provided for the effluent gases which may contain carbon monoxide.

**Sampling procedure.** Immerse the gas-absorption apparatus containing 60 milliliters of isooctane in the coolant bath so that the solvent is completely immersed. Cool for at least 15 minutes and then pass 120 liters of the test gas through the absorption train at a rate of 30 liters per hour or less. Maintain the coolant bath at 0 °C throughout. Remove the absorption vessel from the bath, disconnect, and warm to room temperature. Add isooctane to bring the contents of the absorption vessel to 60 milliliters, and mix. Determine the absorbance of the solution in the 5-centimeter cell in the range 255 millimicrons to 310 millimicrons, inclusive, compared to isooctane. The absorbance of the solution of combustion product gas shall not exceed that of the isooctane solvent at any wavelength in the specified range by more than one-third of the standard reference absorbance.

#### § 173.355 Dichlorodifluoromethane.

The food additive dichlorodifluoromethane may be safely used in food in accordance with the following prescribed conditions:

- (a) The additive has a purity of not less than 99.97 percent.

(b) It is used or intended for use, in accordance with good manufacturing practice, as a direct-contact freezing agent for foods.

(c) To assure safe use of the additive:

(1) The label of its container shall bear, in addition to the other information required by the act, the following:

(i) The name of the additive, dichlorodifluoromethane, with or without the parenthetical name "Food Freezant 12".

(ii) The designation "food grade".

(2) The label or labeling of the food additive container shall bear adequate directions for use.

#### § 173.357 Materials used as fixing agents in the immobilization of enzyme preparations.

Fixing agents may be safely used in the immobilization of enzyme preparations in accordance with the following conditions:

(a) The materials consist of one or more of the following:

(1) Substances generally recognized as safe in food.

(2) Substances identified in this subparagraph and subject to such limitations as are provided:

Substances	Limitations
Acrylamide-acrylic acid resin: Complying with § 173.5(a)(1) and (b) of this chapter.	May be used as a fixing material in the immobilization of glucose isomerase enzyme preparations for use in the manufacture of high fructose corn syrup, in accordance with § 184.1372 of this chapter.
Cellulose triacetate .....	May be used as a fixing material in the immobilization of lactase for use in reducing the lactose content of milk.
Diethylaminoethyl-cellulose .....	May be used as a fixing material in the immobilization of glucose isomerase enzyme preparations for use in the manufacture of high fructose corn syrup, in accordance with § 184.1372 of this chapter.
Dimethylamine-epichlorohydrin resin: Complying with § 173.60(a) and (b) of this chapter.	May be used as a fixing material in the immobilization of glucose isomerase enzyme preparations for use in the manufacture of high fructose corn syrup, in accordance with § 184.1372 of this chapter.
Glutaraldehyde .....	Do.
Periodic acid (CAS Reg. No. 10450-60-9) ..	

~~MEMORANDUM OF CONFERENCE~~

May 22, 1962

BETWEEN: Mr. Donald W. Thomas, Legal Counsel, The Whirlpool Corporation  
Benton Harbor, Michigan

and

Mr. A. T. Spiher, Jr., Food Additive Petitions Control Branch

SUBJECT: Combustion product gas.  
Food Additive Petition 751.

Mr. Thomas called without previous appointment to discuss the above petition. He said that he had received my letter of May 10, 1962, in which we filed the petition, and said that we may need additional data on meat. These data would be needed to establish that the treatment of meat would not serve to cause the meat to retain its fresh red color longer than meat not so treated.

I explained to Mr. Thomas the way in which petitions are handled, and explained the question which we have concerning possible deception of the consumer where treatment of the meat leads to longer retention of the fresh red color. I said that they could either submit additional data on this point or they could request withdrawal of the portion of the petition for meat, and explained the different courses of action.

Mr. Thomas said that they had data concerning the retention of red color in meat, and they will get it together. He was concerned, however, about whether he should submit this as an amendment which would start the time clock over, or should withdraw animal products from the petition, to submit later on.

I said that this was a decision which he would have to make in the light of the explanation we had given him, and I suggested that he submit the data which they have and let us look at it before they did anything additional, because what they had done might be sufficient for our people.

I further suggested that when he submit the information for meat, he should supplement the data in the petition to explain exactly how the combustion product gas is to be used on the various commodities named in their petition. He said that he would do so. Briefly, he said that the gas was to be used as the atmosphere in a cold storage room.

In response to a question, he said that they had tested the effluent from their generator and were satisfied that the gas complied with the requirements established in the food additive regulation.

cc: FA:DF:DP:EE:DOH:BPS  
ATSpiher:nrg:5/23/62  
R/D:ATS:nrg:5/24/62



ADMINISTRATIVE CENTER • BENTON HARBOR, MICHIGAN

July 23, 1962

Mr. Alan T. Spiher, Jr.  
Food and Drug Administration  
Department of Health, Education and Welfare  
Washington 25, D. C.

Subject: Food Additive Petition No. 751

7A 3/9/62

Dear Mr. Spiher:

We are in receipt of your letter of May 10, 1962, advising us of the filing of Food Additive Petition No. 751 with an effective filing date of March 24, 1962.

In view of your comments in the above-mentioned letter, we now request that our petition as originally presented be amended so as to delete any reference to animal products wherein paragraph 121.1060, section (c) of Part 121, Sub-Part D of Title 21 would now read as follows:

- (c) It is used or intended for use to displace or remove oxygen in the processing, storage, or packaging of citrus products, vegetable fats and vegetable oils, coffee, wine, fruit and fruit products and vegetable and vegetable products.

The following comments are submitted to further supplement the Remarks section of our first letter of March 6, 1962.

In food studies conducted at the Whirlpool Research Laboratories involving the use of combustion product gas as set forth in paragraph 121.1060 of Title 21, fruits and vegetables were stored under refrigeration at temperatures between

- 10 copies sent  
DF DP  
7/13/62

12A

FAPB

JUL 25 1962



32° and 45° F. and in their normal distribution containers, that is, baskets, crates and boxes. Products so stored had a shelf life of from three to five times that of air-stored food held at the same temperature. The results of one such study involving apples stored in air versus apples stored in conventional controlled atmosphere versus apples stored in combustion product gas are presented in the attached table. It will be noted that apples stored in combustion product gas had firmer flesh and a lower incidence of scald than did apples stored either in air or conventional controlled atmosphere even though the apples in combustion product gas were in storage for a longer period of time.

The combustion product gas under study at Whirlpool would most likely be used in the following general areas:

1. Fresh fruit and vegetable storage
2. Processors - storage, packaging and processing
3. Transportation

Because of these diverse applications, our petition requests approval for fruit and vegetable "products" as well as the natural, original raw fruits and vegetables.

To expand on the use of combustion product gas by food processors, the following examples are presented:

1. Storage of fruits and vegetables in order to have better quality control, improve yield and extend packaging season.
2. Packaging of processed foods in inert gases, i. e., nitrogen and/or carbon dioxide to prevent oxidative changes that may develop during storage.
3. Use of gas mixtures in certain processing steps as a "blanket" to keep out oxygen and prevent the associated undesirable changes.


Mr. Alan T. Spiher, Jr.

Page Three

We are hopeful that the requested amendment to the petition as well as the supplemental information presented above will clear up any questions concerning Food Additive Petition No. 751 and that favorable action will be shortly forthcoming.

Very truly yours,

WHIRLPOOL CORPORATION

  
By John E. Mahaffey  
Vice President

ERIC F. GREENBERG  
ATTORNEY AT LAW

ERIC F. GREENBERG  
Of Counsel  
Ungaretti & Harris

*Contains Confidential Business Information*

August 29, 2001

Division of GRAS Notice Review  
Office of Food Additive Safety  
Center for Food Safety and Applied Nutrition  
Food and Drug Administration  
200 C St., SW  
Washington, DC 20204

Re: NOTIFICATION OF CLAIM FOR GENERAL RECOGNITION OF  
SAFETY OF CARBON MONOXIDE IN A MODIFIED ATMOSPHERE  
SYSTEM FOR PACKAGING FRESH MEAT, submitted by Pactiv  
Corporation

To the FDA:


Enclosed is the NOTIFICATION OF CLAIM FOR GENERAL RECOGNITION OF  
SAFETY OF CARBON MONOXIDE IN A MODIFIED ATMOSPHERE SYSTEM FOR  
PACKAGING FRESH MEAT, submitted by Pactiv Corporation, 1900 West Field Court,  
Lake Forest, Illinois 60045, c/o the undersigned counsel, consisting of pages 000001.001  
through 000250.

Please note that this submission contains Confidential Business Information that  
Pactiv Corporation desires not to be revealed to Freedom of Information Act  
requestors and other members of the public. In the first copy of the submission  
following this letter, the Confidential Business Information has been redacted. For  
ease of reference, a list indicating which pages contain redactions is attached.

Five complete copies of the submission are enclosed, including the one that has been  
redacted.

If you have any questions, please contact me at 312 977-4647.

Sincerely,

  
Eric F. Greenberg  
Enc.

000001.001

REDACTIONS

000003  
000005  
000006  
000008  
000009  
000013  
000016  
000017  
000018  
000019

000020  
000026  
000027  
000028  
000033  
000038  
000040  
000041  
000043  
000054

000056  
000057  
000160  
000162  
000166  
000199  
000222  
000248

Notification of Claim for General Recognition of Safety of Carbon Monoxide in A  
Modified Atmosphere System for Packaging Fresh Meat

TABLE OF CONTENTS

	<u>Page</u>
GRAS Notice .....	000003
I. Claim of Exemption .....	000004
a. Name and address of the notifier.....	000004
b. Common or unusual name of the notified Substance .....	000004
c. Conditions of use (foods, levels, purposes).....	000005
Summary .....	000005
"Traditional" ActiveTech .....	000006
AT2001 .....	000007
d. Basis of GRAS determination: Scientific procedures .....	000011
<u>CO Safety</u> .....	000011
<i>Effects on Fresh Meat and Consumption</i> .....	000015
<u>Safety: Effects on carboxymyoglobin levels</u> .....	000016
<u>Safety: The Norwegian experience</u> .....	000025
<u>Safety: Exposure in environment</u> .....	000026
<i>Corroborative information about AT2001</i> .....	000029
e. Statement of availability of information .....	000038
II. Identity of notified substance .....	000038
III. Self-limiting levels of use .....	000040
IV. Basis of GRAS determination .....	000040
Conclusion .....	000043

**Attachments:**

Attachment 1	Authorization letter from Pactiv Corporation for representation by Eric F. Greenberg . . . . .	000050
Attachment 2	Photographs of meats treated with ActiveTech and ActiveTech 2001. . . . .	000053
Attachment 3	"Application for Assessment of the Food Additive Carbon Monoxide (CO) Prior to its Authorization", Norwegian Meat Cooperative; Norwegian Independent Meat Association (1999) . . . . .	000058
Attachment 4	"Evaluation of Beef Steaks and Ground Beef in the Pactiv ActiveTech Packaging System: Effects of Carbon Monoxide in the Package Atmosphere", Hachmeister, K; Hunt, M.; Milliken, G; May 2001 . . . . .	000157
Attachment 5	Certificate of Conformance, Carbon monoxide, Haun Welding Supply, Inc., Syracuse, New York, May 8, 2001. . . . .	000190
Attachment 6	"Standard Practice for Analysis of Reformed Gas by Gas Chromatography", ASTM D 1946-90 (Reapproved 2000) . . . . .	000192
Attachment 7	Daren Cornforth, Ph.D., letter and curriculum vitae . . . . .	000198
Attachment 8	Vasilios H. Frankos, Ph.D., letter and curriculum vitae . . . . .	000205
Attachment 9	Melvin C. Hunt, Ph.D., letter and curriculum vitae . . . . .	000221
Attachment 10	Oddvin Sorheim, Ph.D., letter and curriculum vitae . . . . .	000247

**ERIC F. GREENBERG**  
**ATTORNEY AT LAW**

ERIC F. GREENBERG  
Of Counsel  
Ungaretti & Harris

**Contains Confidential Business Information**

August 29, 2001

Division of GRAS Notice Review  
Office of Food Additive Safety  
Center for Food Safety and Applied Nutrition  
Food and Drug Administration  
200 C St, SW  
Washington, DC 20204

**Re: NOTIFICATION OF CLAIM FOR GENERAL  
RECOGNITION OF SAFETY OF CARBON MONOXIDE  
IN A MODIFIED ATMOSPHERE SYSTEM FOR  
PACKAGING FRESH MEAT**

To the FDA:

This letter and its attachments contains the notification, pursuant to the Federal Food, Drug and Cosmetic Act and FDA's regulations, by Pactiv Corporation, 1900 West Field Court, Lake Forest, Illinois 60045, c/o attorney Eric F. Greenberg, 3500 Three First National Plaza, Chicago, Illinois 60602<sup>1</sup>, for the General Recognition of Safety of carbon monoxide ("CO") at a level of 0.4% in a modified atmosphere system for packaging fresh meat.

---

<sup>1</sup> Attachment 1 contains Pactiv's authorization of undersigned counsel, as well as a Summary regarding Pactiv Corporation.

[efgreenberg@uhlalaw.com](mailto:efgreenberg@uhlalaw.com)

3500 THREE FIRST NATIONAL PLAZA, CHICAGO, ILLINOIS 60602-4283  
PHONE: 312-977-4647 · FAX: 312-977-4405

000003

As set forth more fully below and in the attachments to this document, Pactiv believes its intended use of CO is GRAS based on scientific procedures within the meaning of 21 U.S.C. Sec. 201(s) and FDA's implementing regulations in 21 CFR Sec. 170.30, and including FDA's proposed rule published on April 17, 1997 (62 FR 18937). FDA regulations provide that the scientific evidence available and reviewed for a GRAS determination is of the same quantity and quality as that required for a food additive approval, and that the scientific evidence of safety be generally known and accepted by qualified experts in the appropriate scientific and technical fields. 21 CFR Sec. 170.30(a).

**I. Claim of Exemption**

**a. Name and address of the notifier.**

Pactiv Corporation  
1900 West Field Court  
Lake Forest, Illinois 60045  
c/o Eric F. Greenberg  
Attorney at Law  
3500 Three First National Plaza  
Chicago, IL 60602

**b. Common or usual name of the notified substance.**

Carbon monoxide ("CO")

000004



**c. Conditions of use (foods, levels, purposes).**

When used as described in this Notice, CO meeting appropriate purity specifications is a processing aid in packaging of fresh cuts of muscle meat and ground meat, as a component of a gas mixture utilized in a specific modified atmosphere packaging system. 21 CFR Sec. 170.3(o)(24). A technology utilizing 0.4% CO within a modified atmosphere packaging system will maintain wholesomeness, permit greater flexibility in distribution, and reduce shrinkage, all within a system that results in traditional product display to consumers. All elements of the system, excluding the CO, are already in use in the United States as part of a modified atmosphere meat packaging system called ActiveTech™. Notifier refers to the new system incorporating CO as "AT2001".

**Summary**

ActiveTech™ is a system that is designed to permit more extended storage of meats, but, as explained below, has no effects on retail display time or characteristics as compared with other modified atmosphere technologies currently in use. It employs materials that are either approved additives used consistently with their approvals, or GRAS substances. AT2001 adapts that system for additional storage scenarios. AT2001 serves to reduce the time needed for enzymatic reduction after modified

000005

atmosphere packaging, and allows consistent display color of whole muscle meats. AT2001's advantages are in the resulting flexibility and consistency during storage and distribution.

The GRAS use of CO described in this Notice involves use as a component of the flush gas mixture used in replacement of ambient air in the packaging for distribution of refrigerated fresh red meat. The meats are in all instances fresh, and are intended to be cooked prior to consumption.

#### "Traditional" ActiveTech™

The ActiveTech™ modified atmosphere system, in commercial use in the United States since 1998, is a modified atmosphere system for packaging fresh cuts of muscle meat, or portions of ground meat. AT2001 is a refinement of ActiveTech™, and differs from it only in the addition of 0.4% CO to the modified atmosphere.

In the "traditional" ActiveTech™ system, the meats are placed in polystyrene trays and covered with oxygen-permeable, flexible polyvinyl chloride ("PVC") overwraps. The wrapped trays of meat are then placed within an outer barrier bag from which ambient air is removed and replaced with a blend of 30% carbon dioxide (CO<sub>2</sub>) and 70% nitrogen (N<sub>2</sub>). An activated oxygen-absorbing sachet is also added within the outer bag.

This modified atmosphere maintains the packaged meat in an oxygen-free deoxymyoglobin state, with its distinctive purplish appearance that is generally considered undesirable by consumers. The traditional ActiveTech™ system relies on the rapid reduction of the oxygen content of the outer bag. Once the oxygen is removed, a "seasoning" phase begins during which enzymatic effects take place so that the meat will be able to "re-bloom" when once again in the presence of oxygen. As the residual oxygen in the package is consumed by the activated oxygen scavenger, red meat oxymyoglobin is first subject to rapid conversion to metmyoglobin (brown) at very low partial pressures of oxygen, e.g. 0.5% oxygen. This low partial pressure region of oxygen is necessarily passed through prior to ultimately reaching 0% in the package and the conversion to deoxymyoglobin (purple). This seasoning phase can take up to 5 days.

Before display to consumers at retail, the outer bag, and thus the modified atmosphere, is removed, and the traditionally wrapped product (in polystyrene foam tray with PVC overwrap) is permitted to "re-bloom" to its familiar appearance through creation of oxymyoglobin on the meat's surface.

000007

## AT2001

In the AT2001 modified atmosphere system, as with traditional ActiveTech™, fresh cuts of muscle meat, or portions of ground meat, are placed in polystyrene trays and covered with oxygen permeable flexible PVC overwraps. The wrapped trays are placed within the outer barrier bag, the air is removed and replaced with a blend of 0.4% CO, 30% carbon dioxide (CO<sub>2</sub>) and the balance nitrogen (N<sub>2</sub>). As with the traditional AT system, an activated oxygen-absorbing sachet is added within the outer bag to create and maintain an oxygen-free environment for the packaged meat during storage.

As noted, meat packaged in traditional ActiveTech™ undergoes a myoglobin pigment conversion from oxymyoglobin (red) to metmyoglobin (brown) to deoxymyoglobin (purple) in the oxygen free environment. The metmyoglobin formed generally will convert to deoxymyoglobin in the oxygen free storage environment in about 5 days, a period of time referred to as the "seasoning period". However, the meat's ability to convert all of the metmyoglobin formed to deoxymyoglobin during the seasoning period and then fully rebloom to oxymyoglobin upon re-exposure to normal, oxygen-rich atmosphere at retail, is a function of a multitude of unpredictable, uncontrollable factors in the meat such as age, muscle

000008

location, and enzyme energy level. This is a key weakness of all current low oxygen packaging systems.

Meat packaged in the AT2001 atmosphere will instead convert from oxymyoglobin to carboxymyoglobin (red) in the package due to the inclusion of 0.4% CO in the modified atmosphere. This conversion occurs during the initial 24 hours as the free oxygen in the headspace is consumed. Thus, the CO effectively protects the myoglobin from converting to metmyoglobin as the oxygen in the package is removed. This feature is especially important for the most pigment sensitive meats such as those from the round. The meat will continue to maintain its red color while in storage until the package is opened for retail display, when the outer bag (and modified atmosphere) is removed. Since carboxymyoglobin and oxymyoglobin are essentially the same colors, no seasoning period is necessary. The meat can be opened for retail display the day following packaging.

Once in retail display, the meat's myoglobin will begin the rather slow, natural conversion to metmyoglobin (brown), akin to that seen with untreated meat, allowing for a conventional retail display life of 3 to 4 days, closer to consumers' expectations of color than results from use of high

000009

oxygen packaging systems. Attachment 2 consists of photographs depicting the ActiveTech™ and AT2001 systems.

The AT2001 formulation will assure that the meat will have the familiar color during and following storage, eliminating the seasoning period, allowing for placement in retail display beginning at 1 day, and up to 30-40 days, after packing. For cuts of meat from the round, and other color sensitive cuts, the AT2001 will help them have a more uniform red color for retail display.

In AT2001 (as in traditional ActiveTech™ ), the trays and films utilized are made from familiar, FDA-approved polymers that are used in accordance with their existing approvals or GRAS status. The activated oxygen-absorbing sachet inserted into the outer bag to absorb oxygen does not contact food and is not expected to become a component of the food. Therefore, it is not a food additive under the definition in 21 USC Sec. 321(s). As an added assurance of safety, each of the sachet's components has some GRAS status or food-related approvals.

Thus, the AT2001 system adds a refinement to the existing ActiveTech™ system that will allow its utilization with whole and ground meat products that meet the processors' desire to get to market as soon as the day following processing.

000010

d. Basis of GRAS determination: Scientific procedures

CO safety

Pactiv believes its proposed use of CO is GRAS based on scientific procedures within the meaning of 21 U.S.C. Sec. 201(s) and FDA's implementing regulations in 21 CFR Sec. 170.30 and including FDA's proposed rule published on April 17, 1997 (62 FR 18937).

CO is a colorless, odorless gas that is poisonous to humans if inhaled at much higher levels than are involved with the use that is the subject of this Notice. It is formed when carbon is not completely burned, for example, in the combustion of fuels.

It is well known that CO creates negative health effects at elevated levels because it out-competes oxygen for attachment to the hemoglobin molecule. The resulting carboxyhemoglobin levels in the blood are associated with severe health effects. In addition, the equilibrium rate for the exchange of carboxyhemoglobin for oxyhemoglobin is very slow, and the resulting level of carboxyhemoglobin is a function of the CO level in the respired air, the time of exposure and the level of activity of the individual. Typical atmospheric CO levels are  $<20 \text{ mg/m}^3$  as an 8 hour mean (higher in

urban and high traffic areas), and typical carboxyhemoglobin levels due to natural background CO range between 1.2 and 1.5%.

CO is recognized as a significant air pollutant at higher levels.

Automobile exhausts, industrial processes and boilers and incinerators all contribute to air quantities of CO. According to the U.S. EPA Office of Air and Radiation:

Carbon monoxide enters the blood stream and reduces oxygen delivery to the body's organs and tissues. The health threat from exposure to CO is most serious for those who suffer from cardiovascular disease. Healthy individuals are also affected, but only at higher levels of exposure. Exposure to elevated CO levels is associated with visual impairment, reduced work capacity, reduced manual dexterity, poor learning ability, and difficulty in performing complex tasks. EPA's health-based national air quality standard for CO is 9 parts per million (ppm) [10 mg/m<sup>3</sup>] measured as an annual second-maximum 8-hour average concentration.

"Summary regarding carbon monoxide" as part of discussions of 6 principal pollutants, U.S. EPA Office of Air and Radiation.

No health effects result when carboxyhemoglobin levels are under 4% to 5% in healthy adults. Carboxyhemoglobin levels of 2 to 3% may have negative effects on those with cardiovascular disease or other sensitivity. See, Environmental Health Criteria 13, Carbon Monoxide, World Health Organization, Geneva, Switzerland (1979), p. 15.



The US Occupational Safety and Health Administration's air contaminants regulation, 29 CFR Sec. 1910.1000, lists 50 ppm and approximately 55 mg/m<sup>3</sup> of CO as the 8-hour Time Weighted Average of exposure for the substance. 29 CFR Sec. 1910.1000.

By contrast, as explained below, the worst case estimated intake of CO attributable to AT2001 is 1.88 mg CO/meal.

The US FDA has not established an Acceptable Daily Intake for CO. Nevertheless, CO exposure, at levels much higher than those attributable to AT2001, for decades has been permitted within the existing FDA and USDA food additive regulatory provisions:

- Wood smoke ("smoke flavoring"), conventionally including CO as a component, is permitted by regulation as an ingredient in meat and poultry products pursuant to 9 CFR Secs. 318.7(c)(4)[meat], 381.147(c)(4)[poultry], 424.21(c).
- The specifications for Combustion product gas in 21 CFR Sec. 173.350 permit CO up to 4.5 percent by volume in such gases, which may be used in the processing and packaging of beverages and other foods except fresh meats, to remove and displace oxygen. Such gases are commonly used to package fruits and vegetables.

- In 2000, FDA responded favorably to GRAS Notice 000015 from Hawaii International Seafood, Inc. for the use of tasteless smoke, before freezing of tuna, as a preservative, specifically to preserve taste, aroma, texture and color. GRAS Notice (GRN) No. 000015, March 10, 2000. CO is a primary component of conventional smoke, which that Notice asserts is Generally Recognized As Safe based on decades of safe use in a variety of foods, which uses are recognized by FDA and incorporated into numerous food standards that permit smoking of cheeses. CO is also a primary component of tasteless smoke, along with nitrogen, oxygen, carbon dioxide and methane. The tasteless smoke is used to impart a "preservative effect." As noted in FDA's March 10, 2000 letter about the GRAS Notice, "In Hawaii International's view, tuna treated with tasteless smoke and tuna treated with conventional smoke contain comparable levels of carbon monoxide, carbon dioxide, hydrocarbons, and phenols." FDA's letter notes that it "has no questions at this time" regarding Hawaii International's conclusion that the use described is GRAS, though, in keeping with current regulatory practice, it had not made its own determination.

000014

CO is listed as a reproductive toxicant by the State of California pursuant to its Safe Drinking Water and Toxic Enforcement Act of 1986 ("Proposition 65"). California law contemplates that exposures to listed reproductive toxicants will be accompanied by a warning, unless the exposure is less than 1/1000<sup>th</sup> of an established no observable effect level. Cal. Health and Safety Code, Sec. 25249.6. No such level has been established for CO. Almost without question, though, any such future level (which will have a dubious connection to safety principles in any event, due to the design of Prop. 65), will be more than 1,000 times any possible exposure that could result from AT2001. The worst-case potential exposures from AT2001 are tiny fractions of the established occupational and environmental exposure levels (see below), which themselves are certain to be well below any level at which reproductive toxicity is ultimately is deemed to result.

#### *Effects on Fresh Meat and Consumption*

Analysis of the AT2001 system makes plain the lack of any safety issue from consumption of treated meats. Additionally, similar technologies employing CO as part of a modified atmosphere gas mixture analyzed the technologies for effects on meat in terms of microbial load and organoleptic

properties including color, and for the safety of consumption of treated meats, specifically, any tendency of the consumed meat to expose consumers to levels of carboxymyoglobin. Further important evidence is obtained from examination of the actual experience since 1985 in Norway of packaging fresh red meats in 0.3 – 0.5 % CO for retail.

Safety: Effects on carboxymyoglobin levels

Consumption of meat treated with AT2001 is not expected to result in any measurable levels of carboxymyoglobin in the blood of those who consume treated meat.

An Estimated Daily Intake ("EDI") of CO attributable to the AT2001 use can be calculated as follows. First, we assume the following reasonable values for the exposure parameters:

- (1) An AT2001 bag contains 1.5 L modified atmosphere with a CO concentration of 0.4%, that is equivalent to approximately 0.006 L of CO in the bag ( = 6 mL CO).
- (2) At 28 g CO per mole and approximately 22.4 L per mole, the mass of CO per unit volume may be calculated:  $(28 \text{ g/mol}) / (22.4 \text{ L/mol}) = 1.25 \text{ g/L} = 1.25 \text{ mg/mL}$ .

- (3) The AT2001 bag contains 2 lbs (approximately 1.0 kg) of ground meat.
- (4) Approximately 30% of the total amount of CO is absorbed into the meat (based on Watts, D.A.; Wolfe, S.K.; Brown, W.D., "Fate of [ $^{14}\text{C}$ ] Carbon Monoxide in Cooked or Store Ground Beef Samples", J. Agric. Food Chem., Vol. 26, No. 1 (1978), pp. 210-214). Therefore, the amount of CO taken up by the meat is  $[(0.3)*(6 \text{ mL/bag})*(1.25 \text{ mg/mL})] / [1.0 \text{ kg meat/bag}] = 2.25 \text{ mg CO / kg meat}$ .
- (5) If we assume that a person consumes an 8.8 oz steak (250 g = 0.25 kg meat), or ground equivalent, at a single meal<sup>2</sup>, that 85% reduction in CO content occurs during cooking, and that 100% of the ingested CO is absorbed, then the maximum amount of CO exposure is  $(0.15)*(2.25 \text{ mg CO / kg meat})*(0.25 \text{ kg meat/meal}) = 0.084 \text{ mg CO/meal}$ .

Next, comparison may be made of the of the consumer EDI for CO to that amount inhaled during an 8-hour period at the EPA's National Ambient Air Quality Standard ("NAAQS") level. 40 CFR Sec. 50.8, National primary ambient air quality standards for carbon monoxide:

---

<sup>2</sup> Note that this is a conservative assumption. The EDI of beef for the 90<sup>th</sup> percentile intake per user is 139.2 g/d based on the most recent USDA national survey of food intake by individuals. Pactiv chose to use a larger value for beef consumption to simulate a typical to above-average consumption incident rather than an average over all meats.

The calculated, worst case consumer EDI for CO may be compared to that amount inhaled during an 8-hour period at the American Conference of Governmental Industrial Hygienists ("ACGIH") Threshold Limit Value ("TLV"). Documentation of the Threshold Limit Values and Biological Exposure Indices, p. 23, ACGIH, 1330 Kemper Meadow Drive, Cincinnati, Ohio.

- (1) The ACGIH TLV is 25 ppm CO is equivalent to approximately 28.9 mg CO per m<sup>3</sup> air.
- (2) The typical person breathes 15 m<sup>3</sup> air per day or approximately 5 m<sup>3</sup> air per 8-hours.
- (3) The exposure under these circumstances may be calculated as follows:

$$(28.9 \text{ mg/m}^3) * (5 \text{ m}^3/8\text{-hr}) = 145 \text{ mg CO} / 8\text{-hr}.$$

Thus, the ingestion of residual CO from the cooked meat is merely 1.3% of the exposure level at ACGIH TLV  $((1.88 \text{ mg}) / (145 \text{ mg}) = 0.013 = 1.3\%)$

Finally, the calculated worst case consumer EDI for CO may be compared to that amount inhaled during an 8-hour period at the OSHA PEL:

- (1) The OSHA PEL is 50 ppm CO is equivalent to approximately 58 mg CO per m<sup>3</sup> air.

- (2) The typical person breathes  $15 \text{ m}^3$  air per day or approximately  $5 \text{ m}^3$  air per 8-hours.
- (3) The exposure under these circumstances may be calculated as follows:

$$(58 \text{ mg/m}^3) * (5 \text{ m}^3/8\text{-hr}) = 290 \text{ mg CO} / 8\text{-hr}.$$

Thus, the ingestion of residual CO from the cooked meat is 0.65% of the exposure level at OSHA PEL.  $((1.88 \text{ mg}) / (290 \text{ mg}) = 0.0065 = 0.65\%)$ .

Thus, the consumer EDI of CO from a eating meat packaged in the Active Tech 2001 bag is a small fraction of any of the currently allowed exposures by authoritative agencies. As these various limits were established to protect individual safety and health, it is plain that the worst case exposures that may result from AT2001 present no safety concerns whatsoever.

In the 1997 study, "Technological, hygienic and toxicological aspects of carbon monoxide used in modified-atmosphere packaging of meat" Trends in Food Science and Technology, September 1997 [Vol. 8], pp. 307-312, Sørheim, et al. concluded that meat packaged and displayed in an atmosphere combining 60 to 70% carbon dioxide, 30 to 40% nitrogen, and less than 0.5% CO "will result in only negligible levels of carboxyhemoglobin in the blood."

The authors note that there was sparse information in published literature on exposure to CO after consumption of meat treated with CO gas. They note that "the inhalation of air containing CO at a level of 55 mg per m<sup>3</sup> (the acceptable level in working environments in the USA) would provide a COHb level for a prolonged time period (hours) of at least 14 times that of the level reached temporarily on the consumption of 225 g of meat that has been packaged in CO at the saturation level for myoglobin." That estimate assumed saturation of meat myoglobin and hemoglobin was maximal and the transfer of CO from the gastrointestinal tract to the blood was 100%. Sørheim, et al. (1997), p. 310. The authors concluded, "Consequently, even for such a "worst case" scenario, the treatment of meat with CO gas appears to contribute very little to COHb levels, relative to levels that are considered safe in the working environment." Sørheim, et al. (1997), p. 310.

The authors report that "CO is lost from previously CO-treated meat during storage in the absence of CO, with a half life of ~3d." Sørheim, et al. (1997), p. 310. As these fresh meats are to be cooked before consumption, CO lost during cooking is also relevant. The authors report that "When the beef was cooked at 195° C, only 0.1 mg of CO remained

000021



per kg of meat. The loss of CO amounted to ~85%." Sorheim, et al. (1997), p. 310.

The authors also compared CO exposure from the air and estimated exposure from CO-treated meat. Their comparative table is shown below.

Table 5. Theoretical Uptake of Carbon Monoxide (CO) in Blood

Exposure method	CO intake in 1 h	CO intake in 8 h
Lungs ( $15\text{m}^3$ )	$24\text{ mg} \times 0.625 = 15.1\text{ mg}$	$9.2\text{ mg} \times 5 = 46.0\text{ mg}$
Meat (250 g CO treated)	0.025 mg	0.025 mg

Sørheim, et al. (1997), p. 311, Table 5.

Part of the authors' analysis was the premise that absorption of CO from the gastrointestinal tract into blood will in all probability be less effective than absorption from the lungs. The authors summarized the comparison as follows:

In order to prevent a maximum COHb level in the blood of 1.5% being exceeded, the CO concentration in air for a 1h period of moderate physical activity should not exceed  $24\text{ mg}/\text{m}^3$ , or  $9.2\text{ mg}/\text{m}^3$  in 8h (according to Table 4). In contrast, the consumption of meat that had been treated for 3d in an atmosphere containing 1% CO yielded ~0.1 mg of CO per kg of meat on storage and cooking.

Sørheim, et al. (1997), p. 310, citing Watts, D.A.; Wolfe, S.K.; Brown, W.D., "Fate of [ $^{14}\text{C}$ ]Carbon Monoxide in Cooked or Stored Ground Beef Samples", *J. Agric. Food Chem.*, Vol 26, No. 1 (1978), pp. 210-214.

The authors calculate that CO intake in 1h through the lungs taking in  $15\text{m}^3$  per day would result in 15.1 mg of CO, as compared with 0.025 mg of CO from intake of 250 g of CO treated meat. In 8 hours, the authors say the lungs will take in 46.0 mg, and the figure for meat would still be 0.025 mg. As the authors conclude,

Estimates detailed above indicate that, even assuming an improbable 100% absorption of CO from the gastrointestinal tract into the blood, the consumption of meat that has been treated with 1% CO will result in COHb levels that are negligible (approximately 3 orders of magnitude lower) compared with those resulting from exposure in the working environment to CO at an acceptable level. Consequently, it is highly improbable that CO exposure from meat packaged in an atmosphere containing up to 0.5% will represent a toxic threat to consumers through the formation of COHb.

Sørheim, et al. (1997), p. 310.

In another published report, the storage life and characteristics of meats packaged in a modified atmosphere including 0.4% CO were studied, but under circumstances distinguishable from AT2001. Sørheim O; Nissen, H; Nesbakken, T, "The Storage Life of Beef and Pork Packaged in an Atmosphere With Low Carbon Monoxide and High Carbon Dioxide", 52 Meat Science 157-164 (1999). In the study, the meats were packed in

000023

modified atmosphere into retail-ready packages. This study examined off odor and microflora, as well as color, comparing the 0.4% CO/ 60% CO<sub>2</sub> /40% N<sub>2</sub> gas mixture with a gas mixture of 70% oxygen and 30% CO<sub>2</sub>.

Among the points made by these authors was that there is sometimes an objection raised against using CO in retail ready meats because "the colour stability can exceed the microbiological shelf life, with the risk of masking spoilage of the meat." Sørheim, et al. (1999), p. 163. (Citing Kropf, D.H. (1980), "Effects of retail display conditions on meat colour", *Proceedings of the Reciprocal Meat Conference*, 33, pp. 15-32.) The authors assert that in those circumstances, consumers would need to rely on off odors to evaluate microbiological conditions of meat. In addition, they caution, "When a MA with CO is applied commercially, it is important to have a proper control of hygienic condition of the meat raw materials and the chill chain temperatures." See Sørheim, et al. (1999), p. 163.

AT2001, by contrast, presents no such similar problems or needs for caution. AT2001 does not mask spoilage of the meat. AT2001 does not involve use of a modified atmosphere including CO in the retail package. Moreover, as noted below, Pactiv's own commissioned experimentation with AT2001 demonstrates that AT2001 retail packages will deteriorate in color beginning almost immediately after removal of the modified

atmosphere, and that microbial load will not reach unsafe levels while the color of AT2001 meat is still acceptable to the consumer.

Safety: The Norwegian experience

In Norway, CO has been used to package fresh meats, even at retail, since 1985, with commercially satisfactory and safe results.

The 2000 submission by the Norwegian Meat Cooperative and Norwegian Independent Meat Association to the EU Commission seeking Europe-wide approval of the use of CO, "Application For Assessment Of The Food Additive Carbon Monoxide (CO) Prior To Its Authorization", is Attachment 3. The evaluation undertakes a detailed analysis of the CO exposure expected through the described packaging use. See section entitled "IV. Report by Tore Aune: "Fresh Meat in Consumer Packaging-A Toxicological Evaluation of the Use of Up to 0.5% CO in a Gas Mixture".

As the Norwegian risk assessment analysis concludes, assuming a worst-case exposure of 0.1 mg/kg from consumption of 250 grams of heated CO-treated meat, CO intake can be expected to be 0.025 mg in 1 hour or even after 8 hours. Attachment 3, p. 000154. The cited study, Sørheim, et al. (1997), utilized meat that had been treated with 1% CO. According to the authors, to stay under maximum blood levels of carboxyhemoglobin of 1.5%, "the CO concentration in the air must be 24

000025

milligrams per  $\text{mg}/\text{m}^3$  for 1 hour at moderate physical activity at  $9.2 \text{ mg}/\text{m}^3$  for 8 hours...." Attachment 3, p. 000154. Assuming an adult inhales  $15 \text{ m}^3$  per 24 hours, this translates to 15.1 mg of CO taken in 1 hour, or 46.0 mg of CO taken in 8 hours. This is in dramatic contrast to the miniscule amount expected to be ingested through meat. The Norwegian authors conclude, "From a health perspective, the use of CO in concentrations below 0.5-1% for fresh meat thus represents no toxicological risk." Attachment 3, p. 000155.

Safety: Exposure in environment

As a basis for comparison, the possible effect on ambient CO concentration associated with the release from a typical AT2001 barrier bag was estimated. A typical AT2001 barrier bag contains approximately 1.5 liters of modified atmosphere with a CO concentration of 0.4 percent, which is equivalent to approximately 0.006 liters of CO within the bag. On a mass basis, this volume of CO is equivalent to approximately 0.0075 g (7.5 mg) CO per bag.

Consider the possible use of the bag for storage of meat prior to retail display (e.g., at a supermarket). Any unassociated CO within the bag, it can be assumed, would be released to the meat processing area when the bag is opened, resulting in possible exposure by the employee(s) to the

released CO. The extent of such exposure is dependent on several factors, including the size of the meat processing area, air-mixing within the area and between adjacent areas, the number of bags opened, and the amount of free CO unassociated with the meat in the package. For these calculations, it has been conservatively assumed that none of the CO has become associated with the meat and is therefore all free to the ambient atmosphere upon opening of the package.

Assume, however, that the air volume within a meat processing area may reasonably range from  $150 \text{ m}^3$  to  $1,500 \text{ m}^3$ , which would represent several hundred to several thousand square feet of processing area. If each bag introduces 7.5 mg CO to the air within the processing area, the corresponding concentration of CO in air would be in the range of  $0.005 \text{ mg/m}^3$  to  $0.05 \text{ mg/m}^3$ , assuming conservatively that there is no air exchange between the processing area and other rooms or the outdoors. Thus, to exceed the occupational safety standard (i.e., threshold limit value, or TLV) of 25 ppm ( $29 \text{ mg/m}^3$ ), 580 to 5,800 1.5 liter bags would need to be opened within an 8-hour period. As noted above, this assumes no mixing with other areas of the building or with outdoor air.

Thus, applying the reasonable assumption that the air volume within the processing area will be exchanged with external air once per hour,

000027

opening of 580 to 5,800 bags per hour would be required to exceed the TLV, or 4,600 to 46,000 bags per work day. The number of bags opened within a given processing area will be a function of the size of the processing area, to a given extent, but is unlikely to even approach the number of bags required to result in air concentrations at the TLV. Actual concentrations in the work area of a secondary processing facility would likely be one to two orders of magnitude below the standard. Thus, opening of bags within a work area will not alter significantly the environmental exposure to CO.<sup>3</sup>

Regardless, the opening of the bags does not alter significantly the environmental exposure to CO. This action qualifies for a categorical exclusion from preparation of an environmental assessment pursuant to 21 CFR Sec. 25.32 (i), which provides an exemption for, in pertinent part, "Approval of a ...GRAS affirmation petition...." 21 CFR Sec. 25.32(i). The regulation makes no specific mention of the GRAS Notice procedure, but similar treatment is warranted for a GRAS Notice. (We also note that CO as used here also qualifies for exclusion under 21 CFR Sec. 25.32(r), as CO "occurs naturally in the environment" and the noticed use "does not alter significantly the concentration or distribution of the substance, its

---

<sup>3</sup> As an aside, there is no reason to expect any difficulty achieving compliance with the OSHA Threshold

metabolites, or degradation products in the environment." 21 CFR Sec. 25.32(r).)

*Corroborative information about AT2001*

The specific AT2001 system has been thoroughly tested to confirm that it results in the expected limited exposures to CO, and has no adverse effects on the treated meats. A study of meats treated with AT2001 commissioned by Pactiv examined its effects on initial product color, stability of color during display, and the central safety consideration of the relationships between color deterioration and microbial populations.

The study, conducted by faculty of the Department of Animal Sciences & Industry of Kansas State University, Manhattan, Kansas, examined steaks from three cuts of beef (strip loin, tenderloin, and inside round steaks), as well as ground beef. The study report is Attachment 4. The meats were packaged in AT2001 atmosphere, then stored at 35° F or 43° F for up to 35 days. Cuts were then placed under simulated retail conditions by being removed from the AT outer package and displayed at 34° F until their color approached consumer unacceptability. Comparisons

---

Limit Value at plants using the AT2000 technology to fill bags. Experimental use of an exhaust hood over the machinery has resulted in no measurable increase in CO ppm levels near the line.

000029



were made to similar products that had been exposed to oxygen but not CO.

Among the study's conclusions were:

(1) *Color:* AT2001 system resulted in products that were equally red to products packaged with traditional oxygen permeable overwrap. When the AT2001 outer bag was removed, the product's conversion to oxymyoglobin occur red in 60-90 minutes and then had a typical bright red color. Visual appearance was improved, especially in the tenderloin and inner part of the inside round steaks, throughout display. Color deterioration compared well to baseline products exposed to oxygen. For tenderloin and inside round steaks, and to a lesser degree for ground beef, display time was increased only slightly in the AT2001 samples.

(2) *Bacterial growth:* Bacterial growth was neither encouraged nor suppressed by the addition of CO to the ActiveTech™ gas blend (nitrogen and carbon dioxide ), although microbial growth curves changed in slope and exponential growth according to the environment in the packages. Aerobic bacteria and facultative anaerobes followed typical patterns of growth according to environmental conditions.

(3) *Spoilage indicators: CO neither masked spoilage, nor extended color life beyond the point of wholesomeness (i.e., the point of microbial soundness).*

A summary of the study follows.

A random selection of all steaks and ground beef packaged using oxygen-permeable polyvinyl chloride ("PVC") film were placed in display to serve as a baseline for color and microbiological comparisons. Products were expected to have the lowest microbiological load and ideal color stability using traditional packaging and display conditions for products exposed only to atmospheric oxygen. The inherent muscle chemistry responsible for good color life also was optimal. If the product exposed to CO were to have extended meat color life, then it will be compared to the baseline "control" with the "best" possible color.

To measure color changes, visual scores were considered the "standard" with instrumental color being discussed relative to its agreement or disagreement with the visual panel, ie, did the objective measurements confirm what the color panel saw? Visual scores of  $\geq 3.5$  were considered borderline acceptable. When samples reached this discoloration, they were removed from display. Normally,  $a^*$  values (higher values indicate more redness) are highly correlated to visual appraisal.

Inside round steaks typically are two-toned in color. The inner portion (ISM) is much less color stable than the outer portion (OSM). These portions were scored separately since one portion may have acceptable color while the other has unacceptable color that would be discriminated against by consumers resulting in the whole cut being judged unacceptable in color. The effects of CO on this bi-colored muscle were needed to confirm that color was not excessively extended in either portion.

Average fat and moisture contents of the ground beef were 19.5 and 61.6%, respectively. The pH of both intact muscles and the ground beef ranged from 5.3 to 5.7. The initial aerobic plate counts and lactic bacteria counts for all products were relatively low and indicative of good microbial quality of the raw materials and good sanitation. Furthermore, coliforms and *E. coli* were below the detection limit throughout the study.

The color of ground beef and steaks entering display (after MAP storage at 2 temperatures) was an attractive, typical red color. Although there were several significant differences in visual scores and  $a^*$  values (Attachment 9, Table 2 and Figures 1-10 at day 0) for product CO vs. baseline cuts, the variation in color was usually within  $\pm 0.5$  of a color score.

Color results: In general, the initial color of product exposed to CO was very similar to the color of steaks from the baseline display (never exposed to CO). When differences occurred, they were more related to either storage temperature or postmortem age of the product.

Panelists did not consider the color of product exposed to CO atypical. Cuts exposed to CO generally appeared more uniformly bright-red and would be expected to have high consumer appeal. These results were expected, as CO is known to preferentially form a ligand with the colored pigment (myoglobin) in meat resulting in a more intense red pigment known as carboxymyoglobin.

In the AT2001 system, Pactiv uses a low level of 0.4% CO, and obtains a red color very similar to the normal red oxymyoglobin pigment of fresh meats exposed to oxygen.

Color stability results: A critical next question was whether the carboxymyoglobin formed on the surface was more stable than the oxymyoglobin formed in baseline product. Further, did the carboxymyoglobin deteriorate in a predictable way that consumers could continue to use visual color to judge freshness or potential spoilage?

Product exposed to CO during MAP storage had color deterioration during display. (See visual panel scores (Attachment 4, Figures 1-5) and

instrumental color ( $a^*$  values, Attachment 4, Figures 6-10).) As expected, visual scores increased (color deteriorated) and  $a^*$  values decreased (loss of redness) as days in display increased. In several instances, color appeared to improve late in display – as indicated by a decrease in visual scores (see ground beef, strips loins and tenderloins at 43°F). These decreases were not a return of redness, but resulted from removal of discolored packages the preceding period, leaving product with less overall discoloration remaining in the case.

In general, the color deterioration profiles followed an expected pattern. Namely, the freshest product (baseline packages) had the most stable, red color and the most days in display needed to reach borderline discoloration of all treatments. (Attachment 4, Table 3 scores to 3.5) Exceptions occurred for the inside portion of the inside round and tenderloin products, where the product exposed to CO had slightly more stable color than the baseline product (Attachment 4, Table 3). These two muscle areas are well known by retailers as having short color life. Thus, CO appeared to improve color life when the inherent muscle chemistry desired for color was limited.

For product from MAP, the longer the storage time, the faster the deterioration, especially at the higher storage temperature (Attachment 4,

000034

Tables 2 and 3). For packages stored at 43°F, which was a mildly abusive temperature, color deterioration would be expected to accelerate. This phenomenon also is illustrated in Attachment 4, Figures 1-10.

There was no evidence the color shelf life was unexpectedly lengthened by exposure of meat to CO in MAP. Changes in  $a^*$  values (and other instrumental measures of color not shown) followed the same pattern of color deterioration observed by the visual panelists.

Color and microbial data: Initial, pre-display microbiological data suggested that the raw materials were fresh and processed using good hygienic practices. For intact cuts, lactic acid bacteria, generic *E. coli*, and total coliform counts were below the detection limit of 1.76 colony forming units (CFU)/in<sup>2</sup>. Initial, pre-display aerobic plate counts ("APC") for intact muscles ranged from 1 to 1.63 log<sub>10</sub> CFU/in<sup>2</sup>. Post-display counts were higher ( $P < 0.05$ ) than pre-display APC which was an increase in bacterial proliferation and typical deterioration. However, all product had sufficient microbes to be susceptible to spoilage.

Baseline products were pulled from display when the visual panel scores reached  $\geq 3.5$ . However, the APC did not exceed 5 log<sub>10</sub> CFU/unit as shown in Attachment 4, Figures 15-18. Furthermore, off-odor scores for

product at end of display (Attachment 4, Table 3) ranged from no to slight off odor.

Thus, color life in this base population did not exceed microbial soundness, which is generally accepted as  $< 100$  million CFU/g hamburger ( $< 1 \times 10^8$ ). (Principles of Meat Science, 3d Ed., Hedrick, H.B.; Aberle, ED, Forrest, JD; Judge, MD; Merkel, RA, Eds, Kendall/Hunt Publishing Co., Dubuque, Iowa.

Similar trends in microbial growth occurred in post-displayed samples stored in MAP compared to baseline products. Microbial patterns for product deterioration are shown in Attachment 4, Table 4 and Figures 11-18. Products stored under MAP at a slightly abusive temperature showed, as expected, a more rapid increase ( $P < 0.05$ ) in microbial counts compared to samples stored at 35°F. For post-MAP (pre-display) and post-display samples, APC were higher at 45°F than 35°F (Table 4), and during the later days of storage at the higher temperature, differences were more obvious. Significant changes ( $P < 0.05$ ) occurred in all cuts and ground beef with the exception of semimembranosus muscle. Counts for the SM muscle were lower than expected and no significant changes occurring until day 35 of MAP storage. This suggests that quality products that have been handled

000036

in a sanitary fashion can be stored in the AT2001 system up to 35 days without comprising microbial quality.

The APCs for intact strip loin and tenderloin steaks stored at 35°F were lower ( $P < 0.05$ ) on all days of display on days 21 and 35 post-MAP than steaks stored at 43°F (Attachment 4, Figures 12 and 14). Although products did not show a difference in APCs 7 days post-MAP, those products stored at the higher temperature (43°F) were more inferior 21 and 35 days post-MAP.

One goal of this research was to see if the color of CO-treated meat might mask spoilage. Visual color scoring was considered as the "standard" for determining the time to remove products from display. Because the visual panel scores were the deciding factor for length of shelf life, the interdependence between visual color and APC, LAB, and odor were considered quite important.

Attachment 4, Figures 19-21 show aerobic and lactic bacterial growth and odor scores at the end of display plotted against their corresponding visual color scores. All data observations were summed over storage temperature, storage time, and product type and plotted in one graph. If color masked spoilage, then there would be multiple points in the upper left



quadrant of the plot, the area represented by unacceptable microbial counts and off odors but with acceptable color (i.e., scores <3.5).

This did not occur with any frequency in any of the three plots. Thus, it does not appear that exposure of meat to CO during extended (up to 35 days at either 35° or 43°F) caused meat color to hide spoilage.

**e. Statement of availability of information**

Notifier has relied on published studies and generally accepted scientific data and information as the basis of its conclusions, and those of its panel of experts, about the safety and the general recognition of a modified atmosphere system for meat incorporating 0.4% CO in the gas mixture.

**II. Identity of notified substance**

1. Chemical name: Carbon monoxide
2. Chemical Abstracts Service: 630-08-0
3. Composition Specifications for food-grade material: The CO employed in this system is to be of suitable purity for food contact. Specifically, this will mean a 99.99% minimum purity, as supplied by Pactiv's commercial gas supplier, Haun Welding Supply, Inc., 6481 Ridings

000038

Road, Syracuse, NY 13206. Attachment 5. The supplier's CO meets the following specifications, and will be referred to as "commercial grade":

Component	Specification
Carbon Monoxide	99.99% min.
Oxygen	$\leq 0.5$ PPM
Nitrogen	$\leq 10$ PPM
Carbon Dioxide	$\leq 20$ PPM
Methane	$\leq 5$ PPM
Ethane	$\leq 1$ PPM
Propane	$\leq 1$ PPM
Dimethyl Ether	$\leq 1$ PPM
Hydrogen	$\leq 1$ PPM
Moisture	$\leq 1$ PPM

4. Properties:

Relative molecule mass	28.01
Critical point	-140.2 °C at 34.5 atm (3.5 MPa)
Melting point	-205.1 °C
Boiling point	-191.5 °C
Density, at 0 °C, 1 atm	1.250 g/litre
at 25 °C, 1 atm	1.145 g/litre
Specific gravity relative to air	0.967
Solubility in water at 0 °C, 1atm	3.54 ml/100 ml
at 25 °C, 1 atm	2.14 ml/100 ml
at 37 °C, 1 atm	1.83 ml/100 ml <sup>a</sup>
Conversion factors:	
at 0 °C, 1 atm	1 mg/m <sup>3</sup> = 0.800 ppm <sup>b</sup>
	1 ppm = 1.250 mg/m <sup>3</sup>
at 25 °C, 1 atm	1 mg/m <sup>3</sup> = 0.873 ppm
	1 ppm = 1.145 mg/m <sup>3</sup>

<sup>a</sup> Value obtained by graphic or calculated interpolation (Altman et al., 1971).

<sup>b</sup> Parts per million by volume

5. Analyses: ASTM D1946, "Analysis of Reformed Gas by Gas Chromatography (GC) with Thermal Conductivity Detection (TCD)", may be utilized to measure the quantity of CO present in gas mixtures. A copy of the method is Attachment 6.

### III. Self-limiting levels of use

Studies of modified atmospheres for packaging meat that contained both higher and lower levels of CO have established that the 0.4% used in the AT2001 system both has advantageous characteristics and avoids disadvantages seen with lower or higher levels. A CO level of 0.4% is sufficient to produce stable, cherry red color, (Sørheim, et al. (1997), and use of CO through retail display time may result in masked spoilage.

### IV. Basis of GRAS determination.

Pactiv believes its use of CO is GRAS based on scientific procedures, 21 CFR Sec. 170.30(b), and convened a panel of experts qualified by scientific training and experience to evaluate the safety of food, food additives and food ingredients. The experts have reviewed and evaluated the publicly available information summarized in this GRAS Notice. Their testimonial

000040

letters are attached as Attachments 7 through 10. The above discussion and citations to generally available accepted scientific data, information, methods and principles relied upon, together with the anticipated consumption levels for both CO and meat treated with CO, provide ample basis to conclude that the use of CO at 0.4% in a modified atmosphere for packaging fresh meats is both safe and generally recognized as such by qualified experts.

The panel consisted of the following experts, whose GRAS opinions and curricula vitae are attached as attachments 7 through 10.

1. Daren Cornforth, Ph.D.  
Professor  
Department of Nutrition and Food Sciences  
Utah State University  
750 N. 1200 E.  
Logan, Utah 84322-8700

Dr. Cornforth is a professor in Nutrition and Food Sciences at Utah State University, Logan, Utah, and received his Ph.D. in food science and human nutrition from Michigan State University. He has performed extensive research and published multiple articles on the subject of meat color.

2. Vasilios Frankos, Ph.D.  
Principal  
Environ Corp.  
4350 N. Fairfax Dr.

000041

Suite 300  
Arlington VA 22203

Dr. Frankos is a Principal at ENVIRON corporation, Arlington, Virginia, a scientific consulting firm, and has over 20 years of experience in the toxicological and pharmacological evaluation of data used to assess the risks posed by foods, food additives, and other substances. He holds a Ph. D. from the University of Maryland Pharmacy School in Pharmacology and Toxicology.

3. Melvin C. Hunt, Ph.D.  
Professor  
Weber Hall  
Dept. of Animal Sciences and Industry  
Kansas State University  
Manhattan, KS 65506

Dr. Hunt is a professor of food science at the Department of Animal Sciences and Industry at Kansas State University, Manhattan, Kansas. He received his Ph.D. in food science at the University of Missouri. Among his many research projects and publications are multiple studies relating to meat color and the effects of various environments on meat color.

4. Oddvin Sørheim, Ph.D.  
Senior Research Technologist  
MATFORSK – Norwegian Food  
Research Institute  
Osloveien 1

000042

N-1430 Ås  
Norway


Dr. Sørheim is a Senior Research Technologist at the Norwegian Food Research Institute, Osloveien, Norway. He received his Ph.D. in food science from the Agricultural University of Norway, and has performed extensive research and industry consultation, and published numerous articles on meat, including extensive experience with the use of CO in modified atmosphere packaging of meat.

Pactiv is not aware of any reports of investigations that are inconsistent with the GRAS determination relating to the use described.

### Conclusion

Based on all the above information, Pactiv Corporation has concluded that its use of 0.4% CO within the AT2001 modified atmosphere system for packaging fresh meat is Generally Recognized as Safe within the meaning of 21 U.S.C. Sec. 321(s).

Sincerely,

  
Eric F. Greenberg

000043

**Attachments:**

- Attachment 1 Authorization letter from Pactiv Corporation for representation by Eric F. Greenberg
- Attachment 2 Photographs of meats treated with ActiveTech™ and ActiveTech™ 2001
- Attachment 3 "Application for Assessment of the Food Additive Carbon Monoxide (CO) Prior to its Authorization", Norwegian Meat Cooperative; Norwegian Independent Meat Association (1999)
- Attachment 4 "Evaluation of Beef Steaks and Ground Beef in the Pactiv ActiveTech™ Packaging System: Effects of Carbon Monoxide in the Package Atmosphere", Hachmeister, K; Hunt, M.; Milliken, G; May 2001.
- Attachment 5 Certificate of Conformance, Carbon monoxide, Haun Welding Supply, Inc., Syracuse, New York, May 8, 2001.
- .....
- Attachment 6 "Standard Practice for Analysis of Reformed Gas by Gas Chromatography", ASTM D 1946-90 (Reapproved 2000).
- Attachment 7 Daren Cornforth, Ph.D., letter and curriculum vitae
- Attachment 8 Vasilios H. Frankos, Ph.D., letter and curriculum vitae
- Attachment 9 Melvin C. Hunt, Ph.D., letter and curriculum vitae
- Attachment 10 Oddvin Sørheim, Ph.D., letter and curriculum vitae

000044

**THIS PAGE BLANK**

000045



**THIS PAGE BLANK**

000046

**THIS PAGE BLANK**

000047

THIS PAGE BLANK

000048

## List of Attachments

### Attachments:

- Attachment 1 Authorization letter from Pactiv Corporation for representation by Eric F. Greenberg ..... 000050
- Attachment 2 Photographs of meats treated with ActiveTech and ActiveTech 2001 ..... 000053
- Attachment 3 "Application for Assessment of the Food Additive Carbon Monoxide (CO) Prior to its Authorization", Norwegian Meat Cooperative; Norwegian Independent Meat Association (1999) ..... 000058
- Attachment 4 "Evaluation of Beef Steaks and Ground Beef in the Pactiv ActiveTech Packaging System: Effects of Carbon Monoxide in the Package Atmosphere", Hachmeister, K; Hunt, M.; Milliken, G; May 2001 ..... 000157
- Attachment 5 Certificate of Conformance, Carbon monoxide, Haun Welding Supply, Inc., Syracuse, New York, May 8, 2001 ..... 000190
- Attachment 6 "Standard Practice for Analysis of Reformed Gas by Gas Chromatography", ASTM D 1946-90 (Reapproved 2000) ..... 000192
- Attachment 7 Daren Cornforth, Ph.D., letter and curriculum vitae ..... 000198
- Attachment 8 Vasilios H. Frankos, Ph.D., letter and curriculum vitae ..... 000205
- Attachment 9 Melvin C. Hunt, Ph.D., letter and curriculum vitae ..... 000221
- Attachment 10 Oddvin Sorheim, Ph.D., letter and curriculum vitae ..... 000247

000049

**ATTACHMENT 1**



**PACTIV**

Advanced Packaging Solutions

August 9, 2001

Division of GRAS Notice Review  
Office of Food Additive Safety  
Center for Food Safety and Applied Nutrition  
Food and Drug Administration  
200 C St, SW  
Washington, DC 20204

Pactiv Corporation  
Technology Center  
2651 Brickyard Road  
Canandaigua, New York 14424-1026

Re: Authorization of counsel regarding  
**NOTIFICATION OF CLAIM FOR GENERAL  
RECOGNITION OF SAFETY OF CARBON MONOXIDE  
IN A MODIFIED ATMOSPHERE  
SYSTEM FOR PACKAGING FRESH MEAT**

To the FDA:

Please take note that Pactiv Corporation, with headquarters at 1900 West Field Court, Lake Forest, Illinois, 60045, authorizes its attorney, Eric F. Greenberg, 3500 Three First National Plaza, Chicago, Illinois 60602, to represent it and communicate on its behalf in all matters regarding Pactiv's NOTIFICATION OF CLAIM FOR GENERAL RECOGNITION OF SAFETY OF CARBON MONOXIDE IN A MODIFIED ATMOSPHERE SYSTEM FOR PACKAGING FRESH MEAT.

Sincerely,

For PACTIV CORPORATION

By: Vinod K. Luthra  
General Manager  
New Business Development  
2651 Brickyard Road  
Canandaigua, New York 14424

000051

### *Summary regarding Pactiv Corporation*

Pactiv Corporation, 1900 West Field Court, Lake Forest, Illinois, is a leading provider of advanced packaging solutions to customers around the world. The company employs 17,000 people in 87 facilities worldwide. Annual revenues exceed \$3 billion.

Pactiv manufactures, markets and sells plastic and paper-based consumer products and food/foodservice packaging as well as protective and flexible packaging. Approximately 80% of its revenue comes from products made from different types of plastics, with the balance from paper and aluminum products.

The company's products include a wide range of items for consumers, food processors, supermarkets, foodservice entities, and the construction, automotive, computer, electronic, furniture and durable goods industries. The consumer products are sold under such recognized brand names as Hefty® , Baggies® , Hefty One-Zip® , Kordite™ and E-Z Foil®.

Pactiv further fuels internal growth by developing and commercializing proprietary new products and by designing value-added product-line extensions. In 1998, the consumer products and food/foodservice packaging business introduced over 80 new products and product-line extensions. In the protective and flexible packaging business, where custom design services drive revenues, Pactiv developed over 500 custom product applications in 1998. New product innovations include ActiveTech™ packaging, a proprietary modified atmospheric package used by food processors for case-ready meat.

**ATTACHMENT 2**



ActiveTech™

For Case Ready Applications

# Case Ready Packaging System


ActiveTech  
Case Ready Packaging System




000055

First Display Day

Hi-Oxygen



20 Days Stored



Fresh




Beef: 80% Lean Ground Beef

Hi-Ox, 20 Days Stored in "Short Gas", and Fresh  
(0.4% CO/ 35% CO<sub>2</sub>/ 64.6% N<sub>2</sub>)

Third Display Day

Hi-Oxygen



20 Days Stored



Fresh




Beef: 80% Lean Ground Beef

Hi-Ox, 20 Days Stored in "Short Gas", and Fresh  
(0.4% CO/ 35% CO<sub>2</sub>/ 64.6% N<sub>2</sub>)

Second Display Day


Hi-Oxygen



20 Days Stored



Fresh




Beef: 80% Lean Ground Beef

Hi-Ox, 20 Days Stored in "Short Gas", and Fresh  
(0.4% CO/ 35% CO<sub>2</sub>/ 64.6% N<sub>2</sub>)

Fourth Display Day


Hi-Oxygen



20 Days Stored



Fresh



Beef: 80% Lean Ground Beef


Hi-Ox, 20 Days Stored in "Short Gas", and Fresh  
(0.4% CO/ 35% CO<sub>2</sub>/ 64.6% N<sub>2</sub>)

First Display Day

20 Days Stored



Fresh Cut



Beef: Top Round Steak

Fresh Cut and 20 Days Stored in "Short Gas"  
(0.4% CO/ 35% CO2/ 64.6% N2)

Second Display Day

Fresh Cut



20 Days Stored



Beef: Top Round Steak

Fresh Cut and 20 Days Stored in "Short Gas"  
(0.4% CO/ 35% CO2/ 64.6% N2)

Third Display Day

20 Days Stored



Fresh Cut



Beef: Top Round Steak

Fresh Cut and 20 Days Stored in "Short Gas"  
(0.4% CO/ 35% CO2/ 64.6% N2)

Fourth Display Day

20 Days Stored



Fresh Cut



Beef: Top Round Steak

Fresh Cut and 20 Days Stored in "Short Gas"  
(0.4% CO/ 35% CO2/ 64.6% N2)

000057

The pages immediately following illustrate:

1. On the top of the photograph on the first page, an example is shown of the structure utilized for both ActiveTech™ and AT2001 incorporating tray, flexible overwrap, outer bag and activated oxygen scavenging sachet.
2. The second page of photographs show examples of ground meat color during the first, second, third and fourth days of display after packaging in (1) Hi-oxygen; (2) AT2001 atmosphere (referred to in captions as "Short Gas", 0.4% CO/35% CO<sub>2</sub>/64.6 N<sub>2</sub>), and after being held in that atmosphere for 20 days; and (3) Fresh.
3. The third page of photographs show examples of whole muscle meat (top round steak) color during the first, second, third and fourth days of display after packaging in (1) AT2001 atmosphere (referred to in captions as "Short Gas", 0.4% CO/35% CO<sub>2</sub>/64.6 N<sub>2</sub>), and after being held in that atmosphere for 20 days; and (2) Fresh cut.

**ATTACHMENT 3**

## APPLICATION FOR ASSESSMENT OF THE FOOD ADDITIVE CARBON MONOXIDE (CO) PRIOR TO ITS AUTHORIZATION

(This application is based on the document "Presentation of an application for assessment of a food additive prior to its authorisation", Office for Official Publications of the European Communities, Luxembourg, 1989, ISBN 92-826-0135-8).

### PART I. ADMINISTRATIVE DATA

I.1. Applicants: Altogether, the two applicants represent the total meat industry in Norway

Applicant no. 1:

The name of the applicant:

Norsk Kjøttssamvirke (Norwegian Meat Cooperative)

Address:

Lørenveien 37  
P.O.Box 360 Økern  
0513 Oslo, Norway

Other means of communication:

Telephone: +47 22 09 21 00  
Fax: +47 22 15 59 08

Applicant no. 2:

The name of the applicant:

Kjøttbransjens Landsforbund (The Norwegian Independent Meat Association) – represents the private meat industry in Norway

Address:

Karoline Kristiansensvei 2, Fyrstikktorget  
P.O.Box 6279 Etterstad  
0603 Oslo, Norway

Other means of communication:

Telephone: +47 23 24 44 70, Fax: +47 23 24 44 80

I.2. The name of the manufacturer(s) of the substance:

RIVOIRA S.P.A., Stabilimento Chivasso gas, Via cardinal Massaia 75L, I-10147 Torino, Italy

I.3. The name of the person responsible of the dossier:

Research director Truls Nesbakken, Norwegian Meat Research Centre, P. O. Box 396 Økern, 0513 Oslo, Norway. Telephone +47 22 09 23 99, Mobile phone +47 91 87 81 46, Fax +47 22 22 00 16, e-mail: truls.nesbakken@fagkjott.no

I.4. The table of contents of dossier

This dossier is sent through the Norwegian Food Control Authority (Statens næringsmiddeltilsyn). Together with this document follow as enclosures:

- 1) Nissen, H., Alvseike, O., Bredholt, S., Holck, A. and Nesbakken, T. (submitted) Packaging of ground beef in an atmosphere with high carbon dioxide and low carbon monoxide restrains growth of *Yersinia enterocolitica*, *Listeria monocytogenes* and *Escherichia coli* O157:H7. Int. J. Food Microbiol. As long as this work is not published, please handle this information with care.
- 2) Nissen, H., Alvseike, O., Bredholt, S., Holck, A. and Nesbakken, T. (1999) Packaging of ground beef in an atmosphere with low carbon monoxide and high carbon dioxide restrains growth of *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Yersinia enterocolitica* and *Salmonella diarizonae*. In: Tuijtelaars, A.C.J., Samson, R.A., Rombouts, F.M., Notermans, S., (Eds.), Food Microbiology and food safety into the next millenium. Proceedings of the Seventeenth International Conference of the International Committee on Food Microbiology and Hygiene (ICFMH), 13-17 September 1999, Veldhoven, The Netherlands, pp. 285-286.
- 3) Solheim, R. (1996) Consumer purchase probability of beef and pork packaged in different atmospheres. Report, Matforsk, 10 pp.
- 4) Sørheim, O. (1996) Discoloration of meat as an indicator of leakages in packages containing a CO gas mixture. Report, Matforsk, 5 pp.
- 5) Sørheim, O., Aune, T. and Nesbakken, T. (1997a) Technological, hygienic and toxicological aspects of carbon monoxide used in modified-atmosphere packaging of meat. Trends in Food Sci. Technol. 8, 307 - 312
- 6) Sørheim, O., Nissen, H., and Nesbakken, T. (1999) The storage life of beef and pork packaged in an atmosphere with low carbon monoxide and high carbon dioxide. Meat Sci. 52, 157 - 164

- 7) Letter from the director of Swedish Meats (which is the organisation of the Swedish meat cooperative) supporting the Norwegian meat industry's application to the EU Commission
- 8) Letter from the director of Swedish Meat Trade Association (which is the organisation of the private meat industry in Sweden) supporting the Norwegian meat industry's application to the EU Commission
- 9) Letter from the director of the Danish Pig Producers and Slaughterhouses, Copenhagen, Denmark supporting the Norwegian meat industry's application to the EU Commission
- 10) Letter from the Spanish Meat Industry's Association supporting the Norwegian meat industry's application to the EU Commission
- 11) Letter from the Finnish Meat Research Institute supporting the Norwegian meat industry's application to the EU Commission

## PART II. TECHNICAL DATA

### II.1. Name of the substance

- names in the IUPAC nomenclature: carbon monoxide
- other names (usual name/trade name/synonyms: carbon oxide, carbon monoxide)
- abbreviations: CO
- CAS number (if this has been attributed): 630 - 08 - 0

### II.2. Specification of the substance

- composition (% , m/v, mg/kg), e.g. in the case of heterogeneous products): 100%
- empirical and structural formula: CO
- molecular weight: 28.010 g/mole
- degree of purity (%): higher than 99.3%
- nature of known impurities/percentage of significant and main impurities: (< 0.7%):

Impurities:	Concentration:
Oxygen + Argon	< 2500 vpm
Water	< 20 vpm
THC (Total hydrocarbons)	< 500 vpm
Hydrogen + Nitrogen	< 3500 vpm
Carbon dioxide	< 500 vpm



- physical form (liquid, powder, etc.): gas
- solubility (e.g. aqueous, organic solvents, lipid):

solubility in water, 0°C, a CO partial pressure of 101.325 kPa = 3.537 cm<sup>3</sup>/100 cm<sup>3</sup>. Solubility in organic solvents and lipid: not relevant – see Part II.6 Exposure

- other data that the applicant believes may be useful to identify the substance (e.g. physico-chemical properties, analytical data on differences between batches):

Thermodynamic properties of carbon monoxide as ideal gas at 25°C:

Heat capacity,  $c_p$ : 29.142 J/(mol \* °K)

Entropy, S: 197.543 J/(mol \* °K)

Enthalpy: 8.669 kJ/mol

- information on the microbiological characteristics, in particular on the possible presence of pathogens and bacterial or mycotoxins: not relevant

### II.3. Manufacturing process

- Information on the method of manufacture (i.e. the process by which the raw materials are converted to the finished product):

The CO-gas is bought from RIVOIRA S.P.A., Torino, Italy (see I.2.). Hydro Rjukan Næringspark; P.O.Box 43-44, N-3661 Rjukan, Norway, makes the two CO-gas mixtures which are used in the Norwegian meat industry. They are called "Pakkemix NC1" and "DNC 29.7 – 0.3"

1) Pakkemix NC1 = 1.0% CO + 99% N<sub>2</sub>

The production:

- evacuation of an empty cylinder to under 10 mbar
- flushing with N<sub>2</sub>, quality 5.0
- repeat the evacuation of an empty cylinder to under 1 mbar
- manometric filling with CO, quality 2.3
- manometric filling with N<sub>2</sub> quality 5.0
- every tenth cylinder is analysed with gas chromatograph (GC) and thermoconductivity detector (TCD)

The pressure of the cylinder is 200 bar.

2)  $\text{DNC } 29.7 - 0.3 = 0.3\% \text{ CO} + 29.7\% \text{ N}_2 + 70\% \text{ CO}_2$

The production:

- a) evacuation of an empty cylinder to under 10 mbar
- b) flushing with  $\text{N}_2$ , quality 5.0
- c) repeat the evacuation of an empty cylinder to under 1 mbar
- d) manometric filling with  $\text{CO}$ , quality 2.3
- e) manometric filling with  $\text{N}_2$ , quality 5.0
- f) manometric filling with  $\text{CO}_2$ , quality 3.0
- g) every tenth cylinder is analysed with gas chromatograph (GC) and thermoconductivity detector (TCD)

The pressure of the cylinder is 50 bar..

#### II.4. Methods of analysis

- analytical methods to describe the substance, evaluate its purity and measure its physico-chemical and microbiological characteristics:

Oxygen + Argon, and Hydrogen + Nitrogen: Gas chromatograph with thermoconductivity detector (TCD) – min. detect. limits 10 vpm (RIVOIRA S.P.A., Stabilimento Chivasso gas, Via cardinal Massaia 75L, I-10147 Torino, Italy)

THC: Gas chromatograph with flame ionization detector (FID) – min. detect. limit 0.5 vpm (RIVOIRA S.P.A., Stabilimento Chivasso gas, Via cardinal Massaia 75L, I-10147 Torino, Italy)

Carbon dioxide: Gas chromatograph with helium ionization detector (HID) – min. detect. limit 0.5 vpm (RIVOIRA S.P.A., Stabilimento Chivasso gas, Via cardinal Massaia 75L, I-10147 Torino, Italy)

Water: Specific water analyzer – min. detect. limit 0.1 vpm (RIVOIRA S.P.A., Stabilimento Chivasso gas, Via cardinal Massaia 75L, I-10147 Torino, Italy)

- analytical methods for the determination of the additive and its degradation products (where relevant), in the foodstuff of which the substance is to form part:

The isotope,  $\text{C}^{14}$ , might be used for measuring  $\text{CO}$  before and after heat treatment (Watts et al., 1978). Spectrophotometry is used to measure carboxymyoglobin at 540 and 577 nm (native) or 425, 542 and 570 nm (denaturated) (El-Badawi et al., 1964; Cornforth, 1994).

## II.5. Justification for the additive

### - intended use and purpose:

Fresh red meat (mainly beef, pork and lamb, but also horse, goat, reindeer, game etc.) packaged in an atmosphere with 60 - 70% carbon dioxide (CO<sub>2</sub>), 30 - 40% nitrogen (N<sub>2</sub>) and < 0.5% carbon monoxide (CO) (high CO<sub>2</sub>/low CO mixture).

Gas mixtures with low concentrations of CO and high concentrations of CO<sub>2</sub> provide a combination of a long microbiological shelf life and a stable bright red colour of meat (Sørheim et al., 1999).

- the quantity to be added to specific foods and the residues in food: < 0.5% CO.
- investigations on the efficacy of the substance for the intended effect at the level proposed:

The main function of low levels of CO in modified atmospheres (MAs) is to give a stable, cherry red colour of the meat through strong binding of CO to myoglobin and formation of carboxymyoglobin (El-Badawi, 1964). Although a substantial increase in the shelf life of meat can be obtained by using various MAs, it is often limited by discolouration due to oxidation of myoglobin to metmyoglobin. This discolouration can be prevented by including a small fraction of CO in the gas mixture. Carboxymyoglobin is more resistant to oxidation than oxymyoglobin, due to the stronger binding of CO to the iron-porphyrin site on the myoglobin molecule (Wolfe, 1980). CO in concentrations of 1 - 5% had the ability to increase metmyoglobin reduction, even in the presence of air (Lanier et al., 1978).

The high CO<sub>2</sub>/low CO mixture and absence of O<sub>2</sub> provides a unique combination of a long microbiological shelf life (caused by the high CO<sub>2</sub> level) and a stable cherry red colour (caused by the low CO level). CO<sub>2</sub> inhibits growth of many microorganisms, but it has no effect *per se* on the colour of the meat (Renerre and Labadie, 1993). This gas is absorbed in meat and fat tissue at a ratio of approximately 1 litre gas per kg tissue (Gill, 1988). N<sub>2</sub> affects neither the microbiology nor the colour of the meat, but prevents collapse of the packages because it is not absorbed in the product. O<sub>2</sub> supports the growth of aerobic microorganisms, and removal of O<sub>2</sub> from the MA will therefore extend the microbiological shelf life. The shelf life of meat is considerably longer in the high CO<sub>2</sub>/low CO mixture than in the commonly used atmosphere of high oxygen (O<sub>2</sub>) with approximately 70% O<sub>2</sub> and 30% CO<sub>2</sub>. Consumption of meat treated with the high CO<sub>2</sub>/low CO mixture will result only in negligible levels of carboxyhaemoglobin in blood. It is highly improbable that CO from packaging of meat will present a toxic threat to the consumer (Sørheim et al., 1997a).

Shelf life in the high CO<sub>2</sub>/low CO mixture in comparison with alternative packaging methods:

Ground beef, beef loin steaks and pork chops were packaged in MAs of 0.4% CO/ 60% CO<sub>2</sub>/ 40% N<sub>2</sub> (high CO<sub>2</sub>/low CO mixture) and 70% O<sub>2</sub>/ 30% CO<sub>2</sub> (high O<sub>2</sub> mixture). In addition ground beef was packaged in clipped chub packs, beef loin steaks were vacuum packaged, and pork chops were packaged in an atmosphere of 60% CO<sub>2</sub>/ 40% N<sub>2</sub> with each pack containing an

O<sub>2</sub> absorber. The packs were stored in the dark at 4°C or 8°C for up to 21 days. Meat in the high CO<sub>2</sub>/low CO mixture had a stable bright red colour. The storage lives in this gas mixture at 4°C, as limited by off-odours, were 11, 14 and 21 days for ground beef, beef loin steaks and pork chops, respectively. The high O<sub>2</sub> mixture resulted in an initially bright red to red colour of the meat, but the colour was unstable and off-odours developed rapidly. The off-odours probably were caused by *Brochothrix thermosphacta*, which grew in all meat types, and in ground beef by pseudomonads also. Meat stored in chub packs, vacuum packs and 60% CO<sub>2</sub>/40% N<sub>2</sub> with an O<sub>2</sub> absorber developed off-odours and microflora similar to those of meat in low CO/high CO<sub>2</sub> mixture with however less acceptable colours or appearances. These results show that a low CO/high CO<sub>2</sub> atmosphere is effective for preserving retail-ready meat (Sørheim et al., 1999).

#### Aspects of spoilage:

Consumers may evaluate the shelf life of packaged meat based on its colour. A possible negative aspect of using CO in modified atmosphere packaging (MAP) of retail meat is a concern that the consumer might misjudge the product, because the microbiological status may be masked by the stable cherry red carboxymyoglobin colour (Knopf, 1980). However, the consumer is able to detect spoilage by off-odour (Sørheim et al., 1999). This is in contrast to ready to eat products such as cooked, sliced vacuum or gas packaged meat, gas packaged vegetables and vacuum-packaged cheeses where the consumers often have to taste it before judging the product as unacceptable. As ready to eat products, they also represent a higher risk than fresh meat packed in the high CO<sub>2</sub>/low CO mixture which is heat treated before consumption. In the current low concentrations, below 0.5%, CO *per se* seems to have no or only minor effects on bacteria and the shelf life of the meat. The combination of CO with high concentrations of CO<sub>2</sub>, for example 60 - 70%, is necessary for microbiological control. MAP enables centralised packaging operations with quality control, but MAP alone is no guarantee for the shelf life of the product. Sufficient shelf life can only be obtained through a proper quality control of raw materials, production, packaging, chill chain and retail conditions (Sørheim et al., 1999).

#### Pathogens in the high CO<sub>2</sub>/low CO mixture in comparison with alternative packaging methods:

Growth of the pathogens *Yersinia enterocolitica*, *Listeria monocytogenes*, *Escherichia coli* O157:H7 and strains of *Salmonella* was compared in ground beef packed in high CO<sub>2</sub>/low CO mixture, high O<sub>2</sub> mixture and in chub packs. The ground beef was inoculated with rifampicin or nalidixic acid/streptomycin-resistant strains (final concentration 10<sup>2</sup>-10<sup>3</sup> bacteria/g) and stored at 4°C and 10°C for up to 14 days. At 4°C the shelf life based on stable colour and reduced background flora was prolonged for the high CO<sub>2</sub>/low CO mixture, compared to the two other packaging methods, but at 10°C the shelf life was < 8 days for all the packaging methods. Growth of *Y. enterocolitica* was nearly totally inhibited both at 4°C and 10°C in the high CO<sub>2</sub>/low CO mixture, while the bacterial numbers in the samples packed in the high O<sub>2</sub> mixture increased from about 5x10<sup>2</sup> bacteria/g at day 0 to about 10<sup>4</sup> at day 5 at 4°C and to 10<sup>5</sup> at 10°C. Growth in the chub packs was even higher. *L. monocytogenes* showed very little growth at 4°C in all treatments. At 10°C there was slow growth of *L. monocytogenes* from about 5x10<sup>3</sup> bacteria/g to about 10<sup>4</sup> at day 5 in the high CO<sub>2</sub>/low CO mixture, while the numbers in the high O<sub>2</sub> mixture and the chub packs were about 10 times higher. Growth of *E. coli* O157:H7 at 10°C in the ground beef was nearly totally inhibited in both the high CO<sub>2</sub>/low CO mixture and the high O<sub>2</sub> mixture. Growth of *E. coli* O157:H7 in the chub packs was higher reaching 10<sup>5</sup> bacteria/g on day

5. The *Salmonella* strains (*S. Typhimurium*, *S. Dublin*, *S. Enteritidis* and *S. enterica* «61:k1, 5.(7)») in the ground meat stored at 10°C for 5 and 7 days, grew to a higher number in the high CO<sub>2</sub>/low CO mixture than in the high O<sub>2</sub> mixture. The present study shows that the prolonged shelf life at 4°C did not increase growth of *Y. enterocolitica* and *L. monocytogenes* in ground beef stored in the high CO<sub>2</sub>/low CO mixture, but the observed growth of strains of *Salmonella* in this mixture and in chub packs at the abuse temperature of 10°C does emphasise the importance of temperature control during storage (Nissen et al. 1999; Nissen et al., submitted).

#### Consumer purchase probability:

Ground beef, beef loin steaks and pork chops were packaged in high CO<sub>2</sub>/low CO mixture and high O<sub>2</sub> mixture. In addition ground beef was packaged in clipped chub packs, beef loin steaks were vacuum packaged, and pork chops were packaged in an atmosphere of 60% CO<sub>2</sub>/ 40% N<sub>2</sub> with each pack containing an O<sub>2</sub> absorber. The purchase probability data were collected by interviewing 126 consumers usually purchasing meat and meat products. The consumers visually compared the samples within each type of meat after 4 days storage at 4°C. The consumers preferred ground beef packaged in the high CO<sub>2</sub>/low CO mixture or the high O<sub>2</sub> mixture compared to ground beef packaged in clipped chub packs. Purchase probability increased when pork chops were packaged in the high CO<sub>2</sub>/low CO mixture. Pork chops in packs containing an O<sub>2</sub> absorber, were rated lowest in purchase probabilities. The purchase probability for beef loin steaks was similar when packaged in the high CO<sub>2</sub>/low CO mixture or the high O<sub>2</sub> mixture, and these products were preferred compared to beef loin steaks packaged in vacuum (Solheim, 1996)

- documentation on the need for the additive:

#### Alternatives to the high CO<sub>2</sub>/low CO mixture:

The most common MA for retail packaging of meat today contains O<sub>2</sub> at high concentrations combined with CO<sub>2</sub>, approximately 70% O<sub>2</sub>/ 30% CO<sub>2</sub>. The shelf life of meat in high O<sub>2</sub> atmospheres in commercial practice, typically at temperatures of 6 - 8°C, is about 7 days, limited by both microbiological spoilage and discolouration. Meat stored in the high O<sub>2</sub> mixture is often spoiled by bacteria like *Brochothrix thermosphacta* and pseudomonads (Gill, 1996). In MAs with high concentrations of O<sub>2</sub>, the meat normally maintains its bright red oxymyoglobin colour for 4 - 7 days before the colour starts deteriorating into grey/brown due to formation of metmyoglobin (Sørheim et al., 1999). This length of time is often not considered sufficient for displaying and selling the product to 4.5 mill. inhabitants all along the distance from Kristiansand in the south to Kirkenes in the north of Norway (about 2700 km) corresponding to the distance from Oslo to Rome (about 2600 km)!

Using high CO<sub>2</sub>, MAs of either CO<sub>2</sub> alone or CO<sub>2</sub>/N<sub>2</sub> with up to 70% CO<sub>2</sub> increases the microbiological shelf life of the meat compared to MAs of high O<sub>2</sub>. The absence of O<sub>2</sub> combined with the presence of CO<sub>2</sub> retard the microbiological growth. Unfortunately, the colour of the meat in MAs of CO<sub>2</sub> is less satisfactory, either as purple deoxymyoglobin or as grey/brown metmyoglobin. The meat inevitably discolours when concentrations of O<sub>2</sub> are low. Tolerance levels for avoiding metmyoglobin formation are less than 0.1% O<sub>2</sub> for beef (Gill and McGinnis, 1995) and 0.5% O<sub>2</sub> for pork (Sørheim et al., 1997b). These low O<sub>2</sub> levels, particularly for beef, are difficult to achieve in most commercial packaging operations, because small fractions of air will be

incorporated in the MAs of the packages. MAs with high CO<sub>2</sub> concentrations seem to be useful for retail packaging when combined with low concentrations of CO for stabilisation of myoglobin and the meat colour.

Vacuum is a packaging method commonly used for bulk storage, transportation and export of meat. However, vacuum has not been a successful method for retail packaging of meat, because of the purple deoxymyoglobin colour of the meat and the visible exudate that occurs in the packages (Bruce, 1990; Gill, 1996). Meat packaged in vacuum can not be presented in the bright red oxymyoglobin state, which depends on the presence of high concentrations of O<sub>2</sub> (Gill, 1996; Taylor et al., 1990), or alternatively as cherry red carboxymyoglobin with CO included in the MA.

#### Hazard for workers:

One of the objections raised against CO as a component of a packaging gas is the potential hazard it might represent for the workers in the meat plants. Using pure CO for mixing in the plant would certainly be such a risk, however, CO is delivered as a premixture (DNC 29.7 - 0.3) or in a 1% mixture together with 99% N<sub>2</sub> (Pakkemix NC1), which is the practice of gas suppliers to the Norwegian meat industry. This way of supplying CO is recognised to be a very safe handling procedure by the health authorities. MAs with concentrations of 60 - 70% O<sub>2</sub> must be handled carefully, because they are explosive gas mixtures. Strict safety regulations apply to explosive gas mixtures, resulting in additional costs of equipment and packaging operations. The benefit of the CO mixture is that it carries no risk or handling costs due to risk of explosion.

#### Experience of the Norwegian meat industry:

Despite the long term knowledge of CO and its many positive properties as a component of MAs for meat, the CO mixture has not been adopted to any large extent by the global meat industry. In many countries, like the US and EU, CO is presently not permitted for use in MAP of meat (Cornforth, 1994; European Parliament and Council Directive, 1995). However, the Norwegian food control authority has derogated from the EU directive for a two years period. Accordingly, the Norwegian meat industry might use CO as a component of a packaging gas in concentrations up to 0.5% until October 1, 2000. The high CO<sub>2</sub>/low CO mixture is the only MAP which provides a shelf life sufficient for displaying and selling fresh retail meat products in all parts of Norway. The Norwegian meat industry started to use the high CO<sub>2</sub>/low CO mixture in packaging of fresh retail meat products in the mid-eighties. The market share of retail meat packaged in the high CO<sub>2</sub>/low CO mixture in Norway is currently estimated at 50 - 60% (ground beef as high as 85%). The Norwegian food control authority has not registered outbreaks or a higher frequency of sporadic cases of food borne diseases linked to such products (The Scientific Committee, under The Norwegian Food Control Authority, 19.4.99).

#### Support from the European meat industry:

The meat industry in Sweden has also discovered the benefits and advantages of the high CO<sub>2</sub>/low CO mixture in packaging of fresh meat. Both Swedish Meats (which is the organisation of the Swedish meat cooperative) and the Swedish Meat Trade Association (which is the organisation of the private meat industry in Sweden) support the Norwegian meat industry's

application to the EU Commission (letters enclosed). Also, the Danish Pig Producers and Slaughterhouses, the Spanish Meat Industry and the Finnish Meat Research Institute support the application (letters enclosed).

- benefit for the consumer:

The high CO<sub>2</sub>/low CO mixture is the only MAP which provides a shelf life sufficient for displaying and selling fresh retail meat products in a large geographical area like Norway.

Food safety and traceability:

The high CO<sub>2</sub>/low CO mixture enables centralised packaging operations with quality control with less risk for cross-contamination than in local butcher shops or by supermarket back-store operations. The Norwegian meat industry already produces pork products traceable in integrated systems back to the farm and beef products traceable to the individual animal.

The ability of *Y. enterocolitica* to multiply at low temperature is of considerable concern to food producers, particularly in countries like Australia, Canada, Denmark, Germany, New Zealand, Norway and Sweden where *Y. enterocolitica* has surpassed *Shigella* and now rivals *Salmonella* and *Campylobacter* as a cause of acute bacterial gastroenteritis (Nesbakken, 2000). The growth of *Y. enterocolitica* was totally inhibited in ground beef packed in the high CO<sub>2</sub>/low CO mixture both at 4°C and 10°C while it grew fairly well both in the high O<sub>2</sub> mixture and in the chub packs.

At the abusive storage temperature of 10°C *E. coli* O157:H7 in the chub packs grew about as fast as the background flora. However, growth was nearly totally inhibited in the high CO<sub>2</sub>/low CO mixture and in the high O<sub>2</sub> mixture (Nissen et al., 1999; Nissen et al., submitted).

Quality:

Centralised pre-packaging of retail meat in the meat industry is cost-effective compared to on-site packaging in food stores. Self-service food stores and supermarkets often require to be supplied with pre-packaged meat. The long shelf life of meat packaged in the high CO<sub>2</sub>/low CO mixture provides a possibility of a wider selection of fresh meat on display in the stores. If the Norwegian meat industry loses the possibility to use this mixture, food stores in rural and remote areas will have to be supplied by frozen meat, which has a low acceptability of the consumer.

The high CO<sub>2</sub>/low CO mixture provides extended freshness: fresh meat will last for many days (often more than a week) in the consumer's home refrigerator, and the consumer might shop fresh meat once a week, and fresh meat is available 24 hrs a day 7 days a week in hypermarkets, supermarkets, discount stores, service stores; and the consumer might get quality premium brand fresh meat in her/his local discount store. The consumer will find a wider variety of fresh meat products than otherwise possible.

The consumers seem to prefer fresh meat products packaged in the high CO<sub>2</sub>/low CO mixture or the high O<sub>2</sub> mixture compared to other packaging methods (Solheim 1996).

Leakages in packages containing high CO<sub>2</sub>/low CO mixture might be detected by the consumer. The discoloration might be an indicator of leakages for ground beef packed in the high CO<sub>2</sub>/low CO mixture (Sørheim, 1996).

#### Prices:

Industrialised handling with centralised packaging in MAP means lower consumer prices. Waste due to "sell by date" in the distribution chain in Norway (high CO<sub>2</sub>/low CO mixture) is less than 1%, as compared to 2 - 3% in countries using the high O<sub>2</sub> mixture, according to interviews with operators/supermarket chains in UK, The Netherlands and Spain (Dag Hallan, Norwegian Meat Cooperative, personal communication).

#### II.6. Exposure

Carbon monoxide (CO) is a colourless, odourless and tasteless gas. It is produced by incomplete combustion of carbon-containing organic material. The production of CO from natural processes is quite significant. Nevertheless, CO from antropogenic activities is far more important concerning human health, since this formation takes place in heavily polluted areas.

Natural background levels of CO are 0.01 - 0.9 mg/m<sup>3</sup> (0.01 - 0.8 ppm). In urban areas, 8-h mean concentrations of CO are generally < 20 mg/m<sup>3</sup>, but levels exceeding 60 mg/m<sup>3</sup> have been reported (WHO, 1979). Among tobacco smokers, CO from smoking is by far the dominating source of CO exposure (WHO, 1987).

According to WHO experts (WHO, 1979; WHO, 1987), the only way of exposure which is of relevance to human health, is via inhalation of CO gas. Upon absorption from the lungs into the blood, CO forms a strong coordination bond with the iron atom in haemoglobin forming carboxyhaemoglobin (HbCO). The affinity of haemoglobin for CO is roughly 240 times that of its affinity for oxygen. CO is absorbed through the lungs and the concentration of HbCO in the blood will depend on several factors, mainly the concentration of CO in inhalation air, the exposure time and the level of activity of the individual (pulmonary ventilation).

The Norwegian meat industry is using a gas mixture of 60 - 70% CO<sub>2</sub>, 30 - 40% N<sub>2</sub> and 0.3 - 0.4% CO for the packaging of fresh retail meat of beef, pork and lamb. According to Watts et al. (1978) beef which is exposed to an atmosphere containing 1% CO for 3 days result in about 30% saturation of the meat myoglobin. When the meat was cooked (hotplate maintained at 195°C for up to 8 minutes), only 0,1 mg CO remained in the meat per kg resulting in a loss of CO about 85%.

Data are very scarce concerning comparison of CO exposure from air and consumption of CO-treated meat. According to Sørheim et al. (1997a) consumption of 250 g CO-treated meat (after cooking) yield a theoretical intake of maximum 0.025 mg CO, compared with inhalation of 15 mg CO per hour at the acceptance level suggested by Norwegian authorities (giving 1.5% HbCO, including endogenous formation). Even though the estimates are crude, the calculations show without doubt that intake of CO from meat consumption is negligible. Furthermore, absorption of CO from the gastrointestinal tract will be very much lower (if it happens at all), compared with absorption via the lungs.



## II.7. Reaction and fate in food

The main function of low levels of CO in MAs is to give a stable, cherry red colour of the meat through strong binding of CO to myoglobin and formation of carboxymyoglobin (El-Badawi, 1964). Although a substantial increase in the shelf life of meat can be obtained by using various MAs, it is often limited by discolouration due to oxidation of myoglobin to metmyoglobin. This discolouration can be prevented by including a small fraction of CO in the gas mixture. Carboxymyoglobin is more resistant to oxidation than oxymyoglobin, due to the stronger binding of CO to the iron-porphyrin site on the myoglobin molecule (Wolfe, 1980). CO in concentrations of 1 - 5% had the ability to increase metmyoglobin reduction, even in the presence of air (Lanier et al., 1978).

## PART III. TOXICOLOGICAL DATA

### III. 1 - 4.

Health effects of carbon monoxide has been evaluated by two WHO expert groups (WHO, 1979; WHO, 1987). The health effects are associated with the degree of HbCO formation. According to the aforementioned expert groups, the most sensitive individuals should be protected from CO exposures leading to a HbCO level exceeding 2.5%. In healthy adults, no adverse health effects are described at concentrations resulting in HbCO levels < 5%.

A small amount of CO is formed naturally in the human body, from breakdown of haemoproteins. This production leads to a HbCO concentration of about 0.5%. The average HbCO concentration in non-smokers is 1.2 - 1.5%, while the level in smokers usually is 3 - 4%.

The WHO experts (WHO, 1987) recommended a maximum HbCO level of 2.5 - 3% in order to protect the population at large, included sensitive individuals. In order to obtain this, they recommended maximum levels of CO in ambient air which will meet this requirement for different times of exposure:

Maximum levels of 100 mg/m<sup>3</sup> for < 15 min  
Average levels < 60 mg/m<sup>3</sup> for 30 min  
Average levels < 30 mg/m<sup>3</sup> for 1 hour  
Average levels < 10 mg/m<sup>3</sup> for 8 hours

The European Union has not evaluated CO for use as a packaging gas for meat. However, in 1990 (European Commission, 1991), several other gases (carbon dioxide, oxygen, nitrogen, nitrous oxide, hydrogen and argon) were evaluated by the Scientific Committee for Food (SCF) for use as packaging gases and propellants. In this case it was considered unnecessary to adopt ADIs because of general knowledge of their safety in use, and the estimated insignificant intakes compared with exposure from other sources. Furthermore, in 1996, the SCF reviewed the safety of modified and controlled atmosphere packaging again, and placed particular emphasis on the importance of HACCP for the avoidance of microbiological risk in this context (European Commission, 1996). The SCF concluded that it does not see specific hazards for human health by

the use of controlled or MAs, but that a prerequisite is that the principles of HACCP are observed. A similar approach should also be feasible concerning CO used at very low concentrations in mixture with CO<sub>2</sub> and N<sub>2</sub>.

Accordingly, the issue which should be solved concerning health effects of CO used in gas packaging, is the question of preventing the consumers from exposure to meat of unacceptable microbiological quality. Thus, two studies on shelf life, off-odour and colour (Sørheim et al., 1999) and pathogens (Nissen et al., 1999; Nissen et al., submitted) follow as enclosures. Summaries of the two studies are also given in "Part II.5. Justification for the additive - investigations on the efficacy of the substance for the intended effect at the level proposed".

### III.5. Review of results and conclusions

As can be seen from the foregoing, exposure to CO via consumption of meat products treated with a MA containing < 0.5% CO represent a negligible source of CO, and will probably not contribute to any increase in the carboxyhaemoglobin level. From a toxicological point of view, packaging gas with < 0.5% CO presents no threat to human health (Sørheim et al. 1997a). This is in accordance with an assessment performed by members of The Scientific Committee for Food, under The Norwegian Food Control Authority (30.11.98).

## PART IV. SUMMARY DOCUMENT

Gas mixtures with low concentrations of CO and high concentrations of CO<sub>2</sub> provide a combination of a long microbiological shelf life and a stable bright red colour of meat. Meat packaged in a MA with high O<sub>2</sub> can achieve an initial bright red colour, but the microbiological shelf life and the colour stability are considerably lower than those of the CO mixture. Using CO eliminates the need for having O<sub>2</sub> as a component of the MA. Other MAs and packaging methods, like high CO<sub>2</sub> with O<sub>2</sub> absorbers, chub packs and vacuum packs may give a microbiological shelf life similar to that of the high CO<sub>2</sub>/low CO mixture, but with a less acceptable colour or appearance of the meat. Thus, there appears at present to be no fully satisfactory alternatives to the CO mixture used in packaging of retail-ready red meats in Norway (Sørheim et al., 1999).

In an investigation, growth of *E. coli* O157:H7 at 10°C in ground beef was nearly totally inhibited in the high CO<sub>2</sub>/low CO mixture. The prolonged shelf life at 4°C did not increase growth of *L. monocytogenes* in ground beef stored in the high CO<sub>2</sub>/low CO mixture. The growth of *Y. enterocolitica* was totally inhibited in ground beef packed in the high CO<sub>2</sub>/low CO mixture both at 4°C and 10°C while it grew fairly well both in the high O<sub>2</sub> mixture and in the chub packs. However, the observed growth of strains of *Salmonella* both in the high CO<sub>2</sub>/low CO mixture and in chub packs at the abuse temperature of 10°C does emphasise the importance of temperature control during storage (Nissen et al., 1999; Nissen et al., submitted).

From a toxicological point of view, packaging gas with < 0.5% CO presents no threat to human health (Sørheim et al. 1997a). The European Union has not evaluated CO for use as a packaging gas for meat. However, in 1990 (European Commission, 1991), several other gases (carbon dioxide, oxygen, nitrogen, nitrous oxide, hydrogen and argon) were evaluated by the Scientific

Committee for Food (SCF) for use as packaging gases and propellants. In this case it was considered unnecessary to adopt ADIs because of general knowledge of their safety in use, and the estimated insignificant intakes compared with exposure from other sources. A similar approach should also be feasible concerning CO used at very low concentrations in mixture with CO<sub>2</sub> and N<sub>2</sub>.

The Norwegian meat industry started to use the high CO<sub>2</sub>/low CO mixture in packaging of fresh retail meat products in the mid-eighties. The market share of retail meat packaged in the high CO<sub>2</sub>/low CO mixture in Norway is currently estimated at 50 - 60% (ground beef as high as 85%). The Norwegian food control authority has not registered outbreaks or a higher frequency of sporadic cases of food borne diseases linked to such products (The Scientific Committee, under The Norwegian Food Control Authority, 19.4.99).

#### Conclusions:

Gas mixtures with low concentrations of CO, up to 0.5%, and high levels of CO<sub>2</sub>, approximately 70%, have many advantages regarding shelf life, inhibition of pathogenic bacteria like *E. coli* O157 and *Y. enterocolitica*, colour stability, labour safety and costs. CO used as described in these concentrations, does not present any toxic threat to the consumer. Considering the benefits the Norwegian meat industry has experienced with the CO gas mixture over the past decade, this gas mixture should have a potential for a wider application in retail packaging of meat in the EU.

#### REFERENCES

Bruce, J. (1990) A comparison of American and European markets for the MAP of fresh red meats, in Proc. Int. Conf. Modified Atmosphere Packaging (Day, B.P.F., ed.), Stratford-upon-Avon, Part 2, pp. 1 - 14, Campden Food & Drink Res. Assoc., Chipping Campden, UK

Commission of the European Union (CEU) (1981) Food - Science and Technique. Report from the Scientific Committee for Food, 11th Series, p. 6, Brussels, Belgium

Cornforth, D. (1994) Color - its basis and importance, in Quality attributes and their measurement in meat, poultry and fish products. Adv. Meat Res. - Vol. 9 (Pearson, A.M. & Dutson, T.R., eds.), pp. 34 - 78, Blackie

El-Badawi, A.A, Cain, R.F., Samuels, C.E. and Angelmeier, A.F. (1964) Color and pigment stability of packaged refrigerated beef. *Food Technol.* 18, 159-163

European Commission. Reports of the Scientific Committee for Food. 25<sup>th</sup> series, Annex II, pp. 17 - 18, 1991

European Commission, Scientific Committee for Food, Minutes of the 103<sup>rd</sup> meeting, 19 - 20 September 1996, p. 8, 1996

European Parliament and Council Directive (1995) of 20 February 1995 on Food Additives Other than Colours and Sweeteners. Off. J. Eur. Commun. No. L61, 1 - 38, 18.3.95

Gill, C.O. (1988) The solubility of carbon dioxide in meat. Meat Sci. 22, 65 - 71

Gill, C.O. (1996) Extending the storage life of raw chilled meats. Meat Sci. 43 (suppl. issue), S99 - S110

Gill, C.O. and McGinnis, J.C. (1995) The effects of residual oxygen concentration and temperature on the degradation of the colour of beef packaged under oxygen-depleted atmospheres. Meat Sci. 39, 387 - 394

Kropf, D.H. (1980) Effects of retail display conditions on meat color. Proc. Reciprocal Meat Conference 33, 15 - 32

Lanier, T.C., Carpenter, J.A., Toledo, R.T. and Reagan, J.O. (1978) Metmyoglobin reduction in beef as affected by aerobic, anaerobic and carbon monoxide containing environments. J. Food Sci. 43, 1788 - 1792, 1796

Nesbakken, T. *Yersinia species* (2000) In: The microbiological safety and quality of food. Lund, B. M., Baird-Parker, T. C., Gould, G. W. (eds.), Aspen Publishers, Inc., Gaithersburg, MD, USA, pp. 1363-1393.

Nissen, H., Alvseike, O., Bredholt, S., Holck, A. and Nesbakken, T. (1999) Packaging of ground beef in an atmosphere with low carbon monoxide and high carbon dioxide restrains growth of *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Yersinia enterocolitica* and *Salmonella diarizonae*. In: Tuijtelaars, A.C.J., Samson, R.A., Rombouts, F.M., Notermans, S., (Eds.), Food Microbiology and food safety into the next millenium. Proceedings of the Seventeenth International Conference of the International Committee on Food Microbiology and Hygiene (ICFMH), 13-17 September 1999, Veldhoven, The Netherlands, pp. 285-286.

Nissen, H., Alvseike, O., Bredholt, S., Holck, A. and Nesbakken, T. (submitted) Packaging of ground beef in an atmosphere with high carbon dioxide and low carbon monoxide restrains growth of *Yersinia enterocolitica*, *Listeria monocytogenes* and *Escherichia coli* O157:H7. Int. J. Food Microbiol.

Renner, M. and Labadie, J. (1993) Fresh red meat packaging and meat quality. Proc. Int. Congr. Meat Sci. Technol., 39, 361-387

Solheim, R. (1996) Consumer purchase probability of beef and pork packaged in different atmospheres. Report, Matforsk, 10 pp.

Sørheim, O. (1996) Discoloration of meat as an indicator of leakages in packages containing a CO gas mixture. Report, Matforsk, 5 pp.

Sørheim, O., Aune, T. and Nesbakken, T. (1997a) Technological, hygienic and toxicological aspects of carbon monoxide used in modified-atmosphere packaging of meat. *Trends in Food Sci. Technol.* 8, 307 - 312

Sørheim, O., Erlandsen, T., Nissen, H., Lea, P. and Høyem, T. (1997b) Effects of modified atmosphere storage on colour and microbiological shelf life of normal and pale, soft and exudative pork. *Meat Sci.* 47, 147 - 155

Sørheim, O., Nissen, H., and Nesbakken, T. (1999) The storage life of beef and pork packaged in an atmosphere with low carbon monoxide and high carbon dioxide. *Meat Sci.* 52, 157 - 164

Taylor, A.A., Down, N.F. and Shaw, B.G. (1990) A comparison of modified atmosphere and vacuum skin packing for the storage of red meats. *Int. J. Food Sci. Technol.* 25, 98 - 109

Watts, D.A., Wolfe, S. K., and Brown, W. D. (1978). Fate of [ $^{14}\text{C}$ ] carbon monoxide in cooked or stored ground beef samples. *J. Agric. Food Chem.*, 26, 210 - 214

Wolfe, S.K. (1980) Use of CO- and CO<sub>2</sub>-enriched atmospheres for meats, fish, and produce. *Food Technol.* 34 (3), 55 - 63

WHO (1979) Environmental health criteria. 13. Carbon monoxide. World Health Organization, Geneva, Switzerland

WHO (1987) Air quality guidelines for Europe. World Health Organization Regional Publ., European Series, no. 23

1 PACKAGING OF GROUND BEEF IN AN ATMOSPHERE WITH HIGH  
2 CARBON DIOXIDE AND LOW CARBON MONOXIDE RESTRAINS  
3 GROWTH OF *YERSINIA ENTEROCOLITICA*, *LISTERIA*  
4 *MONOCYTOGENES* AND *ESCHERICHIA COLI* O157:H7

5

6 H. Nissen\*, O. Alvseike<sup>1,2</sup>, S. Bredholt, A. Holck and T. Nesbakken<sup>2</sup>

7

8 MATFORSK, Norwegian Food Research Institute, Oslovn. 1, N-1430 Ås, Norway

9

10 <sup>1</sup>The Norwegian School of Veterinary Science, P.O. Box 8146 Dep., 0033 Oslo,

11 Norway

12 <sup>2</sup>Norwegian Meat Research Centre, P.O. Box 396, 0513 Oslo, Norway

13

14 Corresponding author and request for reprints: Hilde Nissen, MATFORSK,

15 Norwegian Food Research Institute, Oslovn. 1, N-1430 Ås, Norway

16 Tel: +47 64970100

17 Fax: +47 64970333

18 E-mail: [hilde.nissen@matforsk.no](mailto:hilde.nissen@matforsk.no)

## 1 Abstract

2 Growth of the pathogens *Yersinia enterocolitica*, *Listeria monocytogenes*,  
3 *Escherichia coli* O157:H7 and strains of *Salmonella* was compared in ground beef  
4 packed in modified atmospheres of 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub> /0.4 % CO (high CO<sub>2</sub>/ low  
5 CO mixture), 70 % O<sub>2</sub>/ 30 % CO<sub>2</sub> (high O<sub>2</sub> mixture) and in chub packs. The ground  
6 beef was inoculated with rifampicin-resistant or nalidixic acid/streptomycin-resistant  
7 strains (final concentration 10<sup>2</sup>-10<sup>3</sup> bacteria/g) and stored at 4 and 10 °C for up to 14  
8 days. At 4 °C the shelf life based on stable colour and reduced background flora was  
9 prolonged for the high CO<sub>2</sub>/ low CO mixture compared to the two other packaging  
10 methods, but at 10 °C the shelf life was < 8 days for all the packaging methods.  
11 Growth of *Y. enterocolitica* was nearly totally inhibited both at 4 and 10 °C in the high  
12 CO<sub>2</sub>/ low CO mixture, while the bacterial numbers in the samples packed in the high  
13 O<sub>2</sub> mixture increased from about 5x10<sup>2</sup> bacteria/g at day 0 to about 10<sup>4</sup> at day 5 at  
14 4°C and to 10<sup>5</sup> at 10°C. Growth in the chub packs was even higher. *Listeria*  
15 *monocytogenes* showed very little growth at 4 °C in all treatments. At 10 °C there  
16 was slow growth from about 5x10<sup>2</sup> bacteria/g to about 10<sup>4</sup> at day 5 in the high CO<sub>2</sub>/  
17 low CO mixture, while the numbers in the high O<sub>2</sub> mixture and the chub packs were  
18 about 10 times higher. Growth of *E. coli* O157:H7 at 10 °C in the ground beef was  
19 nearly totally inhibited in both the high CO<sub>2</sub>/ low CO mixture and the high O<sub>2</sub> mixture.  
20 Growth in the chub packs was higher, reaching 10<sup>5</sup> bacteria/g on day 5. The  
21 *Salmonella* strains (*S. Typhimurium*, *S. Dublin*, *S. Enteritidis* and *S. enterica*  
22 61:k:1,5,(7)) in the ground meat stored at 10 °C for 5 and 7 days grew to a higher  
23 number in the high CO<sub>2</sub>/ low CO mixture than in the high O<sub>2</sub> mixture. This study  
24 shows that the prolonged shelf life at 4 °C did not increase growth of *Y. enterocolitica*

000076

1 and *L. monocytogenes* in ground beef stored in the high CO<sub>2</sub>/ low CO mixture  
2 mixture, but the observed growth of strains of salmonella at 10 °C in this mixture and  
3 in chub packs does emphasise the importance of temperature control during storage.

4

5 **Keywords:**

6 Ground beef, modified atmosphere packaging, high CO<sub>2</sub>, carbon  
7 monoxide, *Yersinia enterocolitica*, *Listeria monocytogenes*, *Escherichia coli*  
8 O157:H7.

9

000077



## 1. Introduction

Ground beef for retail sale is most often ready-packed in modified atmospheres (MA) or in chub packs. MA-packed ground beef has a longer microbiological shelf life and also maintains an attractive red colour. For the past decade the Norwegian meat industry has been using a gas mixture of 60-70 % CO<sub>2</sub>, 30-40 % N<sub>2</sub>, 0.3-0.5 % CO. (The CO comes ready mixed in the N<sub>2</sub> from the supplier.) The reason for adding CO to the gas mixture is that it will produce a long-lasting cherry-red colour of the meat (Sørheim et al., 1999), but the low concentration of CO has little effect on the microflora of the meat (Clark et al., 1976; Gee and Brown, 1978; Luno et al., 1998). The use of CO at such low concentrations does not present any toxic threat to the consumers (Sørheim et al., 1997). The most commonly used gas mixture for retail-ready meat in other European countries is 70 % O<sub>2</sub>/30 % CO<sub>2</sub> (Gill, 1996). The high oxygen concentration is needed to keep the red colour of the meat (Lambert et al., 1991). It is therefore only possible to obtain half the CO<sub>2</sub> concentration used in the high CO<sub>2</sub>/ low CO mixture. The microbiological shelf life of the high O<sub>2</sub> mixture will be longer than in air, but less than in the high CO<sub>2</sub>/ low CO gas mixture (Sørheim et al., 1999).

The inclusion of CO is controversial because the stable cherry-red colour can last beyond the microbiological shelf life of the meat and thus mask spoilage (Kropf, 1980). The extended shelf life obtained by MAP may under some conditions imply increased risk of growth of pathogens (Silliker and Wolfe, 1980; Hintlian and Hotchkiss, 1986; Farber, 1991; Lamberts et al., 1991). This issue has also been discussed by the European Commission (1997).

However, even if meat packed in high CO<sub>2</sub>/ low CO mixture acquires a stable colour, the shelf life based on odour is significantly longer in the high CO<sub>2</sub>/ low CO

1 mixture only at 4 °C (Sørheim et al., 1999). At this temperature *Yersinia enterocolitica*  
2 and *Listeria monocytogenes* are considered to be the most serious pathogens in  
3 meat. At abuse temperatures (>8 °C) *Escherichia coli* O157:H7 and *Salmonella* spp.  
4 also may grow and increase the health risk to the consumers. In the present study  
5 we wanted to compare growth of these pathogens in ground beef packed in a  
6 commercial Norwegian 60 % CO<sub>2</sub>/40 % N<sub>2</sub>/0.4 % CO (high CO<sub>2</sub>/low CO mixture) with  
7 growth in a high O<sub>2</sub> (70 % O<sub>2</sub>/30 % CO<sub>2</sub>) gas mixture and in ground beef in chub  
8 packs during storage at 4 and 10 °C in order to evaluate the microbiological safety of  
9 the product.

10

## 11 2. Materials and methods

### 12 2.1. Preparation and packaging of the ground beef

13 The beef carcasses were de-boned, and trimmings with 14 % fat were ground  
14 through a 4 mm plate. The batch of ground beef was divided into 500 g portions  
15 which were packaged in 0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub> (high CO<sub>2</sub>/ low CO mixture),  
16 70 % O<sub>2</sub>/ 30 % CO<sub>2</sub> (high O<sub>2</sub>) or packed in clipped chub packs. The beef was packed  
17 at a commercial meat plant within 1 hour of grinding as described by Sørheim et al.  
18 (1999).

19

### 20 2.2. Bacterial cultures and growth conditions

21 Strains of the following pathogens were inoculated in the ground beef: *Yersinia*  
22 *enterocolitica* (mixture of 3 strains), *Listeria monocytogenes* (mixture of 3 strains  
23 isolated from cooked sausage, Blom et al., 1997, Nissen and Holck, 1999),  
24 *Escherichia coli* O157:H7, NCTC 1200 (National Collection of Type Cultures,

000079

1 Colindale, London), non-toxic strain (resistant to 100 µg/ml nalidixic acid and 1000  
2 µg/ml streptomycin) and *Salmonella enterica* subspecies *diarizonae* serovar  
3 61:k:1,5,(7) (*S. enterica* 61:k:1,5,(7)), mixture of 3 strains (National Institute of Public  
4 Health, Oslo). The listeria and yersinia strains were made resistant to rifampicin by  
5 spreading 0.1 ml of overnight cultures onto agar plates of TSB medium (Oxoid, CM  
6 129) containing 50 µg/ml rifampicin (Sigma, St.Louis, MO, USA). The growth rates of  
7 the resistant strains were practically equal to those of the parent strains when tested  
8 in TSB medium in a Bioscreen instrument (Labsystem Co., Helsinki, Finland) at the  
9 same temperature, pH and  $a_w$  (NaCl) concentrations.

10 In a second experiment four rifampicin-resistant salmonella strains, *S.*  
11 Typhimurium, *S. Dublin*, *S. Enteritidis* and *S. enterica* 61:k:1,5,(7) were used to  
12 inoculate the MAP- packed ground beef. The growth rates (measured as above) of  
13 the resistant strains of *S. Enteritidis* and *S. enterica* 61:k:1,5,(7) were essentially the  
14 same as the parent strains while the growth rates of *S. Dublin* and *S. Typhimurium*  
15 were slightly lower.

### 16 17 2.3. Inoculation and storage

18 After packaging the ground beef was inoculated with stationary cultures (the  
19 bacteria were cultivated overnight at 30°C and kept in the refrigerator for 1 day  
20 before use) of the different pathogenic bacteria. The stock cultures were diluted in  
21 peptone water (PW) (Bacto peptone, Difco, 1g/l; NaCl, Merck, 8.5 g/l) and the strains  
22 belonging to the same species or serovars were mixed. Fifty µl of each pathogen  
23 were inoculated with a syringe through a gas probe self-sealing tape (Toray  
24 Engineering Co. Ltd, England) into one of the corners of the MA packages. The  
25 packages thus had one pathogen inoculated in each corner. In the chub packs the

1 pathogens were inoculated at least 3 cm apart. Packages inoculated only with *Y.*  
2 *enterocolitica* and *L. monocytogenes* only were stored at 4°C and analysed after 0, 2,  
3 5, 8 and 14 days while packages inoculated with all 4 pathogens were stored at  
4 10 °C and analysed after 0, 2, 5 and 8 days.

5 In the second experiment four serovars of «*Salmonella*» were inoculated in one  
6 corner each of the package of ground beef and which was stored at 10 °C and  
7 analysed after 0, 2, 5 and 7 days. Non-inoculated packages used as controls were  
8 also stored at 10 °C.

#### 9 10 2.4. Microbial analyses

11 Samples of 25 g ground beef containing the inoculated pathogens were  
12 transferred to a stomacher bag and mixed with 150 ml peptone water (8.5 g NaCl,  
13 1.0 g peptone/1000 ml water). One hundred µl of a ten-fold dilution series were  
14 plated on blood agar containing 50 µg/ml rifampicin for *L. monocytogenes* and *Y.*  
15 *enterocolitica* or 100 µg/ml nalidixic acid and 1000 µg/ml streptomycin sulphate for *E.*  
16 *coli* O157:H7. From the undiluted mixture an aliquot of 1 ml was also plated out. For  
17 enumeration of *Salmonella* spp. the selective medium Brilliant Green Agar (modified)  
18 (BGA; Oxoid, Basingstoke, Hampshire, England) was used. The colonies were  
19 confirmed on Triple Sugar Iron Agar (TSI; Difco, Detroit, MI,) and Urea agar (Urea  
20 Agar Base, Oxoid CM53 and Urea Solution, Oxoid SR20) followed by agglutination  
21 by monovalent antisera (provided by the National Institute of Public Health). In the  
22 second experiment, samples for detection of the four salmonella strains were plated  
23 on blood agar containing 50 µg/ml rifampicin samples from non-inoculated packages  
24 were treated the same way and plated on MRS plates (CM359, Oxoid ), pH 5.7, for  
25 determination of lactic acid bacteria and PCA (Difco, Detroit, MI, USA) plates for total

1 counts of bacteria. The plates were incubated at 30°C for up to 2 days, all  
2 aerobically. On each sampling date the packs with MA were analysed for O<sub>2</sub> and CO<sub>2</sub>  
3 and the pH for all samples was measured in the stomacher solution. Samples from  
4 two replicate packages were used for all analyses, except after 7 days storage in  
5 experiment 2 where three replicate packages were analysed.

## 6 7 *2.5. Statistical analyses*

8 Microbial data were subjected to analysis of variance (ANOVA) and Tukey's  
9 pairwise comparisons. It was deemed appropriate to perform ANOVA on these data  
10 after a log<sub>10</sub> transformation, thereby obtaining a distribution more akin to the normal  
11 distribution on which ANOVA is based.

## 12 13 **3. Results**

14 As expected the shelf life of the ground beef stored at 4 °C was prolonged in the  
15 high CO<sub>2</sub>/ low CO mixture compared with the other packaging methods. This was due  
16 to the stable colour and reduced background flora resulting in little off-odour.  
17 Thus the ground beef packed in the high CO<sub>2</sub>/ low CO mixture still had an acceptable  
18 smell after 14 days of storage at 4 °C, while the beef packed in high O<sub>2</sub> mixture and  
19 in the chub packs had some off-odours. The difference in shelf life was less at 10 °C.  
20 After 5 days storage the ground beef packed in the high CO<sub>2</sub>/ low CO mixture had an  
21 acceptable smell (except the packages inoculated with salmonella, while beef packed  
22 in the high O<sub>2</sub> mixture and the chub packs had a slight off-odour.

23 After 8 days storage there was a strong off-odour for all treatments, but the ground  
24 beef in the high CO<sub>2</sub>/ low CO mixture still looked bright red, in accordance with  
25 Sørheim et al. (1999). The O<sub>2</sub> content in the high CO<sub>2</sub>/ low CO mixture was virtually

1 zero throughout storage at both temperatures. At 10 °C the O<sub>2</sub> content in the high O<sub>2</sub>  
2 gas mixture decreased from 70 to about 35 % after 8 days storage, probably due to  
3 aerobic bacterial metabolism. The chub packs had an O<sub>2</sub>-permeable casing which  
4 probably was the cause of the high bacterial growth in these packs at both  
5 temperatures.

6 Growth of *Y. enterocolitica* was totally inhibited both at 4 and 10 °C in the high  
7 CO<sub>2</sub>/ low CO mixture (Fig. 1a and b), while the number in the samples packed in the  
8 high O<sub>2</sub> mixture increased from about 5x10<sup>2</sup> cfu/g at day 0 to about 10<sup>4</sup> cfu/g at day 5  
9 at 4 °C and to 10<sup>5</sup> cfu/g at 10°C. Growth in the chub packs at 4 °C was even higher  
10 than in the other treatments. Growth in chub packs was also higher than in high O<sub>2</sub> at  
11 10 °C (p=0.007). *L. monocytogens* (Fig. 2a) showed very little growth at 4 °C in all  
12 treatments. At 10 °C (Fig. 2b) there was slow growth (from about 5x10<sup>3</sup> bacteria/g to  
13 about 10<sup>4</sup> at day 5) in the high CO<sub>2</sub>/ low CO mixture. This was more than 10-fold  
14 higher cfu/g at day 5 than in the high O<sub>2</sub> mixture (p= 0.040) and the chub packs  
15 (p=0.035). Ground beef inoculated with *E. coli* O157:H7 and strains of salmonella  
16 was stored at 10°C. Growth of *E. coli* O157:H7 was slow both in the high CO<sub>2</sub>/ low  
17 CO mixture and the high O<sub>2</sub> mixture (Fig. 3) and the numbers were less than 10<sup>4</sup>  
18 cfu/g at day 5. Growth in the chub packs was greater than in the high CO<sub>2</sub>/ low CO-  
19 mixture (p=0.011) and in the high O<sub>2</sub> mixture (p=0.019), reaching 10<sup>5</sup> cfu/g. Growth of  
20 lactic acid bacteria in the non-inoculated packages was somewhat inhibited in the  
21 high CO<sub>2</sub>/ low CO mixture, especially at 4 °C (Fig. 4). At start of the experiment the  
22 pH in the ground beef was about 5.8 in all packages. After 5 days storage the pH  
23 was about 5.7 in the high CO<sub>2</sub>/ low CO mixture, 5.5 in the high O<sub>2</sub> mixture and 5.3 in  
24 the chub packs.

1 Due to growth of other bacteria on the selective plates, only approximate numbers  
2 of *S. enterica* 61:k:1,5,(7) were obtained, but growth of about 1.5 log units was  
3 observed both in the CO mixture and the chub packs (results not shown). This  
4 increase was not seen in the high O<sub>2</sub> mixture. To verify these results and check  
5 whether they were valid for other serovars more virulent to humans, such as *S.*  
6 Typhimurium, *S. Dublin* and *S. Enteritidis*, a second experiment was performed. The  
7 results (Fig. 5 a, b, c and d) show that after 2 days of storage at 10 °C there was  
8 essentially no growth of the salmonella strains in ground beef packed in the high  
9 CO<sub>2</sub>/ low CO mixture and in the high O<sub>2</sub> mixture, while the numbers of salmonella in  
10 the chub packs were about 10 fold higher. After 5 days there was a slight off-odour in  
11 all the packages except for one package with high CO<sub>2</sub>/ low CO mixture which  
12 smelled strongly of H<sub>2</sub>S. In this package the numbers of all the salmonella strains  
13 were higher than in the replicate package and were of the same magnitude as the  
14 numbers in the chub packs. In the O<sub>2</sub> mixture there was no growth of *S. Dublin* and  
15 *S. Enteritidis* and only a low growth of *S. enterica* 61:k:1,5,(7) and *S. Typhimurium*.  
16 The growth of the salmonella strains was still greatly inhibited in the high O<sub>2</sub> mixture,  
17 while growth in the high CO<sub>2</sub>/ low CO mixture was just as high or even higher than in  
18 the chub packs.

19 In the non-inoculated packages the lactic acid bacteria rapidly constituted most of  
20 the background flora (not shown). After 5 days storage the numbers were higher in  
21 the chub-packed samples, but after 8 days there were no obvious differences  
22 (Fig. 6). The pH in the non-inoculated ground beef followed the same pattern as in  
23 experiment 1.

24

#### 1 4. Discussion and Conclusions

2 Ground beef is a high-risk product because pathogens may be mixed into the  
3 ground product which may not be sufficiently heated before consumption. To inhibit  
4 growth of spoilage bacteria and increase shelf life, MAP is often used by retailers.  
5 The question «Do modified atmospheres enhance risk to the consumers health, but  
6 delay signs of spoilage» raised by Hintlian and Hotchkiss (1986) is therefore relevant.  
7 When evaluating the safety of ground beef in the high CO<sub>2</sub>/ low CO mixture  
8 compared to other commercially available packaging methods, we have focused on  
9 bacteria that show good growth below 10 °C and are most relevant for meat  
10 products.

11 The ability of *Y. enterocolitica* to multiply at low temperature is of considerable  
12 concern to food producers, particularly in countries like Australia, Canada, Denmark,  
13 Germany, New Zealand, Norway and Sweden where *Y. enterocolitica* has surpassed  
14 *Shigella* and now rivals *Salmonella* and *Campylobacter* as a cause of acute bacterial  
15 gastroenteritis (Nesbakken, 1999). In our study, growth of *Yersinia enterocolitica* was  
16 totally inhibited in ground beef packed in the high CO<sub>2</sub>/ low CO mixture even at 10 °C  
17 while it grew fairly well both in the high O<sub>2</sub> mixture and in the chub packs. Manui-  
18 Tawiah et al. (1993) found that pork shops packed in different MA with 20 or 40 %  
19 CO<sub>2</sub> with or without O<sub>2</sub> allowed growth of *Yersinia enterocolitica*, but here the CO<sub>2</sub>  
20 concentration was lower than in the high CO<sub>2</sub>/ low CO mixture (60 %) used in our  
21 study.

22 *Listeria monocytogenes* is also a pathogen that grows well at low temperatures,  
23 but in our study there was no growth of this bacterium in the ground beef in any of  
24 the packages at 4 °C, and only slow growth at 10 °C. This agrees with results of



1 Farber and Daley (1994) who found no growth of *L. monocytogenes* in different meat  
2 products when stored at 4 °C.

3 At the abusive storage temperature of 10 °C, *E. coli* O157:H7 in the chub packs  
4 grew about as fast as the background flora. However, growth was nearly totally  
5 inhibited in the high CO<sub>2</sub>/ low CO mixture and in the high O<sub>2</sub> mixture. This is in  
6 accordance with the predictive model of Sutherland et al. (1997). Their study showed  
7 that *E. coli* O157:H7 is relatively tolerant of CO<sub>2</sub>, but growth could be inhibited at  
8 10 °C at high CO<sub>2</sub> concentrations and pH < 6.0.

9 In our study, growth of *Salmonella* spp. was not inhibited in ground beef packed in  
10 high CO<sub>2</sub>/ low CO mixture and stored at 10 °C, contrary to what is found in many  
11 other studies (e.g. D'Aoust, 1991). Although salmonella may grow well and out-  
12 compete the background flora on fresh meat stored at 10 °C (Alford and Palumbo,  
13 1969; Mackey and Kerridge, 1988), most reports claim that growth will be inhibited in  
14 MAP at this temperature (Siliker and Wolfe, 1980; D'Aoust, 1991; Gill and DeLacy,  
15 1991). Nychas and Tasson (1996) found that high CO<sub>2</sub> atmospheres were more  
16 inhibitory for growth of *S. Enteritidis* on fresh poultry at 10 °C than were high O<sub>2</sub>  
17 atmospheres, the opposite of what we found for ground beef. Inhibition of bacterial  
18 growth may, however, be influenced by pH, texture and the composition of the  
19 product, and Gill and DeLacy (1991) did find growth of *S. Typhimurium* in high-pH  
20 beef packed in CO<sub>2</sub> and stored at 10 °C. Oxidative stress reactions in salmonella  
21 have recently been reported (Stephen et al., 1999). This may explain the inhibition of  
22 growth (longer lag phase) in the high O<sub>2</sub> mixture in our study.

23 The present study shows that the prolonged shelf life (due to stable colour and  
24 reduced background flora) at 4 °C did not increase the risk of growth of *Y.*  
25 *enterocolitica* and *L. monocytogenes* in ground beef stored in the high CO<sub>2</sub>/ low CO

1 gas mixture. This is probably due to the high CO<sub>2</sub> concentration that is inhibitory to  
2 most microorganisms (Dixon and Kell, 1989). Even at the abusive temperature of  
3 10 °C, the numbers of pathogens at the end of the shelf life (5 days) were less or the  
4 same as were found in the chub packs. The observed growth of salmonella in the CO  
5 mixture and chub packs does however emphasise the importance of temperature  
6 control during storage. There is a wide range of temperature criteria for chilled foods  
7 at retail in European countries. The values range from -1 °C to 10 °C, with most  
8 temperatures being between 4 and 8 °C (European Commission, 1996). These  
9 aspects should also be considered together with the conclusions of the EU report  
10 (European Commission, 1997) which state that MAP has proven to enhance the  
11 product quality by inhibiting the spoilage bacteria. MAP may also constitute a hurdle  
12 to the growth of some pathogens, and the safety of MAP products are mostly  
13 threatened by temperature abuse.

1    **Aknowledgements** We thank Oddvin Sørheim, MATFORSK, Vestfold-Buskerud-  
2    Telemark Slakteri A/L, Sem and Åse Spangelo, Hydrogas AS Utviklingssenter,  
3    Porsgrunn for assistance in the production and packing of the ground meat. We also  
4    thank Janina Berg, Birgitta Baardsen and Ingvild Rosshaug for able technical  
5    assistance and Per Lea for statistics. The Agricultural Food Research Foundation  
6    and "Omsetningsrådet" financially supported the work.

7

#### 8    **References**

- 9    Alford, J.A., Palumbo, S.A. (1969) Interaction of salt, pH and temperature on the  
10    growth and survival of salmonellae in ground pork. Appl. Microbiol. 17, 528-532.
- 11    Blom, H., Nerbrink, E., Dainty, R.H., Hagtvedt, T.A., Borch, E., Nissen, H.,  
12    Nesbakken, T. 1997. Addition of 2.5 % lactate and 0.25 % acetate controls growth  
13    of *Listeria monocytogenes* in vacuum packed sensory acceptable served  
14    sausage and cooked ham stored at 4 °C. Int. J. Food Microbiol. 38, 71-76.
- 15    Clark, D.C., Lentz, C.P., Roth, W.D. (1976) Use of carbon monoxide for extending  
16    shelf-life of prepackaged fresh beef. Can. Inst. Food Sci. Technol. J. 9, 114-117.
- 17    D'Aoust, J.Y. (1991) Psychrotrophy and foodborne *Salmonella*. Int. J. Food Microbiol.  
18    13, 207-216.
- 19    Dixon, N.M., Kell, D.B. (1989) A review. The inhibition by CO<sub>2</sub> of the growth and  
20    metabolism of micro-organisms. J. Appl. Bacteriol. 67, 109-136.
- 21    European Commission (1996) Scientific co-operation on questions relating to food.  
22    Task 2.2: Studies relating to temperature control. Directorate-General III,  
23    10.5.1996, pp. 31-33.
- 24    European Commission (1997) Food science and techniques. Reports of the scientific  
25    committee for food (40<sup>th</sup> series) Directorate-General for Industry, pp. 51-62.

- 1 Farber, J.M. (1991) Microbiological aspects of modified-atmosphere packaging  
2 technology – a review. J. Food Protect. 54, 58-70.
- 3 Farber, J.M., Daley, E. (1994) Presence and growth of *Listeria monocytogenes* in  
4 naturally-contaminated meats. Int. J. Food Microbiol. 22, 33-42.
- 5 Gee, D.L., Brown, W.D. (1978) Extension of shelf life in refrigerated ground beef  
6 under an atmosphere containing carbon dioxide and carbon monoxide. J. Agric.  
7 Food Chem. 25, 274-276.
- 8 Gill, C.O., Delacy, K.M. (1991) Growth of *Escherichia coli* and *Salmonella*  
9 *typhimurium* on high-pH beef packed under vacuum or carbon dioxide. Int. J. Food  
10 Microbiol. 13, 21-30.
- 11 Gill, C.O. (1996) Extending the storage life of meats. Meat Sci. 43 (Suppl), S99-  
12 S109.
- 13 Hintlian, C.B., Hotchkiss J.H. (1986) The safety of modified atmosphere packaging: a  
14 review. Food Technol. 40, 70-76.
- 15 Kropf, D.H. (1980). Effects of retail display conditions on meat color. Proc. of the  
16 Reprocial Meat Conference, 33, 15-32.
- 17 Lambert, A.D., Smith, J.P., Dodds, K.L. (1991) Shelf life extension and  
18 microbiological safety of fresh meat – a review. Food Microbiol. 8, 267-297.
- 19 Luno, M., Beltran, J.A., Roncales, P. (1998) Shelf-life extension and colour  
20 stabilisation of beef packaged in a low O<sub>2</sub> atmosphere containing CO: loin steaks  
21 and ground meat. Meat Sci. 48, 75-84.
- 22 Mackey, B.M., Kerridge, A.L. (1988) The effect of incubation temperature and  
23 inoculum size on growth of salmonellae in minced beef. Int. J. Food Microbiol. 6,  
24 57-65.

000089

- 1 Manu-Tawiah, W., Myers, D. J., Olson, D.G., Molins, R.A. (1993) Survival and growth  
2 of *Listeria monocytogenes* and *Yersinia enterocolitica* in pork chops packaged  
3 under modified gas atmospheres. J. Food Sci. 58, 475-479.
- 4 Nesbakken, T. (1999) *Yersinia* spp. In: Lund, B., Baird-Parker, T., Gould, G.  
5 (Eds.), The microbiological safety and quality of food, in press.
- 6 Nissen, H., Holck, A.L. (1998) Survival of *Escherichia coli* O157:H7, *Listeria*  
7 *monocytogenes*, and *Salmonella kentucky* in Norwegian fermented, dry sausage.  
8 Food Microbiol. 15, 273-279.
- 9 Nychas, G.-J.E., Tassou, C.C. (1996) Growth/survival of *Salmonella enteritidis* on  
10 fresh poultry and fish stored under vacuum or modified atmosphere. Lett. Appl.  
11 Microbiol. 23, 115-119.
- 12 Silliker, J.H., Wolfe, S.K. (1980) Microbiological safety considerations in controlled-  
13 atmosphere storage of meats. Food Technol. 34, 59-63.
- 14 Stephens, P.J., Druggan, P., Nebe-von Caron, G. The significance of oxidative stress  
15 in food preservation. Food Microbiology and food safety into the next millenium. In:  
16 Tuijelaars, A.C.J., Samson, R.A., Rombouts, F.M., Notermans, S., (Eds.),  
17 Proceedings of the Seventeenth International Conference of the International  
18 Committee on Food Microbiology and Hygiene (ICFMH), 13-17 September 1999,  
19 Veldhoven, The Netherlands, pp. 764-766.
- 20 Sutherland, J.P., Bayliss, A.J., Braxton, D.S. Beumont, A.L. (1997) Predictive  
21 modelling of *Escherichia coli* O157:H7: Inclusion of carbon dioxide as a fourth  
22 factor in a pre-existing model. Int. J. Microbiol. 37, 113-120.
- 23 Sørheim, O., Aune, T., Nesbakken, T. (1997) Technological, hygienic and  
24 toxicological aspects of carbon monoxide used in modified-atmosphere packaging  
25 of meat. Trends Food Sci. Technol. 8, 307-312.

000090

- 1 Sørheim, O., Nissen, H., Nesbakken, T. (1999) The storage life of beef and pork
- 2 packaged in an atmosphere with low carbon and high carbon dioxide. Meat Sci.
- 3 52, 157-164.

000091

1 Fig. 1. Growth of *Yersinia enterocolitica* inoculated in ground beef packed in high  
2 CO<sub>2</sub>/ low CO mixture (0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub>), high O<sub>2</sub> (70 % O<sub>2</sub>/ 30 % CO<sub>2</sub>)  
3 or in chub packs. The ground beef was stored at a, 4 °C or b, 10 °C.

4  
5 Fig. 2. Growth of *Listeria monocytogenes* inoculated in ground beef packed in high  
6 CO<sub>2</sub>/ low CO mixture (0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub>), high O<sub>2</sub> (70 % O<sub>2</sub>/ 30 % CO<sub>2</sub>)  
7 or in chub packs. The ground beef was stored at a, 4 °C or b, 10 °C.

8  
9 Fig. 3. Growth of *Escherichia coli* O157: H7 inoculated in ground beef packed in high  
10 CO<sub>2</sub>/ low CO mixture (0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub>), high O<sub>2</sub> (70 % O<sub>2</sub>/ 30 % CO<sub>2</sub>)  
11 or in chub packs, stored at 10 °C.

12  
13 Fig. 4. Growth of lactic acid bacteria (cfu/g on MRS, pH 5.7) in non-inoculated ground  
14 beef packed in high CO<sub>2</sub>/ low CO mixture (0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub>), high O<sub>2</sub>  
15 (70 % O<sub>2</sub>/ 30 % CO<sub>2</sub>) or in chub packs. The ground beef was stored at a, 4°C or b, 10  
16 °C.

17  
18 Fig. 5. Growth of strains of *Salmonellae* inoculated in ground beef packed in high  
19 CO<sub>2</sub>/ low CO mixture (0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub>), high O<sub>2</sub> (70 % O<sub>2</sub>/ 30 % CO<sub>2</sub>)  
20 or in chub packs, stored 10 °C. a. *S.* Typhimurium b. *S.* Dublin c. *S.* Enteritidis d. *S.*  
21 *enterica* 61:k:1,5,(7).

22  
23 Fig. 6. Growth of lactic acid bacteria (cfu/g on MRS, pH 5.7) in non-inoculated ground  
24 beef packed in high CO<sub>2</sub>/ low CO mixture (0.4 % CO/ 60 % CO<sub>2</sub>/ 40 % N<sub>2</sub>), high O<sub>2</sub>

000092


- 1 (70 % O<sub>2</sub>/ 30 % CO<sub>2</sub>) or in chub packs. The ground beef was stored at a, 4 °C or b,
- 2 10 °C.
- 3

000093




*Y. la*

*Yersinia enterocolitica*, 4°C



h<sub>1</sub> h<sub>2</sub>

# *Yersinia enterocolitica*, 10°C



*Listeria monocytogenes*, 4°C





Fig. 2b







Fig. 1a

# Lactic acid bacteria, 4°C



Lactic acid bacteria, 10°C




Fig. 5a

# Salmonella Typhimurium, 10°C








Fig. 5b

# *Salmonella* Dublin, 10°C




*Salmonella* Enteritidis, 10°C




# Salmonella enterica 61:K:1,5,(7)

11/1/81



11/15

Lactic acid bacteria, 10°C



(2)

# FOOD MICROBIOLOGY AND FOOD SAFETY INTO THE NEXT MILLENNIUM

Proceedings of the  
Seventeenth International Conference of the  
International Committee on  
Food Microbiology and Hygiene (ICFMH)  
Veldhoven, The Netherlands, 13-17 September 1999

Edited by  
A.C.J. Tuijtelzaars<sup>1</sup>  
R.A. Samson<sup>2</sup>  
F.M. Rombouts<sup>3</sup>  
S. Notermans<sup>1</sup>

<sup>1</sup>TNO Nutrition and Food Research Institute, P.O.Box 370, 3700 AJ Zeist, The Netherlands

<sup>2</sup>Centraal Bureau voor Schimmelcultures, P.O.Box 273, 3740 AG Baarn, The Netherlands

<sup>3</sup>Wageningen University and Research Centre, Bomenweg 2, 6703 HD Wageningen,  
The Netherlands

misinterpreted in the absence of storage trials or in trials of short duration such as 14 days. If *Brachytherius thermosphacta* is a problem organism in a processing facility or on a particular type of meat, lysozyme, Chrsin or mixtures of the two could be used to control its growth during refrigerated anoxic storage.

**PACKAGING OF GROUND BEEF IN AN ATMOSPHERE WITH LOW CARBON MONOXIDE AND HIGH CARBON DIOXIDE RESTRAINS GROWTH OF *ESCHERICHIA COLI* O157:H7, *LISTERIA MONOCYTOGENES*, *YERSINIA ENTEROCOLITICA* AND *SALMONELLA DIARIZONAE***

Nissen H., Alvseike O.<sup>1,2</sup>, Bredholt S., Holck A. and Nesbakken T.<sup>2</sup>  
MATFORSK, Norwegian Food Research Institute, Oslovn. 1, N-1430 Ås, Norway,  
<sup>1</sup> Norwegian College of Veterinary Medicine, P.O. Box 8146 Dep., 0033 Oslo  
Norway, Norwegian Meat Research Centre, P.O. Box 396, 0513 Oslo, Norway  
Fax: +47 6497 0333, E-mail: hilde.nissen@matforsk.no

Ground beef for retail sale is most often ready packed in modified atmosphere (MA) or in chub packs. MA packed ground beef prolongs the microbiological shelf life and also maintains an attractive red colour. For the past decade the Norwegian meat industry has been using a gas mixture of 0.3-0.5% CO, 60-70% CO<sub>2</sub> and 30-40% N<sub>2</sub> (the CO comes ready mixed in the N<sub>2</sub>). The reason for adding CO to the gas mixture is that it will produce a long-lasting cherry-red colour of the meat (Sørheim et al 1999). The most commonly used gas mixture for retail-ready meat in other European countries is 70% O<sub>2</sub>/30% CO<sub>2</sub> (Gill 1996). The high oxygen concentration is needed to keep the red colour of the meat. It is therefore only possible to obtain half the CO<sub>2</sub> concentration used in the CO gas mixture. The microbiological shelf life will be longer than in air, but less than in the CO gas mixture (Sørheim et al 1999).

The inclusion of CO is controversial because the stable cherry-red colour can last beyond the microbiological shelf life of the meat and thus mask spoilage (Kropf 1980). However, the consumer is able to evaluate the microbiological conditions of the meat by off-odours and the shelf life based on odour is significantly longer in the CO mixture only at 4°C. Thus, extended shelf life does not necessarily imply an increased risk of growth of pathogens. In the present study we wanted to compare growth of the pathogens *Escherichia coli* O157:H7, *Listeria monocytogenes*, *Yersinia enterocolitica* and *Salmonella diarizonae*, in ground beef packed in a commercial Norwegian 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> mixture with growth in a high O<sub>2</sub> (70% O<sub>2</sub>/30% CO<sub>2</sub>) gas mixture and in ground beef in chub packs during storage at 4 and 10°C.

Commercial packages of ground beef (500 g) stored at 10°C were inoculated with the pathogens *Escherichia coli* O157:H7, *L. monocytogenes*, *Y. enterocolitica* and *S.*

#### Chapter 4: Preservation

*diarizona*, and the ground beef stored at 4°C with *L. monocytogenes* and *Y. enterocolitica*. The inocula of *L. monocytogenes* and *Y. enterocolitica* were cocktails of 3 stationary-phase, rifampicin-resistant strains, the inoculum of *E. coli* O157:H7 was one non-toxic nalidixic/streptomycin resistant strain and that of *S. diarizonae* was a cocktail of 3 strains that were not made antibiotic resistant (plated on selective media for *Salmonella* spp.). Controls of ground beef without inoculated pathogens were stored at both temperatures.

After 5 days storage at 10°C the ground beef packed in the CO mixture had an acceptable smell while beef the packed in the high O<sub>2</sub> mixture and the chub packs had a slight off-odour. After 8 days storage there was a strong off-odour for all the treatments. At 4°C the smell was still acceptable after 14 days of storage in the CO mixture, but the high O<sub>2</sub> mixture and the chub packs had some off-odours. The growth of pathogens was restrained in all samples that had been packed in the gas mixture containing CO. Thus, growth of *Y. enterocolitica* was nearly totally inhibited both at 4 and 10°C, while the number in the samples packed in the high O<sub>2</sub> mixture increased from about  $5 \times 10^2$  bacteria per g at day 0 to about  $10^4$  at day 5 at 4°C and to  $10^5$  at 10°C. The number in the chub packs were even higher. *L. monocytogenes* showed very little growth at 4°C in all of the treatments. At 10°C there was slow growth (from about  $5 \times 10^3$  bacteria/g to about  $10^4$  at day 5) in the CO mixture while the number in the high O<sub>2</sub> mixture and the chub packs were about 10 times higher. Growth of *E. coli* O157:H7 at 10°C storage was slow both in the CO-mixture and the high O<sub>2</sub> mixture. Growth in the chub packs was higher reaching  $10^5$  bacteria/g on day 5. The growth of *S. diarizonae* followed the same pattern as *E. coli* O157:H7.

Ground beef is a high-risk product because pathogens may be mixed into the product which may not be properly heated before being eaten. The present study shows that the reduced background flora of beef packed in the CO mixture did not result in increased growth of the pathogens. This was probably due to the high concentration of CO<sub>2</sub> in this mixture which particularly inhibits Gram negative bacteria. The O<sub>2</sub> content in the CO mixture was virtually zero throughout storage at both temperatures. At 10°C the O<sub>2</sub> content in the high O<sub>2</sub> gas mixture decreased from 70% to about 35% after 8 days, probably due to aerobic bacterial metabolism. The chub packs had air-permeable casing which probably was the cause of the high bacterial growth in these packs.

The conclusion of the present study is that for the conditions studied, the risk of growth of the pathogens *Y. enterocolitica*, *L. monocytogenes*, *E. coli* O157:H7 and *S. diarizonae* in ground beef stored in CO gas mixture is the same as or less than in the ground beef stored in high O<sub>2</sub> or under vacuum (chub packs).

#### References

- Gill, C.O. (1996) Extending the storage life of meats. *Meat Science*, 43 (Suppl), S99-S109.
- Kropf, D.H. (1980). Effects of retail display conditions on meat colour. *Proceedings of the Reciprocal Meat Conference*, 33 15-32.
- Sørheim O., Nissen H., Nesbakken T. (1999) The storage of beef and pork packaged in an atmosphere with low carbon and high carbon dioxide. *Meat Science* 52, 157-164.

English summary

3

## CONSUMER PURCHASE PROBABILITY OF BEEF AND PORK PACKAGED IN DIFFERENT ATMOSPHERES

Ragnhild Solheim, MATFORSK, Norwegian Food Research Institute, Oslovn. 1, 1430 Ås, Norway

Ground beef, beef loin steaks and pork chops were packaged in modified atmospheres of 0.4% CO/ 60% CO<sub>2</sub>/ 40% N<sub>2</sub> (high CO<sub>2</sub>/low CO mixture) and 70% O<sub>2</sub>/ 30% CO<sub>2</sub> (high O<sub>2</sub> mixture). In addition ground beef was packaged in clipped chub packs, beef loin steaks were vacuum packaged, and pork chops were packaged in an atmosphere of 60% CO<sub>2</sub>/ 40% N<sub>2</sub> with each pack containing an O<sub>2</sub> absorber. The purchase probability data were collected by interviewing 126 consumers usually purchasing meat and meat products. The consumers visually compared the samples within each type of meat. The consumers preferred ground beef packaged in the high CO<sub>2</sub>/low CO mixture or the high O<sub>2</sub> mixture compared to ground beef packaged in clipped chub packs. Purchase probability increased when pork chops were packaged in the high CO<sub>2</sub>/low CO mixture. Pork chops in packs containing an O<sub>2</sub> absorber, were rated lowest in purchase probabilities. The purchase probability for beef loin steaks was similar when packaged in the high CO<sub>2</sub>/low CO mixture or the high O<sub>2</sub> mixture, and these products were preferred compared to beef loin steaks packaged in vacuum.

000109



**COMMISSIONED REPORT**

MATFORSK – Norwegian Food  
Research Institute

Osloveien 1, N-1430 Ås  
Tel: (+47) 64 97 01 00

**Report No.:**

O-7224

**Availability:**  
Confidential

<b>Report Title:</b> Consumer Survey of Meat Products	<b>Date:</b> July 11, 1996
<b>Project Manager/Author:</b> Ragnhild Solheim	<b>Signature of Project Manager:</b> [Signature]
<b>Head of Department:</b> Björg Egelanddal	<b>Signature of Head of Department:</b> [Signature]
<b>Department:</b> Analysis Methodology	<b>Project No:</b> O-7224 FBT
<b>Commissioned by:</b> Norsk Kjøtt [Norwegian Meat]	<b>Commissioner's contact:</b> Truls Nesbakken

000110



**Summary/Abstract:**

The consumers' (N=126) purchasing tendency for ground meat, pork chops and top loin of beef packaged in various ways was measured by means of a central location test. The consumers indicated their purchasing tendency on a verbal five-point scale from "Will definitely not buy" to "Will definitely buy." Moreover, the consumers estimated their buying frequency for the three different products, and provided their age and gender.

The consumer population was composed of 62 % women and 38 % men. The age distribution was roughly equivalent for both genders.

*Purchasing Frequency:*

The major segment (47.6 %) of the participants said that they buy ground meat two to three times a month, while 20.6 % buy ground meat once a month.

Pork chops were purchased two to three times a month by 29.4 % of the participants, while 38.1 % bought pork chops once a month. 29 % of the participants bought pork chops less frequently than once a month.

Top loin was purchased less frequently than once a month by 58.1 % of the participants in the study. 18 % said that they buy top loin once a month, and 19.4 % indicated that they buy top loin once a week.

*Purchasing Tendency:*

*Ground meat:* Ground meat packaged with CO gas received the same average score for purchasing tendency as ground meat packaged in O<sub>2</sub> gas, while ground meat packaged as sausage had the lowest score for purchasing tendency.

*Pork Chops:* Pork chops packaged in CO gas received the highest total score for purchasing tendency, while pork chops packaged in O<sub>2</sub> gas received the second highest total score and pork chops packaged with oxygen absorber received the lowest score.

*Top Loin:* Top loin packaged in CO gas and in O<sub>2</sub> gas had roughly the same average score for purchasing tendency, while vacuum-packaged top loin received the lowest score for purchasing tendency.

Excerpts of the report/results shall not be reproduced without prior written permission from MATFORSK. Also see special requirements for the use of MATFORSK's name in marketing.



MATFORSK – Norwegian Food Research Institute

Table of Contents

1. Premise .....	1
2. Implementation .....	1
2.1 Materials and Survey Conditions .....	1
2.2 Method .....	1
2.3 Consumer Population .....	3
3. Results .....	3
3.1 Purchasing Frequency for Meat Products .....	3
3.2 Purchasing Tendency for Meat Products .....	3
4. Conclusion .....	7



MATFORSK – Norwegian Food Research Institute

1. Premise

The consumers' choice of meat products may be attributed to factors such as how the products are presented in their packaging. The consumers' purchasing tendency for meat products packaged in accordance with various principles was therefore measured.

2. Implementation

**2.1 Materials and Survey Conditions**

The survey was taken at Drøbak City [shopping center] on June 12, 1996 (Tuesday) from 10 am to 4 pm.

Ground meat, pork chops and top loin of beef packaged in accordance with various principles (Table 1) was presented to consumers who eat these types of products. The products were delivered to MATFORSK on June 8, 1996 and stored at 4°C until the survey date. Products from the same group were placed side by side on a table under lighting with a strength of approximately 2000 lux (equivalent to the light intensity of a refrigerated meat counter in a store). The products were replaced with cold stored products every three hours.

*Table 1. Packaging methods for meat products tested in consumer survey*

Meat Product	Packaging Method				
	Sausage	CO	O <sub>2</sub>	Vacuum	O <sub>2</sub> w/absorber
Ground Meat	x	x	x		
Pork Chops		x	x		x
Top loin of beef		x	x	x	

x = packaging method used

**2.2 Method**

The products were coded with three-digit random numbers and evaluated in a systematically rotating order. The consumers indicated purchasing probability on a verbal scale and purchasing frequency for the product, and gave their age and gender (Figure 1). Following the evaluation, the verbal scale was translated into numerical values from 1 to 5, where 1=Will definitely not buy, and 5=Will definitely buy. The consumers spent between 5 and 10 minutes answering the questions.



MATFORSK – Norwegian Food Research Institute

Dear Consumer!

We are taking a survey on the consumer opinion concerning a selection of meat products as they appear in the meat counter. We hope that you will take a minute to let the meat producers know your opinion!

What to do:

1. Please take a look at the samples of top loin.
2. Consider whether you would buy these products the way they appear, assuming they are priced the same. Evaluate the products in the order listed below and check one box for each product.

Sample marked 763

Will definitely buy ☐  
May buy ☐  
May/may not buy ☐  
May not buy ☐  
Will definitely not buy ☐

Sample marked 288

Will definitely buy ☐  
May buy ☐  
May/may not buy ☐  
May not buy ☐  
Will definitely not buy ☐

Sample marked 911

Will definitely buy ☐  
May buy ☐  
May/may not buy ☐  
May not buy ☐  
Will definitely not buy ☐

In closing please answer the questions below about your age, gender and how often you buy top loin of beef.

My age (check one):

18-25 years old ☐  
26-35 years old ☐  
36-45 years old ☐  
46-55 years old ☐  
56-65 years old ☐  
over 65 years old ☐

Gender (check one):

Female ☐ Male ☐

I buy top loin of beef (refrigerated):

Less than once a month ☐  
Once a month ☐  
Two to three times a month ☐  
Once a week ☐  
More than once a week ☐

THANK YOU FOR YOUR ASSISTANCE!

Figure 1. Questionnaire for consumer survey of meat products. Corresponding forms were used for pork chops and ground meat.

000114

Commission – Confidential/Rso/IBu July 11, 1996/O-7224 FBT



### 2.3 Consumer Population

124–126 consumers over 18 years of age participated in the survey of the three different meat products. There were a few more women than men among the participants, and there were fewer participants over 56 years of age than in the other age groups (Table 2). The age distribution among men and women was the same.

*Table 2. Age and gender distribution among consumers participating in the survey.*

Meat Product	No. of consumers	Gender (%)		Age (year, % distribution)					
		Women	Men	18–25	26–35	36–45	46–55	56–65	Over 65
Ground Meat	125	61.6	38.4	15.9	19.8	18.3	20.6	11.1	14.3
Pork Chops	126	61.9	38.1	15.1	18.3	19.8	19.8	11.1	15.9
Top Loin	124	63.7	36.3	14.5	21.0	16.9	21.8	12.9	12.9

## 3. Results

### 3.1 Purchasing Frequency for Meat Products

Ground meat was most frequently bought, followed by pork chops, while top loin was rarely bought by the consumers participating in this study (Table 3). There were similar purchasing frequencies in the various age and gender groups.

*Table 3. Purchasing Frequency for Three Different Meat Products*

Purchasing Frequency	Meat Product		
	Ground Meat (N=125)	Pork Chops (N=126)	Top Loin (N=124)
Less than once a month	11.1	29.4	58.1
Once a month	20.6	38.1	18.5
Two to three times a month	47.6	29.4	19.4
Once a week	15.1	3.2	3.2
More than once a week	5.6	0.0	0.8

### 3.2 Purchasing Tendency for Meat Products

A detailed overview of the purchasing tendency is provided in Attachment 1.

All differences described in the following were significant to a degree of 95 %.

000115



MATFORSK – Norwegian Food Research Institute

*Ground Meat*

Ground meat packaged as sausage was least appreciated by the consumers (see Figure 2). Ground meat packaged in CO gas and ground meat packaged in O<sub>2</sub> gas received the same average score for purchasing tendency. This result occurred regardless of consumer age and gender.

Comments from the consumers:

The wrapping film used to pack meat as sausage hides the contents.

[Above bar chart]

Purchasing Tendency for Ground Meat

[Beside bar chart] Average score (N=126) [Bar chart]

[Below bar chart]

As Sausage

CO gas  
Packaging Method

O<sub>2</sub>

*Figure 2. Purchasing tendency for meat products. 1=Will definitely not buy, and 5=Will definitely buy the product as presented in the survey.*

000116

Commission – Confidential/Rso/IBu July 11, 1996/O-7224 FBT



MATFORSK – Norwegian Food Research Institute

*Pork Chops*

Pork chops packaged in CO gas received the highest average score for purchasing tendency, pork chops packaged in O<sub>2</sub> gas received the second highest score and pork chops packaged with oxygen absorber received the lowest average score (see Figure 3). This result occurred regardless of consumer age and gender.

Comments from consumers:

The pork chops packaged with absorber look gray/brown – are they old? expired?

Sample packaged with CO and O<sub>2</sub>: nitrite added?

[Above bar chart]

Purchasing Tendency for Ground Meat

[Beside bar chart]

Average score (N=126)

[Bar chart]

[Below bar chart]

O<sub>2</sub> absorber

CO gas  
Packaging Method

O<sub>2</sub>

*Figure 3. Purchasing tendency for meat products. 1=Will definitely not buy, and 5=Will definitely buy the product as presented in the survey.*

000117

Commission – Confidential/Rso/IBu July 11, 1996/O-7224 FBT





MATFORSK – Norwegian Food Research Institute

*Top Loin*

Top loin packaged in CO gas and in O<sub>2</sub> gas received approximately the same average score for purchasing tendency (see Figure 4). Vacuum packaged top loin received a lower score for purchasing tendency than the two aforementioned samples. Men indicated roughly equivalent purchasing tendencies for the three varieties, while women indicated the highest purchasing tendency for top loin packaged in CO gas, followed by top loin packaged in O<sub>2</sub> gas, and the lowest purchasing tendency for vacuum packaged top loin.

Comments from consumers:

Sample packaged in CO and O<sub>2</sub>: nitrite added?

Sample packaged in CO: "artificial" sides.

Vacuum packaged sample: looks as if it has been squeezed.

[Above bar chart]

Purchasing Tendency for Top Loin

[Beside bar chart] Average score (N=125) [Bar chart]

[Below bar chart]

Vacuum

CO gas  
Packaging Method

O<sub>2</sub>

*Figure 4. Purchasing tendency for meat products. 1=Will definitely not buy and 5=Will definitely buy the product as presented in the survey.*

000118

Commission – Confidential/Rso/IBu July 11, 1996/O-7224 FBT



#### 4. Conclusion

*Ground Meat:* Ground meat packaged in CO gas and ground meat packaged in O<sub>2</sub> gas received the same average score for purchasing tendency, while ground meat packaged as sausage received the lowest score for purchasing tendency.

*Pork Chops:* Pork chops packaged in CO gas received the highest score for purchasing tendency, pork chops packaged in O<sub>2</sub> gas received the second highest total score, while pork chops packaged with oxygen absorber received the lowest score.

*Top Loin:* Top loin packaged in CO gas and in O<sub>2</sub> gas received approximately the same average score for purchasing tendency, and vacuum packaged top loin received the lowest score for purchasing tendency.

#### Comments

This type of survey does not have representative sampling of consumers in terms of the population as a whole or a specific population segment. The make up of the survey represents a model for a purchasing situation. These circumstances must be taken into account when interpreting the results.

000119

Commission – Confidential/Rso/IBu July 11, 1996/O-7224 FBT



Overview of Results from Consumer Study of Meat Products

	<i>Average</i>	<i>Standard Deviation</i>	<i>Median</i>
<i>Ground Meat (N=126)</i>			
As sausage	2.5	1.5	2
CO gas	3.7	1.4	4
O <sub>2</sub>	3.7	1.4	4
<i>Pork Chops (N=126)</i>			
O <sub>2</sub> absorber	1.9	1.3	1
CO gas	4.6	0.7	5
O <sub>2</sub>	3.6	1.4	4
<i>Top Loin (N=125)</i>			
Vacuum	2.9	1.6	3
CO gas	4.2	1.1	4.5
O <sub>2</sub>	4.0	1.1	4

000120



**Prosjektmedarbeidere:**

Ansvarlig for *opplegg og rapport*: Ragnhild Solheim

Ansvarlig for *prøvemottak og lyssetting*: Oddvin Sørheim

Ansvarlig for *praktisk gjennomføring*: Inger Johanne Fjøsne

Øvrige deltakere i gjennomføring:

Asgeir Nilsen  
Janne Haugdahl  
Vibeke Høst  
Oddvin Sørheim

Ansvarlig for *statistisk bearbeiding*: Per Lea

000121

English summary

**DISCOLORATION OF MEAT AS AN INDICATOR OF LEAKAGES IN PACKAGES CONTAINING A CO GAS MIXTURE**

Oddvin Sørheim, MATFORSK, Norwegian Food Research Institute, Oslovn. 1, 1430 Ås, Norway

The aim of the experiment was to study discoloration of meat packaged in a gas mixture of 60 % CO<sub>2</sub>/40 % N<sub>2</sub>/ 0.4 % CO with different concentrations of residual O<sub>2</sub> added. Tests were performed on ground beef with 1 % NaCl, aged beef loin steaks and pork chops. Leakages were simulated by injecting different amounts of air with a syringe into the packages after two days storage. Discoloration of the meat was measured as reduction in a\* (redness) values and evaluated visually. Ground beef had a low tolerance level of residual O<sub>2</sub> because it was discoloured in atmospheres containing more than 1 % O<sub>2</sub>. Beef loin steaks and pork chops were slightly discoloured in more than 2 and 5 % O<sub>2</sub>, respectively. The results suggest that discoloration can be an indicator of leakages for ground beef, but not for beef loin steaks and pork chops.

COMMISSIONED REPORT

MATFORSK – Norwegian Food  
Research Institute

Osloveien 1, N-1430 Ås  
Tel: (+47) 64 97 01 00

Report No.:

O-7224.col

Availability:  
Confidential

<b>Report Title:</b> Meat Discoloration as an Indicator of Leaks in Packaging with CO Gas Mixtures	<b>Date:</b> November 28, 1996
<b>Project Manager/Author:</b> Oddvin Sorheim	<b>Signature of Project Manager:</b> [Signature]
<b>Head of Department:</b> Astrid Nilsson	<b>Signature of Head of Department:</b> [Signature]
<b>Department:</b> Product and Raw Material Science	<b>Project No:</b> O-7224.col
<b>Commissioned by:</b> Norsk Kjøtt [Norwegian Meat]	<b>Commissioner's contact:</b> Truls Nesbakken

**Summary/Abstract:**

Tests were carried out to find the tolerance limits for residual O<sub>2</sub> for discoloration of meat packaged in CO gas mixture with simulated leak. Various concentrations of air were added to the packages of ground meat, top loin and pork chops with a mixture of 60% CO<sub>2</sub> / 40% N<sub>2</sub> / 0.4% CO two days after packaging. Ground meat packaged in gas containing more than 1% O<sub>2</sub> was clearly discolored, while top loin and pork chops, respectively packaged in gas containing more than 2 and 5% O<sub>2</sub> showed only minor discoloration. The results indicate that discoloration can serve as an indicator of leakage for ground meat, but not for top loin and pork chops.

Excerpts of the report/results shall not be reproduced without prior written permission from MATFORSK. Also, see special requirements for use of MATFORSK's name in marketing.

000123



## Meat Discoloration as an Indicator of Leakage in Packages with CO Gas Mixtures

### Purpose

The purpose of the survey was to find the tolerance limits for residual O<sub>2</sub> with regard to discoloration of ground meat, top loin and pork chops packaged in a CO gas mixture.

### Implementation of the Study

Samples of ground meat with 1% NaCl (20 pieces), tenderized top loin of beef (18 pieces) and pork chops (14 pieces) were gas packed on a Ilapak Delta 2000 machine (Ilapak, Switzerland) on a tray in BDF 550 shrink film (Cryovac). The gas mixture consisted of 60% CO<sub>2</sub> / 40% N<sub>2</sub> / 0.4% CO. The samples were stored out of light at 4°C. After two days of storage, air was added to the packages to increase the O<sub>2</sub> content, i.e. a simulated leak. This was done by sucking out the gas in the package and replacing it with air by means of a syringe and a septa. 0-2.0% O<sub>2</sub> was added to the ground meat, 0-3.2% O<sub>2</sub> was added to the top loin, and 0-13.9% O<sub>2</sub> was added to the pork chops, in all cases with a spectrum of O<sub>2</sub> concentration in their respective ranges. After the replacement of gases, the concentrations of O<sub>2</sub> and CO<sub>2</sub> were measured by means of two Toray instruments, type LC 700F and PG-100 (Toray Eng., Japan). The remaining storage time before unwrapping was 2 days for ground meat and 5 days for both top loin and pork chops.

Upon unwrapping, the O<sub>2</sub> and CO<sub>2</sub> concentrations in the packages were once again measured. Two judges then visually judged the color of the unopened packages according to a scale (1=fresh meat red, 2=dark red, 3=somewhat discolored, 4=moderately discolored, 5=extremely discolored). The packages were then opened, and the color was measured with a Minolta Chroma Meter CR-300 (Minolta Camera Co., Japan) directly on the surface of the meat within 1 minute of opening. The instrument had light source D<sub>65</sub> at 8 mm aperture, and the color was measured in CIE (1976) L\* (luminosity), a\* (redness) and b\* (yellowness). Lastly, the pH was measured directly in the meat with a Ingold Xerolyt electrode (Mettler-Toledo A.G., Switzerland).

### Results and Discussion

The correlation between discoloration upon unwrapping and O<sub>2</sub> concentration when replacing the gas proved to best be expressed by an a\* value (redness) and visual color evaluation. Attached are a plot of the a\* and O<sub>2</sub> concentration for ground meat, top loin and pork chops; see figures 1, 2, and 3. The correlation coefficients for the three products were calculated to -0.71, -0.33, and -0.51. The relatively low coefficients are partly due to the large spread of the measured values and partly because the correlation between a\* and O<sub>2</sub> does not appear to be linear.

For ground meat we found a reduction of approximately 4-5 a\* values from 0 to 1% O<sub>2</sub>. A reduction of a\* to this degree is readily apparent. Samples stored in 1% or higher levels of O<sub>2</sub> had a score of between 3 and 4 on the color scale, i.e. slight to moderate discoloration. The results indicate that the tolerance limit for discoloration of ground meat in CO mixture is approximately 1% O<sub>2</sub>.

000124

Commission – Confidential/OSø/November 28, 1996/O-7224



MATFORSK – Norwegian Food Research Institute

For top loin, on the other hand, a smaller decrease in  $a^*$  and a weak correlation between  $a^*$  and  $O_2$  concentration were found. However, there seemed to be a reduction of about 2  $a^*$  values between 0 and 2%  $O_2$ , with a color scale score of about 3 on samples stored in over 2%  $O_2$ . This indicates a weak discoloration with a tolerance limit of approximately 2 %  $O_2$  for top loin.

The color of pork chops proved to be only slightly affected by the  $O_2$  concentration in the packaging gas, even when up to 2/3 of the gas was replaced with air. We found a reduction of 1 to 1.5  $a^*$  values between 0 and 5%  $O_2$  in the package gas, but this barely registered as discoloration with a score of 2-3 on the color scale.

The pH values at the end of the leak test were on average 5.59, 5.62 and 5.42 for ground meat, top loin and pork chops respectively.

Between the start and the end of the leak tests, we measured a reduction in the  $O_2$  concentrations of 80, 40 and 30 % for ground meat, top loin and pork chops respectively. This reduction can be due to meat respiration or consumption of  $O_2$  by bacteria. Ground meat has a high consumption of  $O_2$  due to a large surface area exposed to surrounding gas and frequently higher bacteria counts than whole meat.

The significance of residual  $O_2$  in package gas with regard to discoloration and microbiological storage life has been discussed previously in the report "Fresh Meat in Consumer Packaging - an Evaluation of Various Packaging Methods and Their Effect on Meat Quality." For storage in gas containing  $CO_2$  and/or  $N_2$  without the presence of  $CO$ , tolerance limits for discoloration have been found to be below 0.1 and 0.5%  $O_2$  for beef and pork respectively. Tests on pork has shown that the microbiological storage life was reduced when the package gas contained more than 2-4%  $O_2$ .

The ground meat containing 1% sodium that was tested in this survey, had obvious discoloration when the  $CO$  mixture contained at least 1%  $O_2$ . Sodium functions as a pro-oxidant, and will usually intensify or accelerate the discoloration of the meat. It is therefore likely that consumers will react on the color of ground meat when small leaks in the packaging exist. For top loin and pork chops, however, there is little likelihood that the minor discoloration occurring at above 2 and 5%  $O_2$  will serve as an indicator of leakage to the regular consumer. The lighting in store refrigerating counters will often conceal minor color nuances. All in all, these results show that  $CO$  has a strong bond to the myoglobin in whole, unsalted meat, which prevents the carbon myoglobin from being destabilized by  $O_2$  in the gas. Hence, discoloration is not a good indicator with regard to alerting consumers of leaks and risk of increased bacterial growth in meat such as top loin and pork chops.

Thanks

We are grateful to Frank Lundby for his valuable technical assistance.

000125

Commission – Confidential/OSø/November 28, 1996/O-7224





MATFORSK – Norwegian Food Research Institute

[Left-hand side of graph]  $a^*$

[Plot]

[Below graph]

% Oxygen

*Figure 1 The correlation between  $a^*$  (redness) and  $O_2$  concentration for ground meat packaged in a mixture of 60%  $CO_2$  / 40%  $N_2$  / 0.4% CO after two days storage at 4°C.  $n=20$ ,  $r=-0.71$ .*

[Left-hand side of graph]  $a^*$

[Plot]

[Below graph]

% Oxygen

*Figure 2 The correlation between  $a^*$  (redness) and  $O_2$  concentration for top loin packaged in a mixture of 60%  $CO_2$  / 40%  $N_2$  / 0.4% CO after five days storage at 4°C.  $n=18$ ,  $r=-0.33$ .*

[Left-hand side of graph]  $a^*$

[Plot]

[Below graph]

% Oxygen

*Figure 3 The correlation between  $a^*$  (redness) and  $O_2$  concentration for pork chops packaged in a mixture of 60%  $CO_2$  / 40%  $N_2$  / 0.4% CO after five days storage at 4°C.  $n=14$ ,  $r=-0.51$ .*

000126

Commission – Confidential/OSø/November 28, 1996/O-7224



reprints from

(5)

elsevier trends journals

Reprinted from  
*Trends in Food Science & Technology*

000127



Elsevier Trends Journals  
68 Hills Road  
Cambridge  
UK CB2 1LA  
Tel +44 1223 315961  
Fax +44 1223 321410

Retail meat can be packaged in gas mixtures containing 60–70% carbon dioxide (CO<sub>2</sub>), 30–40% nitrogen (N<sub>2</sub>) and <0.5% carbon monoxide (CO). This gas mixture with CO provides a unique combination of a long microbiological shelf life and a stable, cherry red colour of the meat. The shelf life of meat packaged in the CO mixture is longer than that of meat packaged in the commonly used atmospheres with high oxygen (O<sub>2</sub>), that is, approximately 70% O<sub>2</sub> and 30% CO<sub>2</sub>. The consumption of meat that has been packaged in a CO mixture will result in only negligible levels of carboxy-haemoglobin in the blood. It is highly improbable that the use of CO in the packaging of meat will present a toxic threat to consumers.

Modified-atmosphere packaging (MAP) is gaining increasing application in modern food distribution. Meat intended for retail sale can either be wrapped in vapour-tight, oxygen-permeable films or packaged in gas-tight films with a modified atmosphere (MA). The main purposes of the MAP of meat are twofold: to ensure the microbiological shelf life and the attractive red colour of the product. Consumers frequently interpret the colour of meat on retail display as an indicator of wholesomeness<sup>1</sup>.

CO is a colourless, odourless and tasteless gas. It is produced mainly through incomplete combustion of carbon-containing materials. Natural background levels of CO are 0.01–0.9 mg/m<sup>3</sup> (Ref. 2). In urban areas, 8-h mean concentrations of CO (i.e. mean CO concentrations are measured for each possible 8-h interval during a 24-h period, then averaged) are generally <20 mg/m<sup>3</sup>; however, maximum 8-h concentrations (i.e. the maximum mean concentration found during any one 8-h period) of up to 60 mg/m<sup>3</sup> have been recorded<sup>2</sup>. By far the most common cause of elevated CO concentrations in the blood is tobacco smoking<sup>3</sup>.

A challenge in the MAP of retail meat is the stabilization of the red colour of the product. The positive effect of CO on meat colour was known and patented over 100 years ago<sup>4</sup>. Despite this knowledge, CO has to date been applied commercially only to a limited extent in the MAP of meat. During the past 10 years, the Norwegian meat industry has been using a gas mixture of 60–70% CO<sub>2</sub>, 30–40% N<sub>2</sub> and 0.3–0.4% CO for the packaging of fresh retail meat, namely beef, pork and lamb. This gas mixture with CO maintains a stable, cherry red colour combined with a long microbiological shelf life

## Technological, hygienic and toxicological aspects of carbon monoxide used in modified-atmosphere packaging of meat

Oddvin Sørheim, Tore Aune and Truls Nesbakken

of the meat. The market share of retail meat packaged in this CO mixture in Norway is estimated at 50–60% (Dag Hallan, pers. commun.). In addition, some meat is also initially bulk-packaged in the CO mixture, and thereafter repackaged on trays with O<sub>2</sub>-permeable films in retail outlets. In other European countries not using such CO mixtures, market shares of retail meat packaged in atmospheres with a high O<sub>2</sub> concentration, with considerably shorter shelf lives, have been reported to be only 10–40%<sup>5,6</sup>.

In this article, we have evaluated the toxicological aspects of CO, and its mode of action and application in the MAP of meat.

### Technological and hygienic aspects of CO as a packaging gas for meat Gases for MAP

The most commonly used gases for the MAP of meat are CO<sub>2</sub>, N<sub>2</sub> and O<sub>2</sub>, although other gases, including CO, nitrous oxide, argon and ozone, have been tried to a limited extent<sup>1</sup>. CO<sub>2</sub> inhibits the growth of many microorganisms, but it has no effect *per se* on the colour of meat<sup>6</sup>. CO<sub>2</sub> is absorbed in meat and fat tissue at a ratio of ~1 litre of gas per kg of tissue<sup>7</sup>. N<sub>2</sub> affects neither the microbiology nor the colour of the meat, but prevents packages from collapsing, because it is not absorbed by the product. O<sub>2</sub> supports the growth of aerobic microorganisms; thus, removal of O<sub>2</sub> from the MA will extend the microbiological shelf life. High O<sub>2</sub> concentrations cause meat to have a temporary bright red colour; oxygen binds to the muscle pigment myoglobin, forming oxymyoglobin, which is gradually oxidized to grey-green-brown metmyoglobin<sup>8</sup> (Fig. 1). Gases for the packaging of meat are seldom used alone, but in mixtures, which vary according to the application. Examples of different gas mixtures for the MAP of meat are discussed below.

Oddvin Sørheim is at MATFORSK – Norwegian Food Research Institute, Osloveien 1, N-1430 Ås, Norway (fax: +47-64-970333; e-mail: oddvin.sorheim@matforsk.no). Tore Aune is at the Department of Pharmacology, Microbiology and Food Hygiene, Norwegian College of Veterinary Medicine, PO Box 8145 Dep., N-0033 Oslo, Norway. Truls Nesbakken is at the Norwegian Meat Cooperative, Department of Research and Development, PO Box 360 Økern, N-0513 Oslo, Norway; and at The Royal Veterinary and Agricultural University, Department of Veterinary Microbiology, Bülowsvej 13, DK-1870 Frederiksberg C, Denmark.




Fig. 1

Myoglobin forms and colour of meat.

### CO and colour

The main function of low levels of CO in MAs is to give meat a stable, cherry red colour, as a result of strong binding of CO to myoglobin and the formation of carboxymyoglobin<sup>9</sup> (Fig. 1). Although a substantial increase in the shelf life of meat can be obtained by using various MAs, it is often limited by discolouration due to the oxidation of myoglobin to metmyoglobin. This discolouration can be prevented by the inclusion of a low level of CO in the gas mixture.

Carboxymyoglobin is more resistant to oxidation than oxymyoglobin, owing to the stronger binding of CO to the iron-porphyrin site on the myoglobin molecule<sup>10</sup>. CO at concentrations of 1–5% increased the reduction of metmyoglobin, even in the presence of air<sup>11</sup>.

Examples of different gas mixtures that contain CO for the packaging of meat are given in Table 1. A mixture of 2% CO and 98% air was very effective in stabilizing the colour of beef for 15 d, compared with 5 d in air alone<sup>12</sup>. Ground beef patties stored in an atmosphere

of 1% CO/50% CO<sub>2</sub>/49% air retained a stable, red colour for at least 6 d, whereas the colour of samples stored in air was stable for only 3 d<sup>13</sup>.

The colour of beef was analysed during storage in MAs containing N<sub>2</sub> with 0.5–10% CO. Levels of CO >0.5% resulted in a stable, red colour for >30 d, whereas discolouration occurred after 5 d storage in control samples packaged in air<sup>14</sup>. In addition, samples of this beef were exposed to pure CO for 2–16 h before packaging in air. The colour stability of the CO-treated samples was no greater than that of untreated samples. However, in other experiments the exposure of beef to CO before vacuum packaging increased its redness and colour acceptability during subsequent chilled or frozen storage<sup>9,17</sup>. Beef loin roasts stored in 1% CO/51% CO<sub>2</sub>/30% O<sub>2</sub>/18% N<sub>2</sub> were shown to have lower levels of metmyoglobin on their surface than vacuum-packaged roasts. After a further 4 d on retail display, steaks from the roasts underwent less discolouration if they had previously been stored in the CO mixture<sup>15</sup>. Ground beef and beef loin steaks packaged in 1% CO/50% CO<sub>2</sub>/25% N<sub>2</sub>/24% O<sub>2</sub> or 1% CO/20% CO<sub>2</sub>/9% N<sub>2</sub>/70% O<sub>2</sub> retained a stable colour for 29 d<sup>16</sup>. Similarly, beef loin steaks packaged in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> maintained a stable, cherry red colour for up to 22 d<sup>18</sup>. Experiments with beef and higher levels of CO, that is, 2% CO/20% CO<sub>2</sub>/78% N<sub>2</sub>, resulted in meat that had a stable colour; however, its colour was characterized as 'too artificial' by a sensory panel<sup>6</sup>.

Based on the cited literature, the presence of 0.4–1.0% CO in MAs used for the packaging of meat seems sufficient to produce a stable, cherry red colour.

Cooked, cured meat products can also benefit from storage in MAs containing CO. Packaging in 1% CO/99% N<sub>2</sub> stabilized the colour of sliced bologna, indicating binding between CO and denatured myoglobin<sup>19</sup>.

Under certain circumstances, an undesirable pink or red colour can arise in cooked white meat, such as poultry, and cooked meat products without added nitrite<sup>20</sup>. Such colour problems can sometimes be linked with exposure to CO, which results in similar colours occurring after the use of MAs with CO. For example, roasted turkey was noted to be pink; this was probably due to the presence of CO and nitric oxide in the combustion gases in the oven. The pink colour did not occur when the turkeys were roasted in complete isolation from the oven gases<sup>21</sup>. Combustion engines produce various gases, including CO, which can affect live poultry during transportation to the abattoir. Meat from chickens that were exposed to exhaust fumes immediately before slaughter developed an undesired red colour on cooking<sup>22</sup>.

### CO and microbiology

Generally, the purpose of most of the experiments investigating the use of CO as a small component of MAs for meat has been to study its effect on colour stability, and more seldom its microbiological aspects. The growth of psychrotrophic bacteria on beef stored in MAs containing 0.5–10% CO in N<sub>2</sub> was lower, relative to controls, resulting in an increase in odour shelf life at

Table 1. Applications of carbon monoxide (CO) in the modified-atmosphere packaging of meat

Gas combinations (%)					Refs
CO	CO <sub>2</sub>	N <sub>2</sub>	O <sub>2</sub>	Air	
2				98	12
1	50			49	13
0.5–10		90–99.5			14
1	51	18	30		15
1	50	25	24		16
1	20	9	70		16
2	20	78			6
1–5				95–99	10
100*					9, 14, 17
0.4	60	40			18
0.3–0.4	60–70	30–40			b

\*Exposure before packaging

b>Data supplied by Norwegian meat plants

temperatures in the range of 0–10°C<sup>14</sup>. For example, beef packaged in a MA of 1% CO/99% N<sub>2</sub> had an odour shelf life of 24 d, compared with 18 d in 100% N<sub>2</sub>, and 7 d in air at 5°C. However, in another experiment with a MA of 20% CO<sub>2</sub>/70% O<sub>2</sub>/9% N<sub>2</sub>, the addition of 1% CO had no effect on the microbiological growth on ground beef and beef steaks<sup>16</sup>. The presence of bacteriostatic CO<sub>2</sub> in the latter experiment apparently reduced the importance of the effect of CO on the shelf life of the meat. The odour shelf life of steaks of beef loins stored in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> was 4 d longer than that of steaks stored in 70% O<sub>2</sub>/30% CO<sub>2</sub> at 4°C<sup>14</sup>. Beef steaks that were exposed to pure CO before vacuum packaging had an extended shelf life compared with untreated controls. The total aerobic plate, lactic acid bacteria and psychrotropic counts of CO-treated steaks were 1–2 log cycles lower than those of controls after 8 weeks storage at 4°C<sup>17</sup>. In a study using pure bacterial cultures, the presence of CO at a concentration of 5–30% in air had no effect on the growth of *Pseudomonas aeruginosa*, inhibited the growth rate of *Escherichia coli* (in proportion to the concentration of CO), increased the lag phase of *Achromobacter* and inhibited the growth rate of *Pseudomonas fluorescens*<sup>23</sup>.

### Toxicological aspects of CO

#### Health effects of CO

CO binds to the iron atom of haemoglobin in red blood cells, forming carboxyhaemoglobin (COHb). The affinity of haemoglobin for CO is ~240 times higher than its affinity for O<sub>2</sub>. CO also binds to myoglobin, cytochromes and some enzymes, but these reactions are considered to be of less importance than the formation of COHb<sup>2</sup>. The binding of CO to haemoglobin is reversible, with a half-life of ~4.5 h in individuals who are at rest.

Although CO acts primarily by interfering with O<sub>2</sub> transport, it also reduces the delivery of O<sub>2</sub> to the various tissues<sup>3</sup>. In humans, health effects are mainly manifested in the cardiovascular system, the nervous system and in the foetus.

The COHb concentration in blood, often referred to as the COHb percent (COHb%), is a function of the CO concentration in the air, the exposure time, and the level of physical activity of the individual<sup>24</sup> (see Table 2). At a COHb concentration of ~2.5%, the most sensitive individuals (patients suffering from cardiovascular diseases) display changes in cardiac function and report chest pain. At somewhat higher COHb concentrations, they experience reduced working capacity and the onset of angina pectoris on exercise<sup>25,26</sup>. In healthy adults, no adverse health effects were described at CO concentrations that result in <5% COHb<sup>27</sup>.

A small amount of CO is formed naturally in the human body, owing to the breakdown of haemoproteins.

Table 2. Estimate of carboxyhaemoglobin percent (COHb%) in human blood at different concentrations of carbon monoxide (CO) in the atmosphere, depending on the level of physical activity\*

Exposure		COHb%		
CO concentration (mg/m <sup>3</sup> )	Time (h)	At rest	Moderate activity	Heavy work
10	8	1.3	1.4	1.4
25	1	1.0	1.5	2.0
40	1	1.3	2.2	2.9

\*Data taken from Ref. 24

Table 3. Association between different carboxyhaemoglobin (COHb) levels in blood and health effects\*

COHb%	Observed health effects
≥50	Unconsciousness, lethal if not treated
≥30	Headache, nausea, vomiting, dizziness
≥10	Life threatening for heart and lung patients; headache in other individuals
25	Reduced maximum oxygen consumption during exercise in healthy individuals
25	Reduced visual perception, learning ability and fine motor performance
25	The foetus can be affected on carbon monoxide exposure of pregnant women
22.9	Angina patients endure less physical strain before experiencing attack
22.3	Reduced physical working capacity, especially endurance
22	Possible reduction in attention and ability to concentrate
22	Signs of local lack of oxygen and onset of chest pain in heart patients

\*Data taken from Refs 25–27

Such production leads to a COHb concentration of ~0.5%. The average COHb concentration in non-smokers is 1.2–1.5%, and ~3–4% in smokers<sup>27</sup>.

The absorption and excretion of CO from the body occur relatively slowly; thus, exposure to elevated CO levels over short time periods will not result in a significant increase in the COHb level in the blood. Table 3 details the various health effects observed at different COHb levels. This table confirms that exposure to CO that results in a COHb level greater than ~2% should be avoided to protect the most vulnerable individuals in the population.

In order to protect the most vulnerable in society, a Norwegian expert group on air pollution<sup>27</sup> recommended maximum CO concentrations for different exposure times that will prevent COHb levels from exceeding 1.5%, taking into consideration endogenous CO formation (Table 4).

Table 4. Estimates of carbon monoxide (CO) levels in ambient air that will result in carboxyhaemoglobin (COHb) levels of 1.5%, including endogenous CO production\*

Exposure time	CO concentration in air (mg/m <sup>3</sup> )		
	At rest	Moderate activity	Heavy work
15 min	170	80	52
30 min	86	42	29
1 h	48	24	18
8 h	11.5	9.2	9.2

\*Data taken from Ref. 27

#### Exposure to CO on consumption of fresh meat treated with a CO gas mixture

Very little information exists in the literature on the exposure to CO following the consumption of meat that has been treated with CO gas. The inhalation of air containing CO at a level of 57 mg/m<sup>3</sup> (the acceptable level in working environments in the USA) would provide a COHb level for a prolonged time period (hours) of at least 14 times that of the level reached temporarily on the consumption of 225 g of meat that had been packaged in CO at the saturation level for myoglobin<sup>14</sup>. In this estimate, it was assumed that the saturation of the meat myoglobin and haemoglobin was maximal and that the transfer of CO from the gastrointestinal tract to the blood was 100%. Consequently, even for such a 'worst-case' scenario, the treatment of meat with CO gas appears to contribute very little to COHb levels, relative to levels that are considered safe in the working environment. The exposure of beef to an atmosphere containing 1% CO for 3 d resulted in ~30% saturation of the meat myoglobin<sup>21</sup>. CO is lost from previously CO-treated meat during storage in the absence of CO, with a half-life of ~3 d. When the beef was cooked at 195°C, only 0.1 mg of CO remained per kg of meat. The loss of CO amounted to ~85%.

#### Comparison of CO exposure from air and the consumption of gas-treated meat

Data are very scarce, but comparisons still allow crude estimates to be made. An adult inhales ~10–20 m<sup>3</sup> of air in 24 h (depending on their level of activity). This is equivalent to 0.42–0.84 m<sup>3</sup>/h (or 3.36–6.72 m<sup>3</sup> in 8 h). In order to prevent a maximum COHb level in the blood of 1.5% being exceeded, the CO concentration in air for a 1-h period of moderate physical activity should not exceed 24 mg/m<sup>3</sup>, or 9.2 mg/m<sup>3</sup> in 8 h (according to Table 4). In contrast, the consumption of meat that had been treated for 3 d in an atmosphere containing 1% CO yielded ~0.1 mg of CO per kg of meat on storage and cooking<sup>21</sup>. Based on these data, a comparison can be made from the two methods of exposure to CO, and is shown in Table 5.

Equilibrium between CO present in the atmosphere and the COHb concentration in blood is reached only

after a considerable period of time (depending on the concentration and level of physical activity). Even in a 'worst-case' scenario, equilibration between the CO concentration in the gastrointestinal tract and blood will take time. Furthermore, the absorption of CO from the gastrointestinal tract into the blood will in all probability be less effective than the absorption of CO from the lungs, which are composed of tissues that are designed to facilitate gas exchange between the alveoli and the blood. Consequently, it is highly probable that the consumption of one meal of CO-exposed meat per day will not result in measurable increases in the COHb level in blood.

#### Toxicological evaluation of the use of CO as a packaging gas for meat

Unfortunately, the European Union (EU) has not evaluated CO for use as a packaging gas for meat. However, CO<sub>2</sub> and nitrous oxide (N<sub>2</sub>O) have both been approved for use for extraction purposes, and it was considered unnecessary to adopt an acceptable daily intake (ADI) value for these gases in this application<sup>29</sup>.

In order to avoid possible adverse health effects in those individuals who are the most susceptible, a Norwegian expert group on air pollution recommended maximum CO concentrations in ambient air that result in COHb levels not exceeding 1.5% (including endogenous CO production)<sup>27</sup>. Estimates detailed above indicate that, even assuming an improbable 100% absorption of CO from the gastrointestinal tract into the blood, the consumption of meat that has been treated with 1% CO will result in COHb levels that are negligible (approximately three orders of magnitude lower) compared with those resulting from exposure in the working environment to CO at an acceptable level. Consequently, it is highly improbable that CO exposure from meat packaged in an atmosphere containing up to 0.5% will represent a toxic threat to consumers through the formation of COHb.

#### Alternatives to the MAP of retail meat

Currently, the most commonly used MA for the retail packaging of meat contains O<sub>2</sub> at a high concentration in combination with CO<sub>2</sub>, such as ~70% O<sub>2</sub>/30% CO<sub>2</sub>. The shelf life of meat in a high O<sub>2</sub> atmosphere in commercial practice, typically at temperatures of 6–8°C, is ~7 d, being limited both by microbiological spoilage and discolouration. Meat that is stored in a high O<sub>2</sub> concentration is often spoiled by bacteria such as *Brochothrix thermosphacta* and pseudomonads<sup>30</sup>. In MAs with a high concentration of O<sub>2</sub>, the meat normally maintains its bright red oxymyoglobin colour for 4–7 d before the colour starts deteriorating to grey-brown, owing to the formation of metmyoglobin<sup>31</sup>. This length of time is often not considered to be sufficient to display and sell the product.

The use of MAs with a high concentration of CO<sub>2</sub>, either alone or in combination with up to 70% N<sub>2</sub>, would increase the microbiological shelf life of the meat compared with that of meat in a MA with a high O<sub>2</sub> concentration. The absence of O<sub>2</sub> together with the presence

of CO<sub>2</sub> retards microbiological growth. Unfortunately, the colour of meat packaged in MAs containing CO<sub>2</sub> is less satisfactory, being either purple or grey-brown due to the formation of deoxymyoglobin or metmyoglobin, respectively. The meat inevitably discolours when the O<sub>2</sub> concentration is

low. Metmyoglobin formation can be avoided by maintaining O<sub>2</sub> concentrations <0.01–0.1% for beef<sup>31</sup> and <0.5% for pork<sup>32</sup>. These low O<sub>2</sub> levels, particularly for beef, are difficult to achieve in most commercial packaging operations, because a small amount of air will unavoidably be incorporated in the MAs of the packages. MAs with a high CO<sub>2</sub> concentration seem to be useful for retail packaging if a low concentration of CO is also included to stabilize myoglobin and the meat colour.

Vacuum packaging is commonly used for the bulk storage, transportation and export of meat. However, vacuum packaging has not proved to be a successful method for the retail packaging of meat, because of the purple deoxymyoglobin colour of the meat and the visible exudate that occurs in the packages<sup>33</sup>. Meat that is packaged in a vacuum cannot be presented in the bright red oxymyoglobin state, which depends on the presence of a high concentration of O<sub>2</sub><sup>30,33</sup>, or in the cherry red carboxymyoglobin state, which requires CO to be included in the MA.

One of the objections that has been raised against the use of CO as a packaging gas is the potential hazard it might represent to workers in meat plants. Although the use of pure CO for mixing in the plant would certainly pose such a risk, the delivery of 1% CO in a mixture with 99% N<sub>2</sub>, which has been the practice of gas suppliers to the Norwegian meat industry, is recognized by the health authorities to be a very safe handling procedure.

MAs that contain 60–70% O<sub>2</sub> must be handled carefully, because they are explosive gas mixtures. Strict safety regulations apply to explosive gas mixtures, increasing the costs of equipment and packaging operations. The benefits of a CO mixture is that it carries no risk of explosion and therefore does not increase handling costs.

Despite the long-term knowledge of the many advantages of the use of CO as a component of MAs for meat, CO mixtures have not been adopted to any great extent by the global meat industry. In many countries, including the USA and countries within the EU, CO is presently not permitted for use in the MAP of meat<sup>34</sup>. However, Norwegian food control authorities have not opposed the use of CO as a packaging gas at concentrations of up to 0.5%. As a member of the European Economic Agreement, Norway is expected to adapt gradually to EU food regulations, including those relating to gases for the packaging of foods. The Norwegian meat industry is therefore preparing an inquiry, to be directed at the Norwegian and EU food control authorities, for the continued use of CO in the MAP of red meats, which will be partly based on the toxicological evaluation described in this article.

Table 5. Theoretical uptake of carbon monoxide (CO) in blood

Exposure method	CO intake in 1 h	CO intake in 8 h
Lungs (15 m <sup>3</sup> /d)	24 mg × 0.625 = 15.1 mg	9.2 mg × 5 = 46.0 mg
Meat (250 g, CO treated)	0.025 mg	0.025 mg

Consumers may evaluate the shelf life of packaged meat on the basis of its colour. A possible negative aspect of using CO in the MAP of retail meat is concern that consumers might misjudge the quality of a product, because its true microbiological status may be masked by its stable, cherry red carboxymyoglobin colour<sup>1</sup>. However, consumers will be able to detect spoilage by the presence of off-odours. At the current low concentrations, <0.5%, CO *per se* seems to have no or only minor effects on bacteria and the shelf life of the meat. The combination of CO with a high concentration of CO<sub>2</sub>, for example 60–70%, is necessary for microbiological control. Although MAP enables centralized packaging operations with quality control to be carried out, MAP alone cannot guarantee the shelf life of the product. Sufficient shelf life can be obtained only through the proper quality control of raw materials, production, packaging, chill chain and retail conditions.

## Conclusions

Gas mixtures that contain a low concentration of CO, up to 0.5%, and a high concentration of CO<sub>2</sub>, ~70%, have many advantages with respect to shelf life, colour stability, labour safety and costs. The use of CO at such concentrations does not present any toxic threat to consumers. Considering the benefits the Norwegian meat industry has experienced with CO gas mixtures over the past decade, potential exists for their wider application in the retail packaging of meat.

## Acknowledgement

The financial support from the Research Council of Norway is greatly appreciated.

## References

- Kropl, D. (1980) 'Effects of Retail Display Conditions on Meat Color' in *Proc. Reciprocal Meat Conf.* 33, 15–32
- WHO (1979) *Environmental Health Criteria*. 13. Carbon Monoxide, World Health Organization, Geneva, Switzerland
- WHO (1987) *Air Quality Guidelines for Europe* (World Health Organization Regional Publications, European Series, No. 23), World Health Organization, Geneva, Switzerland
- Church, N. (1994) 'Developments in Modified-atmosphere Packaging and Related Technologies' in *Trends Food Sci. Technol.* 5, 345–352
- Bruce, J. (1990) 'A Comparison of American and European Markets for the MAP of Fresh Red Meats' in *Proceedings of the International Conference of Modified Atmosphere Packaging*, Stratford-upon-Avon (Day, B.P.F., ed.), Part 2, pp. 1–14, Campden Food & Drink Research Association, Chipping Campden, UK
- Renner, M. and Labadie, J. (1993) 'Fresh Red Meat Packaging and Meat Quality' in *Proceedings of the 39th International Congress of Meat Science and Technology*, Calgary, pp. 361–387, Lacombe Research Centre, Lacombe, Alberta, Canada
- Gill, C.O. (1988) 'The Solubility of Carbon Dioxide in Meat' in *Meat Sci.* 22, 65–71
- Cornforth, D. (1994) 'Color – Its Basis and Importance' in *Quality Attributes and Their Measurement in Meat, Poultry and Fish Products. Advances in Meat Research – Vol. 9* (Pearson, A.M. and Dutson, T.R., eds), pp. 34–78, Blackie

- 9 Lentz, C.P. (1979) 'Effect of Light Intensity and Other Factors on the Color of Frozen Prepackaged Beef' in *Can. Inst. Food Technol. J.* 12(2), 47-50
- 10 Wolfe, S.K. (1980) 'Use of CO<sub>2</sub>- and CO<sub>2</sub>-enriched Atmospheres for Meats, Fish, and Produce' in *Food Technol.* 34(3), 55-63
- 11 Lanier, T.C., Carpenter, J.A., Toledo, R.T. and Reagan, J.O. (1978) 'Metmyoglobin Reduction in Beef as Affected by Aerobic, Anaerobic and Carbon Monoxide Containing Environments' in *J. Food Sci.* 43, 1788-1792, 1796
- 12 El-Badawi, A.A., Cain, R.F., Samuels, C.E. and Angelmeier, A.F. (1964) 'Color and Pigment Stability of Packaged Refrigerated Beef' in *Food Technol.* 18(5), 159-163
- 13 Gee, D.L. and Brown, W.D. (1978) 'Extension of Shelf Life in Refrigerated Ground Beef Stored Under an Atmosphere Containing Carbon Dioxide and Carbon Monoxide' in *J. Agric. Food Chem.* 26, 274-276
- 14 Clark, D.S., Lentz, C.P. and Roth, L.A. (1976) 'Use of Carbon Monoxide for Extending Shelf-life of Prepackaged Fresh Beef' in *Can. Inst. Food Sci. Technol. J.* 9(3), 114-117
- 15 Seidemann, S.C., Carpenter, Z.L., Smith, G.C., Dill, C.W. and Vanderzant, C. (1979) 'Physical and Sensory Characteristics of Beef Packaged in Modified Gas Atmospheres' in *J. Food Protect.* 42, 233-239
- 16 Luño, M., Beltrán, J.A. and Roncales, P. (1996) 'Shelf-life Extension and Colour Stabilization of Beef Packaged in a Low O<sub>2</sub> Atmosphere Containing CO: 1. - Loin Steaks. 2. - Ground Meat' in *Proceedings of the 42nd International Congress of Meat Science and Technology, Lillehammer*, pp. 76-79, MATFORSK - Norwegian Food Research Institute, Ås, Norway
- 17 Brewer, M.S., Wu, S., Field, R.A. and Pay, B. (1994) 'Carbon Monoxide Effects on Color and Microbial Counts of Vacuum-packaged Fresh Beef Steaks in Refrigerated Storage' in *J. Food Qual.* 17, 231-244
- 18 Sorheim, O., Nissen, H. and Nesbakken, T. (1997) 'Shelf Life and Colour Stability of Beef Loin Steaks Packaged in a Modified Atmosphere with Carbon Monoxide' in *Proceedings of the 43rd International Congress of Meat Science and Technology, Auckland*, 694-695, Meat Industry Research Institute of New Zealand, Hamilton, New Zealand
- 19 Aasgaard, J. (1993) 'Colour Stability of Packed Meat Products' in *Fleischwirtschaft* 73, 428-431
- 20 Ledward, D.A. and Shaw, R. (1994) 'Meat Processing - Undesirable Pink Coloration in Uncooked Cured Meat' in *Meat Focus Int.* 3, 295-298
- 21 Maga, J.A. (1994) 'Pink Discoloration in Cooked White Meat' in *Food Rev. Int.* 10, 272-286
- 22 Froning, G.W., Aather, F.B., Daddario, J. and Hartung, T.E. (1969) 'Effect of Automobile Exhaust Fume Inhalation by Poultry Immediately Prior to Slaughter on Color of Meat' in *Poultry Sci. Wis.* 48, 485-487
- 23 Gee, D.L. and Brown, W.D. (1980) 'The Effect of Carbon Monoxide on Bacterial Growth' in *Meat Sci.* 5, 215-222
- 24 Coburn, R.F., Forster, R.E. and Kane, P.B. (1965) 'Considerations of the Physiological Variables That Determine the Blood Carboxyhemoglobin in Man' in *J. Clin. Invest.* 44, 1899-1910
- 25 Aronow, W.S. and Isbell, M.W. (1973) 'Carbon Monoxide Effect on Exercise-induced Angina Pectoris' in *Ann. Intern. Med.* 79, 392-395
- 26 Allred, E.N. et al. (1989) 'Short-term Effects of Carbon Monoxide Exposure on the Exercise Performance of Subjects with Coronary Artery Disease' in *New Engl. J. Med.* 321, 1426-1432
- 27 Aunan, K. et al. (1992) *Effects of Ambient Air Pollution on Health and Environment - Air Quality Guidelines*, Report No. 92:16, pp. 154-170, State Pollution Control Authority (SFT), Oslo, Norway (in Norwegian)
- 28 Watts, D.A., Wolfe, S.K. and Brown, W.D. (1978) 'Fate of [<sup>14</sup>C] Carbon Monoxide in Cooked or Stored Ground Beef Samples' in *J. Agric. Food Chem.* 26, 210-214
- 29 Commission of the European Union (CEU) (1981) *Food - Science and Technique (Report from the Scientific Committee for Food, 11th Series)*, p. 6, CEU, Brussels, Belgium
- 30 Gill, C.O. (1996) 'Extending the Storage Life of Raw Chilled Meats' in *Meat Sci.* 43 (suppl. issue), 99-110
- 31 Gill, C.O. and McGinnis, J.C. (1995) 'The Effects of Residual Oxygen Concentration and Temperature on the Degradation of the Colour of Beef Packaged Under Oxygen-depleted Atmospheres' in *Meat Sci.* 39, 387-394
- 32 Sorheim, O., Erlandsen, T., Nissen, H., Lea, P. and Høyem, T. 'Effects of Modified Atmosphere Storage on Colour and Microbiological Shelf Life of Normal and Pale, Soft and Exudative Pork' in *Meat Sci.* (in press)
- 33 Taylor, A.A., Down, N.F. and Shaw, B.G. (1990) 'A Comparison of Modified Atmosphere and Vacuum Skin Packing for the Storage of Red Meats' in *Int. J. Food Sci. Technol.* 25, 98-109
- 34 European Parliament and Council Directive (1995) of 20 February 1995 on 'Food Additives Other Than Colours and Sweeteners' in *Off. J. Eur. Commun.* No. L61, 1-38, 18.3.95



## The storage life of beef and pork packaged in an atmosphere with low carbon monoxide and high carbon dioxide

Oddvin Sørheim<sup>a,\*</sup>, Hilde Nissen<sup>a</sup>, Truls Nesbakken<sup>b</sup>

<sup>a</sup>MATFORSK-Norwegian Food Research Institute, Osloveien 1, N-1430 Ås, Norway

<sup>b</sup>Norwegian Meat Cooperative, Department of Research and Development, PO Box 360 Økern, N-0513 Oslo, Norway

Received 13 February 1998; received in revised form 28 October 1998; accepted 1 December 1998

### Abstract

Ground beef, beef loin steaks and pork chops were packaged in modified atmospheres of 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> and 70% O<sub>2</sub>/30% CO<sub>2</sub>. In addition ground beef was packaged in clipped chub packs, beef loin steaks were vacuum packaged, and pork chops were packaged in an atmosphere of 60% CO<sub>2</sub>/40% N<sub>2</sub> with each pack containing an O<sub>2</sub> absorber. The packs were stored in the dark at 4 or 8°C for up to 21 days. Meat in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> had a stable bright red colour that lasted beyond the time of spoilage. The storage lives in this gas mixture at 4°C, as limited by off-odours, were 11, 14 and 21 days for ground beef, beef loin steaks and pork chops, respectively. The 70% O<sub>2</sub>/30% CO<sub>2</sub> atmosphere resulted in an initially bright red to red colour of the meat, but the colour was unstable and off-odours developed rapidly. The off-odours probably were caused by *Brochothrix thermosphacta*, which grew in all meat types, or by pseudomonads in ground beef. Meat stored in chub packs, vacuum packs or 60% CO<sub>2</sub>/40% N<sub>2</sub> with an O<sub>2</sub> absorber developed off-odours and microflora similar to those of meat in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub>, but with less acceptable appearances. These results show that a low CO/high CO<sub>2</sub> atmosphere is effective for preserving retail-ready meat. © 1999 Elsevier Science Ltd. All rights reserved.

### 1. Introduction

The main reasons for modified atmosphere packaging (MAP) of red meats for retail sale are to prolong the microbiological shelf life and to maintain an attractive red colour of the product. Modified atmospheres (MA) usually consist of carbon dioxide (CO<sub>2</sub>) for inhibiting microbiological growth, oxygen (O<sub>2</sub>) for enhancing colour and, occasionally, nitrogen (N<sub>2</sub>) as a filler. The most common gas mixture for retail-ready meat contains approximately 70% O<sub>2</sub> and 30% CO<sub>2</sub>, and gives the product an extended shelf life compared to air (Gill, 1996). The shelf life and colour stability of meat stored in this gas mixture is still limited. To obtain a stable red colour for the meat, low concentrations (<1%) of carbon monoxide (CO) can be introduced in the MA. Then, O<sub>2</sub> can be removed from the gas mixture and the concentration of bacteriostatic CO<sub>2</sub> can be increased. Anaerobic conditions extend the shelf life of meat considerably compared to air and O<sub>2</sub>-enriched atmospheres (Gill & Molin, 1991). CO binds strongly to the meat

pigment myoglobin to form stable carboxymyoglobin which has a cherry red colour (El-Badawi, Cain, Samuels, & Angelmeier, 1964). Low concentrations of CO have little effect on the microflora of meat (Clark, Lentz, & Roth, 1976; Gee & Brown, 1978; Luño, Beltrán, & Roncalés, 1998).

The Norwegian meat industry has for the past decade been using a gas mixture of approximately 0.3–0.5% CO, 60–70% CO<sub>2</sub> and 30–40% N<sub>2</sub> in retail-ready packages of beef, pork and lamb. Packages with this gas mixture now have a 50–60% share of the domestic, retail, red meat market. The technological, hygienic and toxicological aspects of using CO in MA for meat have recently been reviewed with the conclusion that CO used in concentrations up to 1% does not present a toxic hazard to the consumer (Sørheim, Aune, & Nesbakken, 1997a). However, CO may mask spoilage, because the stable cherry red colour can last beyond the microbiological shelf life of the meat (Kropf, 1980).

The inclusion of CO in MA for meat is controversial. CO is presently not allowed in MA for meat in the USA and in the EU (Cornforth, 1994; European Parliament and Council Directive, 1995). However, Norwegian food control authorities have up to now not opposed

\* Corresponding author. Tel.: +47-64-970100; Fax: +47-64-970333; E-mail: oddvin.sorheim@matforsk.no

the use of up to 0.5% CO in MA for meat. This would change with an adoption of EU food regulations in Norway. Consequently, the Norwegian meat industry is seeking amendments of current EU food regulations relating to the use of CO in MAP of red meats. If the use of CO should be disallowed, other means of maintaining the long shelf life and the attractive red colour of the meat will have to be sought.

The aim of the present experiments was to compare a commercial Norwegian CO/CO<sub>2</sub>/N<sub>2</sub> mixture with alternative gas mixtures and packaging methods for their effects on the off-odour, microflora and colour of ground beef, beef loin steaks and pork chops stored at 4 or 8°C for up to 21 days.

## 2. Materials and methods

### 2.1. Preparation of meat

#### 2.1.1. Ground beef

Twenty cow and bull carcasses of Norwegian Red Cattle, which weighed on average 275 kg, were electrically stimulated with 90 V and were chilled using programmed air temperatures between 12 and -5°C. Two days after slaughter the carcasses were deboned, and trimmings with 14% fat were ground through a 4 mm plate. The batch of ground beef was divided into 500 g portions.

#### 2.1.2. Beef loin steaks

Loins (*m. longissimus lumborum et thoracis*) with ultimate pH values below 5.8 were deboned from 25 bull carcasses of Norwegian Red Cattle. These carcasses, which weighed on average 275 kg, were stimulated, chilled and deboned the same way as the carcasses used in the preparation of ground beef. The loins were vacuum packaged and aged for 11 days at 3°C. Thereafter, the loins were cut into steaks 2.5 cm thick, and were randomly assigned to retail packs which each contained two steaks.

#### 2.1.3. Pork chops

Thirty pig carcasses of Norwegian Land Race, which weighed on average 75 kg, were blast-chilled. Four days after slaughter, bone-in loins were removed and crust-frozen in liquid N<sub>2</sub> at -50°C for 20 min to facilitate cutting of chops. The chops, which were 1.6 cm thick, were randomly assigned to retail packs which each contained two chops.

### 2.2. Packaging

Ground beef, beef loin steaks and pork chops were packaged in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> (CO mixture) and 70% O<sub>2</sub>/30% CO<sub>2</sub> (high O<sub>2</sub>). In addition, ground beef was packaged in clipped chub packs, beef loin steaks were vacuum packaged and pork chops were packaged in 60% CO<sub>2</sub>/40% N<sub>2</sub> with one Ageless® FX-

100 O<sub>2</sub> absorber (Mitsubishi Gas Chem. Co. Inc., Tokyo, Japan) in each pack (mixture with O<sub>2</sub> absorber).

The meat was packaged at a commercial meat plant within 2 h of grinding or cutting. Meat in the CO mixture, the high O<sub>2</sub> mixture and the mixture with O<sub>2</sub> absorber was packaged in an Ilapak Delta 2000 flow-packaging machine (Ilapak Machine Auto S.A., Grancia, Switzerland). The CO mixture was a blend of 1% CO/99% N<sub>2</sub> with 100% CO<sub>2</sub>. The high O<sub>2</sub> mixture was used as a preblend. The mixture with O<sub>2</sub> absorber was a blend of 100% N<sub>2</sub> with 100% CO<sub>2</sub> (all gases, Hydrogas, Porsgrunn, Norway). The initial gas volume to meat weight ratio in the packs was approximately 1.5 to 1. The packs consisted of polyethylene trays (Færch Plast, Holstebro, Denmark) wrapped in Cryovac BDF 550 shrinking film (Cryovac, Milan, Italy) with an O<sub>2</sub> transmission rate of 19 cm<sup>3</sup>/m<sup>2</sup>/24 h/atm at 23°C and 0% RH. Chub packs of ground beef were packaged in a clipping machine (Poly-Clip, Frankfurt, Germany) using a red, fishingnet-patterned, polyethylene film (SFK, Vidovre, Denmark) with an O<sub>2</sub> transmission rate of 500 cm<sup>3</sup>/m<sup>2</sup>/24 h/atm at 23°C and 0% RH. Beef loin steaks were vacuum packaged in a Multivac 5100 thermo-forming machine (Multivac, Wolfertschwenden, Germany) using a terephthalate/polyethylene upper film and polyamide/polyethylene lower film with O<sub>2</sub> transmission rates of 10 and 16 cm<sup>3</sup>/m<sup>2</sup>/24 h/atm at 23°C and 0% RH, respectively (Danisco, Horsens, Denmark).

### 2.3. Storage and sampling of meat

Five samples were collected from the ground beef batch, beef loins and pork loins before packaging, for pH measurements and microbiological analyses.

The packaged meat was stored in dark chilling rooms at 4 ± 0.5 or 8 ± 0.5°C for up to 21 days at least until off-odours developed. Five packs were removed per product, packaging method, storage temperature and sampling day after the following storage times:

- ground beef: 2, 4, 6, 8 or 11 days;
- beef loin steaks: 3, 7, 10 or 14 days; and
- pork chops: 3, 7, 10, 14, 17 or 21 days.

### 2.4. Gas analyses

The atmospheres of packs with MA were analysed for O<sub>2</sub> and CO<sub>2</sub> immediately after packaging (approximately every tenth pack) and at sampling (all packs). O<sub>2</sub> was determined using a Toray LC 700-F gas analyser (Toray Engineering, Osaka, Japan) and CO<sub>2</sub> using a Toray PG-100 gas analyser (Toray). The threshold levels for the O<sub>2</sub> and CO<sub>2</sub> analyses were 0.05 and 1%, respectively. Gas samples of 10 cm<sup>3</sup> were removed with a syringe through selfsealing patches on the packs.

## 2.5. pH

The pH measurements were made directly in the meat with an Ingold Xerolyt gel electrode (Mettler-Toledo A.G., Greifensee, Switzerland).

## 2.6. Odour

The meat was evaluated for odours by a three member trained panel between 0.5 and 1 min after opening of the packs. The off-odour scale used was: 1 = none, 3 = slight and 5 = extreme. Scores of 3 or below were considered acceptable.

## 2.7. Microbiology

Ten gram meat samples were collected from portions of the ground beef, and diluted in 90 g peptone water. A sample 25 cm<sup>2</sup> and 2-3 mm thick was removed from the surface of each beef loin or steak and pork loin or chop with a scalpel, and diluted in 100 ml peptone water. Each sample was macerated in a Stomacher for 1 min. Serial 10-fold dilutions of each Stomacher fluid were prepared, and 20 µl volumes of appropriate dilutions were plated in duplicate on the following media:

- plate count agar (PCA; Difco, Difco Laboratories, Detroit, MI, USA) for total viable counts;
- de Man, Sharpe and Rogosa agar (MRS; Oxoid, Unipath Ltd., Basingstoke, Hampshire, UK) adjusted to pH 5.7 for lactic acid bacteria (de Man, Rogosa, & Sharpe, 1960);
- streptomycin thallous acetate actidione agar base (STAA; CM 881 with selective supplement SR 151; Oxoid) for *Brochothrix thermosphacta*;
- pseudomonads agar base (CFC; CM 559 with selective supplement SR 103; Oxoid) for pseudomonads;

In addition, 1 ml portions of appropriate dilutions were plated in duplicate on petrifilm coliform count plates (3M Microbiology Products, St. Paul, MN, USA) for enumeration of coliforms and *Escherichia coli*.

Plates of PCA, MRS, STAA and CFC were incubated at 20°C for four days, and petrifilm plates at 30°C for up to 2 days, all aerobically. Counts were expressed as colony forming units (CFU) per g or cm<sup>2</sup>.

## 2.8. Colour

A six-member trained panel evaluated the colour of the meat in intact packs under 1200 ± 200 lux Warmton Lumilux L36W/31 yellow-white light (Osram, Drammen, Norway). The colour was assessed on a scale where 1 = bright red (ground beef and beef loin steaks) or light bright red (pork chops), 2 = red (ground beef

and beef loin steaks) or light red (pork chops), 3 = slightly brown, grey or green, 4 = moderately brown, grey or green and 5 = extremely brown, grey or green (National Live Stock and Meat Board, 1991).

A Minolta Chroma Meter CR-300 (Minolta Camera Co., Osaka, Japan) with 8 mm viewing port and illuminant D<sub>65</sub> was used for measuring CIE *a*\* values (redness). The colour was measured directly at the meat surface within 1 min of opening of each pack.

Ground beef in chub packs was not included in the colour analyses because the red packaging film hides the colour of the product. With pork chops, the colour of only the *m. longissimus lumborum et thoracis* was analysed.

## 2.9. Statistics

Analysis of variance by Tukey's multiple comparisons test was performed using the Systat programme, version 6 (Systat Inc., Evanston, IL, USA).

## 3. Results

### 3.1. Gas composition

The initial O<sub>2</sub> concentrations in packs with the CO mixture and the mixture with O<sub>2</sub> absorber were all below 0.5% immediately after packaging. O<sub>2</sub> was not detected in these packs after 2 or 3 days storage. The level of O<sub>2</sub> in packs of high O<sub>2</sub> was reduced from the initial 70 to 60-65% during storage for up to 21 days. Concentrations of CO<sub>2</sub> in the packs were generally reduced by one fifth after 2 or 3 days storage, and were then stable (data not shown).

### 3.2. Storage life of ground beef

The time to develop off-odours was 2 to 3 days longer for ground beef stored in the CO mixture and in chub packs than in high O<sub>2</sub>, and it was 4 or 5 days longer at 4 than at 8°C for all three packaging methods (Table 1). In high O<sub>2</sub>, the total viable counts increased faster and were higher (*p* < 0.01) than for the other two types of packaging after 2 days at either 4 or 8°C [Fig. 1(a)]. The total viable counts were more than 90% lactic acid bacteria (data not shown). The high numbers of lactic acid bacteria in ground beef, up to approximately log<sub>10</sub> 8 CFU/g, caused a decrease in the pH value from the initial 5.7 to 5.2 after 6 days when the meat was stored in the CO mixture or chub packs at 8°C (data not shown). At 4°C, the pH value was reduced to 5.5 after 11 days in both those packaging systems. The numbers of *B. thermosphacta* increased, in meat in high O<sub>2</sub> [Fig. 1(b)]. In meat in high O<sub>2</sub> the numbers of pseudomonads increased up to approximately log<sub>10</sub> 7 CFU/g, but only to log<sub>10</sub> 5 and 6 CFU/g in

meat in the CO mixture or chub packs, respectively (data not shown).

Ground beef in the CO mixture had a stable bright red colour, as shown by both the low colour scores and the high  $a^*$  values [Fig. 1(c) and (d)]. Meat in high  $O_2$  was significantly less red ( $p < 0.05$ ) than meat in the CO mixture, with higher colour scores and lower  $a^*$  values at day 2 and at later storage times at both 4 and 8°C. The colour of meat in high  $O_2$  deteriorated with time, significantly faster ( $p < 0.01$ ) at 8 than at 4°C.

Table 1

Time for development of off-odours in different types of meat in various packagings at storage temperatures of 4 or 8°C

Product	Packaging <sup>a</sup>	Time of off-odour detection (days)	
		4°C	8°C
Ground beef	CO mixture	11	6
	High $O_2$	8	4
	Chub packs	11	6
Beef loin steaks	CO mixture	14	7
	High $O_2$	10	7
	Vacuum packs	14	7
Pork chops	CO mixture	21	14
	High $O_2$	14	7
	Mixture with $O_2$ absorber	17	10

<sup>a</sup> CO mixture = modified atmosphere of 0.4%  $CO_2$ /60%  $CO_2$ /40%  $N_2$ ; High  $O_2$  = modified atmosphere of 70%  $O_2$ /30%  $CO_2$ ; Mixture with  $O_2$  absorber = modified atmosphere of 60%  $CO_2$ /40%  $N_2$  with an  $O_2$  absorber in the pack.

### 3.3. Storage life of beef loin steaks

At 4°C, off-odours developed 4 days later in beef loin steaks in the CO mixture and in vacuum packs than in high  $O_2$  (Table 1). At 8°C, no differences in the development of off-odours were observed. Off-odours developed 4 to 7 days earlier in meat at 8 than at 4°C. The type of packaging did not significantly affect ( $p < 0.05$ ) the total viable counts on the meat, but the counts were significantly higher ( $p < 0.01$ ) at 8 than at 4°C after both 3 and 7 days of storage [Fig. 2(a)]. The numbers of *B. thermosphacta* were less than  $\log_{10} 4$  CFU/cm<sup>2</sup> in meat in all types of packaging at all times, but were significantly higher ( $p < 0.05$ ) on meat in high  $O_2$  at 7 and 10 days than on meat in the CO mixture and in vacuum packs at equivalent times [Fig. 2(b)]. The numbers of pseudomonads did not exceed  $\log_{10} 3.5$  CFU/cm<sup>2</sup> at any sampling time, and were not significantly affected ( $p > 0.05$ ) by the type of packaging or the storage temperature.

The colour of the beef loin steaks in the CO mixture was stable bright red throughout storage at both 4 and 8°C, as shown by the low colour scores and high  $a^*$  values [Fig. 2(c) and (d)]. Steaks in high  $O_2$  were also bright red with high  $a^*$  values at day 3, but these steaks discoloured gradually between days 3 and 10, significantly faster ( $p < 0.05$ ) at 8 than at 4°C. Meat in vacuum packs was slightly discoloured with low  $a^*$  values throughout storage. The colour scores and  $a^*$  values of vacuum packaged steaks were not significantly affected ( $p > 0.05$ ) by the storage temperature.




Fig. 1. Mean values ( $n = 5$ ) for (a) total viable counts, (b) numbers of *Brochothrix thermosphacta*, (c) visual colour scores and (d) CIE  $a^*$  values for ground beef stored in 0.4%  $CO_2$ /60%  $CO_2$ /40%  $N_2$  at 4°C (○) or 8°C (●), in 70%  $O_2$ /30%  $CO_2$  at 4°C (□) or 8°C (■), or in chub packs at 4°C (△) or 8°C (▲). Colour was assessed on a scale where 1 = bright red and 5 = extremely discoloured.

### 3.4. Storage life of pork chops

For pork chops, off-odours developed more slowly in meat in the CO mixture than in meat in the mixture with O<sub>2</sub> absorbers or in high O<sub>2</sub> (Table 1). Off-odours were detected 7 days earlier at 8 than at 4°C for chops in each type of packaging. The type of packaging did not affect the total viable counts on the pork chops [Fig. 3(a)]. However, the counts were greater on meat stored at 8 than at 4°C. The numbers of *B. thermosphacta* on chops in high

O<sub>2</sub> were significantly higher ( $p < 0.01$ ) than on chops in the CO mixture or in the mixture with O<sub>2</sub> absorbers after 7 days at 8°C or 10 days at 4°C, and reached approximately  $\log_{10} 6$  CFU/cm<sup>2</sup> [Fig. 3(b)]. The numbers of pseudomonads did not exceed  $\log_{10} 3$  CFU/cm<sup>2</sup> on any of the pork chops.

The colour of pork chops in the CO mixture was light bright red with high  $a^*$  values throughout storage [Fig. 3(c) and (d)]. Chops in high O<sub>2</sub> were red at day 3, but discoloured during storage, significantly faster ( $p < 0.05$ ) at




Fig. 2. Mean values ( $n = 5$ ) for (a) total viable counts, (b) numbers of *Brochothrix thermosphacta*, (c) visual colour scores and (d) CIE  $a^*$  values for beef loin steaks stored in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> at 4°C (○) or 8°C (●), in 70% O<sub>2</sub>/30% CO<sub>2</sub> at 4°C (□) or 8°C (■), or in vacuum packs at 4°C (▽) or 8°C (▼). Colour was assessed on a scale where 1 = bright red and 5 = extremely discoloured.




Fig. 3. Mean values ( $n = 5$ ) for (a) total viable counts, (b) numbers of *Brochothrix thermosphacta*, (c) visual colour scores and (d) CIE  $a^*$  values for pork chops stored in 0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub> at 4°C (○) or 8°C (●), in 70% O<sub>2</sub>/30% CO<sub>2</sub> at 4°C (□) or 8°C (■), or in 60% CO<sub>2</sub>/40% N<sub>2</sub> with O<sub>2</sub> absorbers at 4°C (○) or 8°C (◆). Colour was assessed on a scale where 1 = light bright red and 5 = extremely discoloured.

8 than at 4°C. Approximately 75% of the chops in high O<sub>2</sub> had black back bones at the time of sampling. Chops in the mixture with O<sub>2</sub> absorbers were moderately discoloured from day 3 to the end of storage. These chops had *a\** values similar to those of chops in high O<sub>2</sub>.

#### 4. Discussion

##### 4.1. Off-odour and microflora

The shelf life of the meat, as determined by the time to develop off-odours, was influenced by the packaging method, the storage temperature and the initial microbiological load on the meat. Storage of meat in the CO mixture, in vacuum packs or in chub packs gave the longest shelf lives. Meat stored in high O<sub>2</sub> generally developed off-odours 2-7 days earlier at 4 or 8°C than meat packaged in the other gas mixtures or by the other methods.

The differences in the rates of development of off-odours, as affected by the packaging method, were seldom related to any differences in numbers of total viable counts. However, the development of off-odours from the three meat types, especially ground beef and pork chops in high O<sub>2</sub>, coincided with the attainment of high numbers of *B. thermosphacta*. For ground beef, storage in the CO mixture retarded growth of *B. thermosphacta* even more than storage in chub packs. At chill temperatures above 1°C, *B. thermosphacta* often causes spoilage of meat stored in high O<sub>2</sub> atmospheres (Dainty & Mackey, 1992). High concentrations of CO<sub>2</sub>, removal of O<sub>2</sub> and low storage temperature inhibit the growth of *B. thermosphacta* (Gill, 1996; Nissen, Sørheim, & Dainty, 1996). Pseudomonads probably contributed to the off-odours of ground beef. Meat in high O<sub>2</sub> is often spoiled by *Pseudomonas* spp., but the growth of pseudomonads is retarded under anaerobic conditions (Dainty & Mackey, 1992; Gill, 1996). A shift in the metabolism of lactic acid bacteria under aerobic conditions can also produce off-odours (Nissen et al., 1996). In the present experiments, the numbers of coliforms or *E. coli* did not exceed log<sub>10</sub> 3 CFU/g or cm<sup>2</sup> in any samples. Therefore, those organisms probably did not contribute to off-odours.

For pork chops, the effect of CO on the microflora can be evaluated because the gas compositions of the CO mixture and of the mixture with O<sub>2</sub> absorber were identical, except for the inclusion of 0.4% CO in the former. Although a 4 day increase in the time to develop off-odours was observed with the CO mixture, there was no significant reduction in the microbiological counts. Luño et al. (1998) used 1% CO in high O<sub>2</sub> atmospheres and noted a delay in the onset of off-odours without any reduction in the numbers of psychrotrophic bacteria. However, Clark et al. (1976) found that the addition of

0.5-10% CO to N<sub>2</sub> atmospheres reduced the number of psychrotrophic bacteria and increased the odour shelf life of beef. For example, 1.0% CO in 99% N<sub>2</sub> increased the time to develop off-odours at 5°C from 18 to 24 days. The lack of such an effect of CO on bacteria in our experiments may be due to the use of 60% CO<sub>2</sub> overshadowing any effect of CO.

The use of CO makes it possible to dispense with O<sub>2</sub> and so to increase the CO<sub>2</sub> concentration in a MA to about 60%. Our data suggest that 0.4% CO probably has little or no direct effect on the growth of bacteria. Other studies have shown that increasing the CO<sub>2</sub> concentration from 20 to 100% increases the bacteriostatic effect of the gas, but the efficiency is highly dependent on low storage temperatures (Gill & Molin, 1991; Nissen et al., 1996). The high CO<sub>2</sub> concentration and absence of O<sub>2</sub> in the CO mixture will favour the growth of lactic acid bacteria, which usually cause a mild form of spoilage only late in the development of the spoilage flora (Gill, 1996).

The present experiments were performed at acceptable and abusive storage temperatures to assess the effects of temperatures commonly encountered in the distribution and sale of retail-ready meat. The storage temperature strongly affected the rates of growth of microflora and the time to develop off-odours. Consequently, independently of the packaging method, the shelf life of meat can be considerably extended by maintaining low temperatures in the chill chain (Gill & Molin, 1991; Nissen et al., 1996).

##### 4.2. Colour

The CO mixture gave a stable bright or light bright red colour with consistent high *a\** values for all three products, irrespective of the storage temperature. The initial level of residual O<sub>2</sub>, up to 0.5%, did not adversely affect the visual scores and instrumental values for the colour of meat stored in the CO mixture.

CO binds to myoglobin and forms cherry red carboxymyoglobin (El-Badawi et al., 1964). This pigment is spectrally similar to the bright red oxymyoglobin which normally develops at the surface of fresh meat in air. Carboxymyoglobin is less readily oxidized to brown metmyoglobin than is oxymyoglobin, because of the strong binding of CO to the iron-porphyrin site on the myoglobin molecule (Lanier, Carpenter, Toledo, & Reagan, 1978; Wolfe, 1980). Consequently, CO in concentrations of 0.5-2.0% enhances and stabilizes a bright red colour of meat (Kropf, 1980; Sørheim et al., 1997a). In a recent study, 1% CO in combination with 24 or 70% O<sub>2</sub> stabilized the colour of beef by reduced formation of metmyoglobin after storage at 1°C for up to 29 days (Luño et al., 1998). However, in a study of beef stored in a MA of 2% CO/78% CO<sub>2</sub>/20% N<sub>2</sub>, the colour of the meat was characterized as "too artificial" by

a sensory panel (Renner & Labadie, 1993). From our studies and experience from the Norwegian meat industry, 0.4% CO seems sufficient to produce a stable, attractive, bright red colour of meat.

All three meat types stored in high O<sub>2</sub> were bright red to red with high *a*\* values early in the storage periods, approaching the colour of meat in the CO mixture. As the microbiological counts of meat in high O<sub>2</sub> increased, the colour deteriorated, faster at 8 than at 4°C. Meat stored in a MA of high O<sub>2</sub> develops a thicker layer of oxymyoglobin than meat stored in air (Renner & Labadie, 1993). However, the oxymyoglobin gradually oxidizes to metmyoglobin, and the oxidation is faster at higher temperatures.

For cut bone, haemoglobin released from disrupted red blood cells in the marrow will accumulate at the surface and ultimately become black after the bone has been exposed to air or O<sub>2</sub> (Gill, 1996). Although bone blackening was not considered in the present visual colour evaluation, it can negatively affect the saleability of bone-in meat at retail display. The cut bones of pork chops stored in high O<sub>2</sub> blackened during storage, but this discoloration was not observed on bones in the CO mixture and the mixture with O<sub>2</sub> absorbers.

Beef loin steaks stored in vacuum packs were slightly discoloured with low *a*\* values at both 4 and 8°C. In these packs, meat juices were observed between the upper and lower films, but that did not influence the colour evaluations.

O<sub>2</sub> absorbers in packs with high CO<sub>2</sub> facilitate the removal of residual O<sub>2</sub> and maintain atmospheres free of O<sub>2</sub> during storage (Smith, Abe, & Hoshino, 1995). Low levels of residual O<sub>2</sub>, above 0.01–0.15% for beef and 0.5–1.0% for pork, will inevitably discolour the meat (Penney & Bell, 1993; Gill, 1996; Sørheim et al., 1997b). When no CO is present in an O<sub>2</sub> depleted MA, it is essential to remove the residual O<sub>2</sub> as fast and completely as possible to avoid formation of metmyoglobin. In these experiments, pork chops stored in the gas mixture with O<sub>2</sub> absorbers were moderately discoloured during the whole storage period at 4 or 8°C. Despite the obvious visible differences, these chops had similar *a*\* values to the chops in high O<sub>2</sub>. The discoloured surface made the chops unfit for sale, even in the early stage of storage. The present findings contrast with previous results, where the colour of porcine *m. longissimus thoracis et lumborum* was significantly improved by using O<sub>2</sub> absorbers in MAs of CO<sub>2</sub> with residual O<sub>2</sub> (Sørheim et al., 1997b). The present discoloration could be caused by incomplete use or function of the absorbers (Gill, 1996).

#### 4.3. Benefits and disadvantages of a MA with low CO/high CO<sub>2</sub>

An objection raised against using CO as a small component of a MA for retail-ready meat is the possi-

bility that the colour stability can exceed the microbiological shelf life, with the risk of masking spoilage of the meat (Kropf, 1980). Therefore, the consumer must evaluate the microbiological condition of meat in a CO mixture by off-odours. When a MA with CO is applied commercially, it is important to have a proper control of the hygienic condition of the meat raw materials and the chill chain temperatures.

CO used in concentrations below 1.0% does not present any hazard to the consumer, because consumption of meat packaged in such concentrations of CO will result in only negligible levels of carboxyhaemoglobin in the blood of consumers (Sørheim et al., 1997a). By delivering CO in a 1% mixture with 99% N<sub>2</sub>, which is the practice of Norwegian gas suppliers, CO is considered safe for use in the working environment. Other MAs with high levels of O<sub>2</sub>, up to 70%, must be regarded as explosive gas mixtures, which must be used with appropriate precautions for safety (Luño et al., 1998).

The suitability of gas mixtures and packaging methods for red meats for retail display depends on their ability to both reduce spoilage and stabilize colour. Gas mixtures with low concentrations of CO and high concentrations of CO<sub>2</sub> provide a combination of a long microbiological shelf life and a stable, bright red colour of meat. Meat packaged in a MA with high O<sub>2</sub> can achieve an initial bright red colour, but the microbiological shelf life and the colour stability are both considerably lower than those of meat in the CO mixture. Using CO eliminates the need to have O<sub>2</sub> as a component of the MA. Other MAs and packaging methods, like high CO<sub>2</sub> with O<sub>2</sub> absorbers, chub packs and vacuum packs may give a shelf life comparable to that of the CO mixture, but with a less acceptable colour or appearance of the meat. Thus, there appears at present to be no fully satisfactory alternative to the CO mixture used in packaging of retail-ready red meats in Norway.

#### Acknowledgements

The financial support of this study from the Research Council of Norway is highly appreciated. Vestfold-Buskerud Slakteri A/L, Sem and Hydrogas AS Utviklings-senter, Porsgrunn, are greatly thanked for packaging of the meat. We appreciate the gift of Ageless® O<sub>2</sub> absorbers from Cryovac Europe, Norderstedt, Germany. The technical staff and Per Lea (statistics) at MATFORSK are thanked for their skilful assistance in the study.

#### References

- Clark, D. S., Lentz, C. P., & Roth, W. D. (1976). Use of carbon monoxide for extending shelf-life of prepackaged fresh beef. *Canadian Institute of Food Science and Technology Journal*, 9 (3), 114–117.

- Cornforth, D. (1994). Color—its basis and importance. In: Pearson, A. M., Dutson, & T. R. (Eds.), *Quality Attributes and their Measurements in Meat, Poultry and Fish Products*, Blackie, London, pp. 34-78.
- Dainty, R. H., & Mackey, B. M. (1992). The relationship between the phenotypic properties of bacteria from chill-stored meat and spoilage processes. *Journal of Applied Bacteriology Symposium Supplement*, 73, 103S-114S.
- de Man, J. C., Rogosa, M., & Sharpe, M. E. (1960). A medium for the cultivation of lactobacilli. *Journal of Applied Bacteriology*, 23, 130-135.
- El-Badawi, A. A., Cain, R. F., Samuels, C. E., & Angelmeier, A. F. (1964). Color and pigment stability of packaged refrigerated beef. *Food Technology*, 18(5), 159-163.
- European Parliament and Council Directive (1995). Food additives other than colours and sweeteners. Official Journal of the European Community L61 (18.3.95), 1-38.
- Gee, D. L., & Brown, W. D. (1978). Extension of shelf life in refrigerated ground beef under an atmosphere containing carbon dioxide and carbon monoxide. *Journal of Agricultural and Food Chemistry*, 26, 274-276.
- Gill, C. O. (1996). Extending the storage life of meats. *Meat Science*, 43 (Suppl.), S99-S109.
- Gill, C. O., & Molin, G. (1991). Modified atmospheres and vacuum packing. In: Russel, N. Y., & Gould, G. W. (Eds.), *Food preservatives*, Blackie, London, pp. 172-199.
- Kropf, D. H. (1980). Effects of retail display conditions on meat color. *Proceedings of the Reciprocal Meat Conference*, 33, 15-32.
- Lanier, T. C., Carpenter, J. A., Toledo, R. T., & Reagan, J. O. (1978). Metmyoglobin reduction in beef systems as affected by aerobic, anaerobic and carbon monoxide-containing environments. *Journal of Food Science*, 43, 1788-1792, 1796.
- Luño, M., Beltrán, J. A., & Roncalés, P. (1998). Shelf-life extension and colour stabilisation of beef packaged in a low O<sub>2</sub> atmosphere containing CO: loin steaks and ground meat. *Meat Science*, 48, 75-84.
- National Live Stock and Meat Board (1991). Guidelines for meat color evaluation. *Proceedings of the Reciprocal Meat Conference*, 44, 232-249.
- Nissen, H., Serheim, O., & Dainty, R. (1996). Effects of vacuum, modified atmospheres and storage temperature on the microbial flora of packaged beef. *Food Microbiology*, 13, 183-191.
- Penney, N., & Bell, R. G. (1993). Effects of residual oxygen on the colour, odour and taste of carbon dioxide packaged beef, lamb and pork during short term storage at chill temperatures. *Meat Science*, 33, 245-252.
- Renner, M., & Labadie, J. (1993). Fresh red meat packaging and meat quality. *Proceedings of the International Congress of Meat Science and Technology*, 39, 361-387.
- Smith, J. P., Abe, Y., & Hoshino, J. (1995). Modified atmosphere packaging—present and future use of gas absorbents and generators. In: Farber, J. M., & Dodds, K. L. (Eds.), *Principles of Modified-Atmosphere and Sous Vide Product Packaging*. Technomic Publishing, Lancaster, PA, pp. 287-323.
- Serheim, O., Aune, T., & Nesbakken, T. (1997a). Technological, hygienic and toxicological aspects of carbon monoxide used in modified-atmosphere packaging of meat. *Trends in Food Science & Technology*, 8, 307-312.
- Serheim, O., Erlandsen, T., Nissen, H., Lea, P., & Høyem, T. (1997b). Effects of modified atmosphere storage on colour and microbiological shelf life of normal and pale, soft and exudative pork. *Meat Science*, 47, 147-155.
- Wolfe, S. K. (1980). Use of CO- and CO<sub>2</sub>-enriched atmospheres for meats, fish and produce. *Food Technology*, 34(3), 55-63.



SWEDISH MEATS

[logo]

Date  
April 9, 1999

Ref. No.  
33

Reference:  
Lars Wedén, (+46) 8-725 81 03

Date Ref. No.

Svein E. Skorstad  
Norsk Kjott [Norwegian Meat]  
P. O. Box 60 Refstad  
N-0513 Oslo 5 NORWAY

RECEIVED:	<i>April 14, 1999</i>
CASE WORKER:	<i>K. Framstad</i>
FILE NO.:	<i>561</i>
J. NO./DOC. NO.:	<i>99/00721</i>

Brother:

Please refer to our conversation at the Scandinavian Butchers' Association meeting on March 19 regarding CO as a packaging gas. Here at Swedish Meats we are committed to creating opportunities to increase the meat packaging done by the manufacturer in Sweden. An approval of CO as a gas for use in foods would entail significantly greater possibilities of brand profiling of meat products in the future, which could help improve our service to the retailers. Moreover, efficient factory packaging could even reduce costs across the entire distribution chain ending with the consumer.

We therefore support Norsk Kjott's proposal to submit a joint application to the EU Commission for the addition of CO to EU's list of additives. The plan is to submit a joint application in June 1999.

Sincerely,

Swedish Meats

[signature]  
Lars Wedén

[letterhead information]



DANSKE SLAGTERIER [Danish Bacon and Meat Council]	Axelborg, Axeltorv 3 1609 Copenhagen V Denmark Tel. (+45) 33 11 60 50	Fax (+45) 33 11 68 14 Telex 22 975 ds dk Telegram "danslagt"	[Logo]
--	--	--	--------

February 11, 2000  
Tel. (+45) 33 73 25 68

Norsk Kjøtt [Norwegian Meat]  
att: Jorunn Vormeland  
PB 360 Økern  
N-0513 Oslo  
Norway

[Stamp: ]Received: *February 14, 2000*  
Case Worker: *JVO*  
File No.: *562*  
J. No./Doc. No. *00/00204*

#### Use of CO as Packaging Gas

In Norway, CO has so far been used in very minute quantities together with other packaging gases for gas packaged cuts of meat and ground meat. This packaging gas has not been approved in the EU yet, and has not been used in Denmark this far.

CO and the other packaging gases help ensure the storage life and color stability of fresh meat, which are important for central packaging and distribution of meat. The trend in Denmark is toward increased central packaging of fresh meat, since this is both efficient and ensures high microbiological quality. It would therefore be interesting to take advantage of the positive Norwegian experiences with the use of CO in this country, as well.

Published research shows that the risk of growth of a range of pathogenic bacteria is the same or reduced when using CO in combination with the traditional packaging gases. The use of this gas can thus help improve food safety (Food Microbiology and Food Safety into the Next Millennium, Proceedings of the Seventeenth International Conference of International Committee on Food Microbiology and Hygiene, Netherlands, 1999).

The use of CO in the given concentration of 0.3–0.5% should not represent any toxicological risk to consumers, CO is generally supplied in a concentration of 1% in a mixture with either N<sub>2</sub> or CO<sub>2</sub> and does therefore not represent any workplace hazard to operators during the packaging process.

We are not aware that the use of CO was discussed in the process of drafting Directive 95/2/EU of February 20, 1995 concerning additives other than colorings and sweeteners. This may be due to the fact that no country had expressed any interest in using this kind of gas at the time.

Since there are advantages to the use of CO as a packaging gas, as mentioned, and since there are no negative effects to either consumers or packaging operators, DANSKE SLAGTERIER can support an



application to the EU Commission to include CO on the list of approved additives, possibly limiting the amount.

Sincerely,

DANSKE SLAGTERIER

[Signature]

Anne Birgitte Lundholt

(Managing Director)

000145



(10)

Asociación de Industrias  
de la Carne de España

## TELEFAX

To: Dirk Dobbelaere  
Secretary (CLITRAVI)

Subject: CO gas at meat packaging gas

Date: 14 of February 2000

Nº of pages: 1

Dear Dirk:

After reading the scientific documents that Mr. Truls Nesbakken handed out in the last T&L working group meeting. We will support the Norwegian proposal for authorization CO gas as a packaging gas within the UE.

We look forward to hearing from you soon.

Yours sincerely,

Miryam de Miguel  
Dpto. Calidad-AICE

For info  
To Norwegian CLITRAVI member  
From CLITRAVI

---

General Rodrigo, 6 • 28003 MADRID  
Teléfono: 91 554.70.45 - Fax: 91 554.78.49 - E-mail: aice@aice.es

000146

21.2.2000

Truls Nesbakken  
Fagsentret for kjøtt  
PB 396 Økem  
0513 Oslo  
Norge


Use of CO as packaging gas for meat and meat products

After going through the scientific documents sent to us and having own projects supporting the results, our institute is ready to support the Norwegian proposal to allow CO gas as a packaging gas in EU.

Finnish Meat Research Institute



Markku Raevuori  
Managing Director



Raymond Tuominen  
Laboratory Director

# The Use of CO as a Packaging Gas for Fresh Meat.

By Magne Yndestad

A previous report on the use of CO as a packaging gas concluded that there is unsatisfactory documentation on factors such as the development of pathogenic bacteria in the gas mixture in question (0.4% CO/60% CO<sub>2</sub>/40% N<sub>2</sub>). The Norwegian Research Center for Meat forwarded recent and complementary documentation on September 13, 1999.

Following scientific review, the content of this documentation can be summed up as follows:

## Bacteriological Conditions

Numerous studies have been undertaken regarding trial storage involving concentrations of CO<sub>2</sub> in a range consistent with the "Norwegian" mixture (60-75% CO<sub>2</sub>). Moreover, there are articles documenting the bactericidal effect of various other concentrations of gas mixtures containing CO<sub>2</sub>. The conclusion to these trials is the following:

The low CO concentration (<0.5% CO) has no apparent effect on bacterial flora in products packaged with gas. This also holds true for N<sub>2</sub> (filler gas).

Concentrations of CO<sub>2</sub> below 5% may stimulate the growth of certain types of bacteria. Between 5 and 50%, we see an approximately linear inhibiting effect. This effect is somewhat significant, since the inhibition of growth of the sensitive flora is as much as 50% at 10% CO<sub>2</sub>. The documented effect of CO<sub>2</sub> in high concentrations primarily applies to the psychrotrophic flora, including the most important spoilage bacteria.

As for the pathogenic bacteria, scientific literature in general points to the same tendency, i.e. inhibition of growth at both 4°C and higher temperatures (e.g. 10° C).

In comparison to other packaging methods or gas mixtures used, the mixture in question seems favorable both in terms of storage life and in terms of the relevant pathogens.

Following the last round of applications, The Norwegian Research Center for Meat has performed a relatively extensive study on freshly ground meat packaged in 0.3-0.5% CO/60-70% CO<sub>2</sub> and 30-40% N<sub>2</sub>. Various pathogens, such as *E.coli* O157:H7, *Listeria monocytogenes*, and *Yersinia enterocolitica* were tested in this trial. The Research Center has evaluated factors such as the important possibility that the strong suppression of the general psychrotrophic flora may favor certain pathogens, which will not be inhibited to the same degree. The main conclusion, however, is that the aforementioned pathogens are inhibited both at 4°C and 10°C. Comparing the CO packaging method to packaging employing a high concentration of O<sub>2</sub> or vacuum, shows that the risk for growth of the applied pathogens is identical or lower when packaging with CO.

The Research Center has studied the circumstances concerning salmonella bacteria and the gas mixture in question in the same products. Since none of the cultures grew at +4°C, studies were only undertaken at 10°C.

In this case, storage with packaging gas containing CO performed worst with regard to *S. dublin*, *S. enteritidis* and *S. diarizonae*, as a relatively strong growth occurred following Day 2. *S. typhimurium* too had considerable growth, although "sausage" packaging scored lower.

000148



This is completely in line with what is known about a whole range of salmonella bacteria in foods, i.e. that they hold up very well when competing with other bacteria, and also grow very well at temperatures around 8-10°C.

These facts emphasize the importance of cooling regardless of what packaging method is chosen.

#### Sensorial Circumstances

The last report pointed to the particular fact that the CO packaged meat could retain a fresh red color for days after spoilage set in. Hence, the consumer cannot see whether the meat he or she buys is spoiled, as opposed to fresh meat packaged in other types of gas packaging.

The Research Center notes that when opening a package, the consumer will detect any spoilage odor, and hence not eat the product. This may be true, but it is a fact that many people won't react to any incipient decay when the products looks completely "fresh." However, the packaging method for which approval is sought is meant for fresh meat that will be treated with heat prior to use. This is an additional safety factor that is important in a comprehensive evaluation.

#### Conclusion

The first bacteriological/sanitary statement made was based on the documentation available at the time. The new data and other relevant information from scientific literature indicate that there is sufficient evidence that the use of CO as a packaging gas as described in the application won't result in any increased risk of transmittal of food-borne diseases among consumers.

000149



[Handwritten:]

*From the report "Fresh Meat in Consumer Packaging" with modified gas containing CO<sub>2</sub> [illegible]*

**IV. Report by Tore Aune: "Fresh Meat in Consumer Packaging – A Toxicological Evaluation of the Use of up to 0.5% CO in a Gas Mixture."**

000150





# FRESH MEAT IN CONSUMER PACKAGING – A TOXICOLOGICAL EVALUATION OF THE USE OF UP TO 0.5% CO IN A GAS MIXTURE

By Tore Aune

Carbon monoxide (CO) is a colorless gas that is primarily generated by incomplete combustion of organic material. The background concentration of CO in the atmosphere is approximately  $0.01\text{--}0.09\text{mg/m}^3$  ( $0.009\text{--}0.08$  ppm), while the concentration in larger cities may exceed  $50\text{mg/m}^3$  as an 8 hour mean, depending on traffic.

## General Health Effects

CO attaches to the iron of the hemoglobin in the red blood cells during generation of carboxyhemoglobin (COHb), and can thus affect the transport of oxygen in the blood and the supply of oxygen to the tissues. Compared to its affinity to oxygen, hemoglobin has approximately 240 times greater affinity to CO. CO also attaches to myoglobin, cytochromes, and some other enzymes, but these reactions are considered less important than the formation of carboxyhemoglobin (WHO 1979). The health impact on humans is mainly restricted to effects on the cardiovascular system, the nervous system, and certain types of proteins and cells in the bloodstream, as well as effects on embryos (SFT 1992).

The carboxyhemoglobin percentage (COHb %) is a function of the CO concentration in the inhaled air, the exposure time and the level of physical activity (Coburn et al., 1965) (see Table 1). A CO exposure resulting in a COHb concentration above 2% in the bloodstream of the most sensitive individuals (cardiovascular patients) has been shown to give symptoms of localized oxygen deficit and chest pains. Reduced work capacity occurs at a somewhat higher COHb%, and persons suffering from angina can tolerate less strain before an attack occurs. No health effects have been detected in healthy adults at COHb concentrations below 5%.

Table 1: Blood carboxyhemoglobin percentage as a function of CO concentration in air, exposure time and different degrees of physical activity (Coburn et al., 1965):

CO Conc.	Exposure Time in Hours:	COHb%		
		At rest	Moderate Activity	Strenuous Activity
$10\text{ mg/m}^3$	8	1.3	1.4	1.4
$25\text{ mg/m}^3$	1	1.0	1.5	2.0
$40\text{ mg/m}^3$	1	1.3	2.2	2.9

000151



CO attachment to the hemoglobin is reversible. The half-life at ventilation at rest is approximately 4.5 hours.

A small amount of CO is continually formed in the body as a result of the decomposition of substances such as hemoproteins. This results in a COHb% of approximately 0.5. The uptake of CO through inhalation comes in addition to that. The average COHb level in non-smokers is estimated at 1.2-1.5%, while the level is 3-4% in smokers.

#### *Survey of Health Effects Associated with CO Exposure*

The negative health effects of CO are due to the fact that CO competes with oxygen for points of attachments on the hemoglobin molecule. Moreover, the release of oxygen in the tissues is reduced (WGHO 1987). Myoglobin is closely related to hemoglobin. It stores oxygen and promotes the diffusion of oxygen to muscle cells. In cardiac and skeletal muscles, myoglobin binds CO with an affinity that is 30-50 times higher than the corresponding affinity for oxygen. No reported studies have shown that the binding of CO to myoglobin can cause any health effect at a COHb level of 4-5%.

Uptake and liberation of CO occur at a relatively slow pace (hours), which means that short-time exposure to elevated CO levels will not result in any noticeable increase in the COHb level. SFT report No. 92/16 (1992) includes an overview of the correlation between blood COHb levels and health effects (Table 2).

Table 2: Correlation between blood carboxyhemoglobin levels and health effects (SFT 1992):

COHb%	Observed Effects in Humans:
50 and above	Unconsciousness, lethal when untreated.
30 and above	Headache, dizziness, nausea, and vomiting.
10 and above	May be lethal to cardiovascular patients. Headache in healthy individuals.
5 and above	Reduction of peak oxygen consumption under strenuous activity in healthy individuals.
5 and above	Impaired vision, learning ability and fine motor response.
5 and above	Exposure during pregnancy may affect the embryo.
2.9 and above	Individuals suffering from angina can tolerate less strain before an attack occurs.
2.3 and above	Reduced capacity for physical work, especially stamina.
2 and above	Possible reduced ability to concentrate and pay attention.
2 and above	Symptoms of localized oxygen deficit and incipient chest pains in cardiac patients.

The literature in the field does not seem to indicate that health effects have been proven in healthy adults exposed to CO resulting in a blood COHb concentration of less than 5%.

000152



However, the data indicate that a COHb level of 2-3% may have negative effects on sick and sensitive individuals, such as people suffering from cardiovascular diseases.

#### *Exposure to CO through the Air*

With regard to CO as an air pollution factor, a team of Norwegian experts (SFT 1992) suggested air quality criteria at CO concentrations resulting in a maximum of 1.5% COHb during light physical activity (including the CO produced endogenically). The correlation between CO concentration, activity level, and exposure time in order not to exceed 1.5% blood COHb is shown in Table 3.

Table 3: Calculation of CO concentrations in the air resulting in a COHb level of 1.5%, including endogenic CO production (SFT 1992):

Exposure Time:	CO Concentration, mg/m <sup>3</sup>		
	At Rest:	Moderate Physical Activity:	Strenuous Physical Activity:
15 min	170	80	52
30 min	86	42	29
1 hour	48	24	18
8 hours	11.5	9.2	9.2

#### *Exposure to CO through Consumption of Fresh Meat Treated with a Gas Mixture*

There is a paucity of information in scientific literature concerning exposure to CO through the consumption of fresh meat treated with a gas mixture containing CO. One of the most interesting references in this regard is a 1954 publication by A. L. Tappel et al., which is unfortunately not easily accessible. However, their work has been cited in other publications, e.g. in the study by Clark et al. (1976): Tappel et al. considered a US industrial sanitary norm for CO of 50 ppm (8 hours/day), and found that such exposure would result in a blood COHb level over a longer period of time that is approximately 14 times higher than the temporary increase caused by consumption of approximately 225 g meat, provided that the myoglobin and hemoglobin in the meat are saturated with CO, and that 100% of CO from this source is transferred to the blood of the consumer (an estimate representing a hypothetical worst-case scenario). According to the authors, such treatment of meat will thus cause only a very minor effect in comparison to what is considered the safety limit, even when assuming maximum uptake of CO. Watts et al. (1978) exposed beef to a gas containing 1% CO for 3 days, and found that this resulted in a CO saturation of approximately 30% of the myoglobin. CO was lost under such storage conditions, with a half-life of approximately 3 days. After cooking, the CO concentration in the meat decreased to

000153



below 0.09 ppm (equivalent to approximately 0.1 mg/kg). Maximum loss after cooking (on burner at 195°C) amounted to approximately 85%.

#### *Comparison of CO Exposure through Air and Meat (CO Treated)*

There is little data available for such a comparison, but a rough overview nevertheless provides some points of reference. An adult inhales 10-20m<sup>3</sup> air per 24 hours (depending on the activity level). This is the equivalent of 0.42-0.84m<sup>3</sup> per hour (or 3.36-6.72 m<sup>3</sup> per 8 hours).

To stay within a maximum blood COHb level of 1.5%, the CO concentration in the air must be 24 mg/m<sup>3</sup> for 1 hour at moderate physical activity, at 9.2 mg/m<sup>3</sup> for 8 hours (according to Table 3). In comparison, the CO exposure is 0.1 mg/kg after consumption of 250 g of heated CO-treated meat that has been treated for 72 hours in a gas containing 1% CO (Watts et al., 1978). Table 4 shows a calculation of CO intake from the air and a meal of CO-treated meat.

Table 4: Comparison of CO intake from air within a range without any health impact and theoretical intake of CO through consumption of a meal of CO-treated meat:

Path of Exposure:	CO Intake, 1 hour:	CO Intake, 8 hours
Lungs (15 m <sup>3</sup> /24 hours)	24mg x 0.625 = 15.1mg	9.2mg x 5 = 46.0mg
Meat	0.025mg	0.025 mg

For CO balance between air and blood is only achieved after a considerable period of time (hours). The absorption of gases from the intestinal canal to the blood is probably considerably less efficient than from the lungs, where the tissue allows for maximum gas exchange between the alveoli and the bloodstream. This implies that intake of CO through meat probably won't cause any demonstrable increase in the blood CO level (in the form of COHb). And at any rate, the exposure from meat is much lower (approximately one thousand times lower) than through the airways, as shown in the calculations above.

According to the Norwegian Institute of Air Research (SFT 1992), the CO concentration in larger Norwegian cities is on average between 1 and 2 mg/m<sup>3</sup> during the winter. Maximum hourly values have been measured to approximately 60 mg/m<sup>3</sup>, and maximum values for 8 hours to about 40 mg/m<sup>3</sup>.

#### *Evaluation of Other Gases Used in Foods in the EU*

EU's Research Committee on Foods (SCF) has not considered CO. However, the expert team has considered other gases (EUR 1981), such as carbon dioxide (CO<sub>2</sub>) and nitrogen oxide (NO). In this connection, the committee employed the following evaluation method, which should be applicable for CO, as well:

000154



CO<sub>2</sub>: This compound is a natural product of metabolism, and people are constantly exposed to carbon dioxide from the atmosphere, food and drink. Compared to this exposure, the residual content from its use as an extraction agent is insignificant. Establishing an ADI for this compound is unnecessary. The committee considers this compound acceptable as an extraction agent. It is unnecessary to determine concentration values for the residue.

N<sub>2</sub>O: The pharmacological and pharmacokinetic properties of this gas are well known from the extensive use of N<sub>2</sub>O as an anaesthetic. Even though no data on residual content are available, such amounts are probably so minor that they are not hazardous to the consumer. The committee finds that it is unnecessary to establish an ADI, and considers the use of N<sub>2</sub>O as an extraction agent acceptable.

#### *Toxicological Evaluation of the Use of CO as a Packaging Gas for Meat*

People are continually exposed to carbon monoxide, both by means of endogenic production and by inhaled air. Toxicologically, it is the amount of CO bound to the blood hemoglobin (the carboxyhemoglobin percentage) that determines any health effects. The very first effects in sensitive individuals occur at COHb concentrations from approximately 2-3%. To prevent possible health effects even in the most sensitive individuals, a team of Norwegian experts has suggested limits for CO in the air that do not result in COHb concentrations above 1.5%, including the endogenic production at 0.5%. The above-mentioned estimates indicate that even if all CO in the prepared meat is transferred to the consumer's blood, the CO concentration – even a temporary concentration – will remain well below accepted limits in air. From a health perspective, the use of CO in concentrations below 0.5-1 % for fresh meat thus represents no toxicological risk.

#### *References:*

- Clark, D.S. C.P. Lentz, and L.A. Roth: "Use of carbon monoxide for extending shelf-life of prepackaged fresh beef." Can. Inst. Food Sci. Technol. J., 9, 114-117, 1976.
- Coburn, R.F., R.E. Forster, and P.B. Kane: "Considerations of the physiological variables that determine the blood carboxyhemoglobin in man." J.Clin.Invest., 44, 1899-1910, 1965.
- EUR, "Food Science and Techniques." Reports from the Scientific Committee on Foods (Series 11), EUR 7421, Luxembourg, 1981.
- SFT, "Health and environmental effects of air pollution. Recommended air quality criteria." SFT report No. 92:16, 1992.
- Watts, D.A., S.K. Wolfe., and W. D. Brown: "Fate of [<sup>14</sup>C] carbon monoxide in cooked or stored ground beef samples." J.Agric. Food Chem., 26, 210-214, 1978.

000155



WHO, Environmental Health Criteria 13. Carbon monoxide, World Health Organization, Geneva, 1979.

WHO, Air Quality Guidelines for Europe, World Health Organization Regional Publ., European Series, No. 23, 1987.

000156



**ATTACHMENT 4**

Final Report

**EVALUATION OF BEEF STEAKS AND GROUND BEEF IN THE PACTIV  
ACTIVE TECH PACKAGING SYSTEM:  
EFFECTS OF CARBON MONOXIDE IN THE PACKAGE ATMOSPHERE**

for

Pactiv Corporation  
Canandaigua, NY  
May 2001

*Prepared by:*

Kathy Hachmeister, MS  
Melvin Hunt, PhD  
Weber Hall  
Dept of Animal Sciences & Industry  
Kansas State University  
Manhattan, KS 66506

Phone: 785-532-1232  
Fax: 785-532-7059

George Milliken, Ph.D  
Dickens Hall  
Dept of Statistics  
Kansas State University  
Manhattan, KS 66506

Phone: 785-532-0514

TABLE OF CONTENTS

Project Summary .....	3
-----------------------	---



Introduction.....	4
Hypothesis and Objectives .....	4
Experimental Procedures .....	5
Results and Discussion .....	8
Tables and Figures.....	at back

## PROJECT SUMMARY

The effects of carbon monoxide (CO) in Active Tech modified atmosphere packages (MAP) were determined for:

- A. Initial product color,
- B. Stability of color during display, and
- C. Relationships of color deterioration and microbial populations.

Steaks from three beef cuts (strip loin, tenderloin, and inside round steaks) and ground beef were packaged in a MAP certified gas blend (0.4% CO, 30% carbon dioxide and 69.6% nitrogen) and stored at 35° or 43°F for up to 35 days. Cuts then were removed from MAP and displayed at 34°F until their color was approaching consumer unacceptability. Color and microbial parameters were measured and compared to baseline data of comparable product exposed to oxygen but not CO.

A fundamental goal of this research was to determine if CO extended the color life of beef cuts and ground beef beyond their microbial soundness, i.e., did color mask spoilage.

## CONCLUSIONS

- The Active Tech MAP system containing CO in the gas blend produced products that were equally as red as products packaged with traditional oxygen permeable overwrap.
- Improvement in visual appearance especially in the tenderloin and inner portion of the inside round steaks were observed on day zero of display and throughout display.
- Color of products exposed to CO was a typical, bright red when the outer MAP bag was removed and products were allowed to bloom for 60 to 90 minutes.
- Color declines for products stored in MAP with CO compared well to baseline products exposed to oxygen. Hence, a typical discoloration pattern was seen in both baseline and MAP studies.
- Color life for tenderloin and inside round steaks (and to a lesser extent ground beef) was slightly longer than their baseline counter parts, especially when stored 35°F vs. 43°F.
- Although microbial growth curves changed in slope and exponential growth based on the environment in the packages, bacterial growth was neither encouraged nor suppressed by the addition of CO to the MAP gas blend.
- Aerobic bacteria and facultative anaerobes followed typical patterns of growth contingent upon the environmental conditions.
- Effects of storage temperature (35° vs. 43°F) and increased storage time (21 or 35 days) resulted in typical redness decline, increase in off-odors and microbiological changes.
- CO neither masked spoilage nor resulted in color life extension beyond the point of microbial soundness.

## INTRODUCTION

Marketing of case-ready meats has moved beyond the concept stage to reality. This method of delivering meat to retailers is expected to be the predominate system within five years. Some of the largest retailers are already paving the way for this makes-sense marketing system.

Modified atmosphere packaging (MAP) systems are a necessity for case-ready meats because current retail meat over wrapping does not fulfill requirements for shelf life and other needs. Processors can choose either high-oxygen or low-oxygen MAP for retail-ready meats. Both systems rely upon the meat having certain functional properties needed to optimize delivery of cuts with excellent display color life and sound microbial quality.

In low-oxygen MAP, such as the Active Tech System of Pactiv Corporation, it is essential that the meat achieve a stable red color that extends throughout storage and display. This usually is accomplished by modifying the package atmosphere so that the meat pigment returns to its purple-red state (deoxymyoglobin). Then, at display, packaged cuts are re-exposed to oxygen (air) to re-form a bright-red color (oxymyoglobin). Some muscles can easily accomplish this function whereas other muscles have a difficult time – due principally to short comings of their inherent muscle chemistry. Thus, novel ways to aid in obtaining desirable color during storage and display would be beneficial.

Gas atmosphere composition plays a critical role in the functionality and efficacy of MAP systems for meat. The atmosphere affects one or more of the following: product appearance, shelf life, microbial and palatability issues, gas dynamics, purge, and myoglobin functionality. Typical atmospheres for low-oxygen MAP utilize carbon dioxide (CO<sub>2</sub>) and/or nitrogen (N<sub>2</sub>) prior to the meat being re-exposed to oxygen. Addition of small amounts of carbon monoxide (CO) to a CO<sub>2</sub> and/or N<sub>2</sub> atmosphere could aid in producing a more functional pigment color in MAP, especially in meat cuts known to have lower color stability. CO is well known for its ability to bind to myoglobin and form a bright, crimson-red colored pigment known as carboxymyoglobin. However, carboxymyoglobin is believed to stabilize meat color beyond its microbial shelf life. Consequently, consumers may not be able to rely on color as an indicator of quality at time of purchase. Research is needed to address the use of low levels of CO in a MAP system.

## HYPOTHESIS AND OBJECTIVES

This research was based on the hypothesis that a small quantity (<0.5%) of CO combined with the typical gases of MAP (CO<sub>2</sub> and N<sub>2</sub>) would produce meat color complimentary to the quality needs of a case-ready meat delivery system without compromising consumer quality issues. More specific objectives evaluated the effects of CO in the Active Tech System for:

- The initial color of intact muscles and ground beef – this objective addressed color differences between meat in MAP containing CO vs. packaging in O<sub>2</sub>.
- The color deterioration of these products during display -- these data defined the color display stability of meat in MAP containing CO vs. packaging in O<sub>2</sub>.
- The microbial profile of the meat stored with or without mild temperature abuse – this portion provided information about microbial growth with CO in MAP relative to the time-honored relationship between color deterioration and spoilage.

## EXPERIMENTAL PROCEDURES

This project involved two phases. The **Baseline Display Study** characterized the color and microbial traits of selected cuts and ground beef using typical oxygen-permeable packaging under typical retail display conditions. The **MAP Display Study** utilized the Pactiv Active Tech Packaging System in combination with a unique, certified gas blend (0.4% CO, 30% CO<sub>2</sub> and 69.6% N<sub>2</sub>) in the package atmosphere during storage conditions (pre-display).

The outer MAP bag was removed and the products were displayed in the same manner as the baseline samples. All data from the MAP Portion were compared to the Baseline product.

### RAW MATERIALS:

Twelve beef strip loins (NAMP #180 containing the *Longissimus* muscle), 18 tenderloins (NAMP #189A containing the *Psoas major* muscle), 12 inside rounds (NAMP #169A containing the *Semimembranosus* muscle), and 6 batches of ground chuck (80% lean) were obtained from a commercial source (Prairieland Processors, Inc., Kansas City, KS) at four to six days postmortem. Vacuum packaged subprimals and trim that were received at the KSU Meats Laboratory had an internal temperature of 34°F and had never been frozen. Prior to product preparation, subprimals were stored at 34°F. This product was allocated to 6 replications (2 each of the strip loins and inside rounds and 3 tenderloins constituted a replication).

### PRODUCT PREPARATION AND PACKAGING:

One inch thick steaks cut from each subprimal and ground beef formed into about one-pound blocks (Beef Steaker, Model 600, Hobart Corp., Troy, OH) were placed on Styrofoam trays (17S for strip loins, 4P for inside rounds, 1 for tenderloins, and 2P for ground beef) containing an absorbent pad (Ultra Zap Soakers, Paper Pak Products, La Verne, CA). Product was overwrapped with polyvinyl chloride (PVC) film (23,000ccO<sub>2</sub>/m<sup>2</sup>/24hrs; Filmco MW4, LinPac, UK or Omnifilm 4P, Huntsman, Salt Lake City, UT) using a mechanical wrapper (Filmizer Model CSW-3, Hobart Corporation, Troy OH) and was assigned randomly to either a **Baseline Display Study** using only PVC-wrapped packages or a **MAP Display Study** using the Active Tech System of Pactiv Corporation. Trays for MAP were placed individually in barrier bags (4.5ccO<sub>2</sub>/m<sup>2</sup>/24hrs: NXE 1-300, Alec Enterprises, Burnsville, MN) along with an oxygen absorber (MRM-200, Multisorb Technologies, Buffalo, NY) activated using Pactiv Active Tech Activator No.1. Barrier bags were evacuated, flushed with a certified gas blend containing 0.4% CO, 30% CO<sub>2</sub>, and 69.6% N<sub>2</sub>, and sealed (Freshvac Model A300, CVP Systems, Inc., Downers Grove, IL).

### TREATMENTS:

**Baseline Display Study:** Twelve packages of ground beef and one steak ( $\leq 1/8$ " fat trim) from each subprimal (12 strip loins, 12 inside rounds, 18 tenderloins, and the 6 batches of ground beef), were evaluated in a baseline study to establish the color and microbial parameters for meat never in MAP and exposed only to atmospheric oxygen. These packages were placed in display about 4 hours post-packaging (see display and measurement details below).

**MAP Display Study:** To test the effects of CO in MAP, one package of each product from each of 6 replications was selected at random for assignment to all possible combinations of two storage temperatures (35 and 43°F) and three storage times (7, 14, and 21 days for ground beef and 7, 21, and 35 days for steaks). The lower temperature represented reasonably good industry practice, and the higher temperature represented a mildly abusive storage conditions. The storage times represented current industry practice.

Prior to display (post-MAP), the O<sub>2</sub> and CO<sub>2</sub> levels in the outer barrier bags were measured using a MOCON head space analyzer (Pac Check™ Model 650, MOCON/Modem Controls, Inc., Minneapolis, MN).

#### **DISPLAY CONDITIONS:**

Meat samples were placed in simulated retail display at  $34 \pm 3^{\circ}\text{F}$  under 1614 lux (150  $\pm$  5 foot candles; Model 201, General Electric, Cleveland, OH) light intensity (Philips, 34 Watt, Ultralume 30) in open-top display cases (Unit Model DMF8, Tyler Refrigeration Corporation, Niles, MI). Cases were programed to defrost two-times per day at 12 hour intervals. Display case temperatures were monitored during display using temperature loggers (Omega Engineering, Inc., Stamford, CT). Display times varied based on product type, initial microbial loads, and storage conditions. Product was removed from display when the color score was deemed unacceptable by a visual panel (a color score of 3.5). Baseline products were displayed 7, 5, 4, and 3 days for strip steaks, inside rounds, ground beef, and tenderloins, respectively.

#### **VISUAL COLOR EVALUATION:**

Ten trained visual panelists evaluated color using a five-point scale where 1 = very bright red, 2 = Bright red, 3 = Slightly dark red or tan, 4 = Moderately dark red or tan, and 5 = Extremely dark red or brown. The cut-off score for consumer acceptable color was  $\geq 3.5$ .

Two portions of the inside round muscle were scored separately. The outer 1/3 portion (OSM) and the deep, inner 1/3 portion (ISM). The middle 1/3 area was not scored. The 10 panel scores were averaged for statistical analysis.

#### **INSTRUMENTAL COLOR AND SPECTRAL DATA:**

Samples were instrumentally analyzed for lightness (L\*), redness (a\*), and yellowness (b\*) for Illuminant D-65 (daylight) using a HunterLab MiniScan Spectrophotometer (1.25 inch diameter aperture, Hunter Associates Laboratory, Inc., Reston, VA). Multiple readings (2 to 4 depending on cut size) were taken and averaged for statistical analysis on each cut at each testing period.

#### **ODORS:**

At the end of display, each package from the MAP Display Study was evaluated for off odors by two experienced panelists using a 5-point scale where 1 = no, 2 = slight, 3 = small, 4 = moderate, and 5 = extreme off odor. A score of 3.5 was assumed to be unacceptable to consumers.

## **MICROBIOLOGICAL PROCEDURES:**

Microbial populations were estimated at the end of MAP storage (day 0 of display) and at the end of display (day of unacceptable color). For each post-display sample, a portion of the surface area (top surface) that had been exposed to light was excised. After each package was opened aseptically, two cores (ca 2 in<sup>2</sup>) were removed (approximately 1/8 inch depth), placed in a sterile stomacher bag, and blended two minutes with 0.1% peptone diluent. Serial dilutions of the homogenate were prepared in 0.1% peptone and appropriate dilutions were plated in duplicate on Aerobic Plate Count Petrifilm™ to determine total aerobic bacterial populations and on *E. coli* Count Petrifilm™ to estimate generic *E. coli* and total coliform bacterial counts. In addition, appropriate dilutions also were plated in duplicate on MRS agar to determine lactic acid bacterial populations. Aerobic Plate Count Petrifilm™ and *E. coli* Count Petrifilm™ (3M Microbiology Products, St. Paul, MN) were incubated at 90°F for 48 hours prior to enumeration. LAB populations were counted after 48 hours of 92°F incubation in a CO<sub>2</sub> chamber. Microbial detection limits for intact muscle and ground beef were 1.76 count/cm<sup>2</sup> and 5.0 count/gram, respectively.

### **pH:**

pH was determined on intact muscle and ground beef samples collected on the day of production. Ten grams of sample were added to 100 mL of distilled water and blended for two minutes. A standardized pH meter with an electrode was used to measure pH according to the procedure outlined in the Handbook for Meat Chemists.

### **FAT AND MOISTURE:**

Ground beef samples collected on the day of production were analyzed in duplicate for moisture and fat using AOAC procedures 985.14 and 985.15, respectively.

### **EXPERIMENTAL DESIGN AND STATISTICS:**

The design was a randomized complete block with six replications. A replication consisted of 1 to 3 subprimals (number depended on the size of each cut). Steaks cut from the subprimals and separate batches of ground beef trim were randomly assigned to replication and the treatment combinations. Data were analyzed using analysis of variance and significant differences determined using least significant difference tests at  $P < 0.05$ .

### **SAMPLING TIMES/PARAMETERS MEASURED:**

#### **1. MAP Gas Composition for oxygen and carbon dioxide levels**

- Subsample of several ActiveTech packages on production day (2-3 hours post-packaging) to verify gas composition being obtained
- End of MAP storage at two temperatures

#### **2. Microbiology:**

- Initial counts for subprimals and ground beef on the day of production
- End of MAP storage at two temperatures
- End of display

### 3. Visual Color:

- Initial color prior to display lighting
- End of MAP storage at each of two temperatures and after 60 to 90 min bloom at 34°F (equal to 0 time of display)
- Daily during display

### 4. Instrumental Color:

- Initial color = After packaging in PVC on production day for baseline data, minimal exposure to light
- End of MAP storage at each of two temperatures and after 60 to 90 min bloom at 34°F (equal to 0 time of display)
- Daily during display

### 5. Odor:

- At end of display (prior to microbial testing)

## RESULTS AND DISCUSSION

**The Baseline Study:** A random selection of all steaks and ground beef packaged in PVC film were placed in display to serve as a baseline for color and microbiological comparisons.

Products were expected to have the lowest microbiological load and ideal color stability using traditional packaging and display conditions for products exposed only to atmospheric oxygen. The inherent muscle chemistry responsible for good color life also was optimal. If the product exposed to CO were to have extended meat color life, then it will be compared to the baseline "control" with the "best" possible color.

**Color Reference Points:** The discussion below involves both visual and instrumental measures of color. Visual scores were considered the "standard" with instrumental color being discussed relative to its agreement or disagreement with the visual panel, ie, did the objective measurements confirm what the color panel saw. Visual scores of  $\geq 3.5$  were considered borderline acceptable. When samples reached this discoloration, they were removed from display. Normally,  $a^*$  values (higher values indicate more redness) are highly correlated to visual appraisal.

Inside round steaks typically are two-toned in color. The inner portion (ISM) is much less color stable compared to the outer portion (OSM). These portions were scored separately since one portion may have acceptable color while the other has unacceptable color that would be discriminated against by consumers resulting in the whole cut being judged

unacceptable in color. The effects of CO on this bi-colored muscle were needed to confirm that color was not excessively extended in either portion.

#### **FAT AND MOISTURE, pH, AND INITIAL MICROBIAL LOAD:**

Average fat and moisture contents of the ground beef were 19.5 and 61.6%, respectively. pH of both intact muscles and the ground beef ranged from 5.3 to 5.7. The initial aerobic plate counts and lactic bacteria counts for all products were relatively low and indicative of microbial quality of the raw materials and good sanitation. Furthermore, coliforms and *E. coli* were below the detection limit throughout the study.

#### **GAS COMPOSITION AT END OF MAP:**

At the end of MAP storage, each package atmosphere was analyzed for O<sub>2</sub> and CO<sub>2</sub> (Table 1). Only 6 (each from a different treatment combination) of 288 packages were removed from the experiment due to leakage.

#### **INITIAL PRODUCT COLOR AND APPEARANCE:**

The color of ground beef and steaks entering display (after MAP storage at 2 temperatures) was an attractive, typical red color. Although there were several significant differences in visual scores and a\* values (Table 2 and Figures 1-10 at day 0) for product in CO vs. baseline cuts, the variation in color was usually within  $\pm 0.5$  of a color score. In general, the initial color of product exposed to CO was very similar to the color of steaks from the baseline display (never exposed to CO). When differences occurred, they were more related to either storage temperature or postmortem age of the product.

Panelists did not consider the color of product exposed to CO atypical. Cuts exposed to CO generally appeared more uniformly bright-red and would be expected to have high consumer appeal. These results were expected, as CO is known to preferentially form a ligand with the colored pigment (myoglobin) in meat resulting in an intensely red pigment known as carboxymyoglobin. At higher levels of CO (0.4% vs. 0.6 to 1%) than used in this experiment, meat color has been described as being an unusual crimson, bright-red color compared to the normal red of oxymyoglobin.

A critical next question was whether the carboxymyoglobin formed on the surface was more stable than the oxymyoglobin formed in baseline product. Further, did the carboxy



pigment deteriorate in a predictable way that consumers could continue to use visual color to judge freshness or potential spoilage.

#### **COLOR DETERIORATION PROFILE:**

Visual panel scores (Figures 1-5) and instrumental color ( $a^*$  values, Figures 6-10) clearly showed that product exposed to CO during MAP storage had color deterioration during display. As expected, visual scores increased (color deteriorated) and  $a^*$  values decreased (loss of redness) as days in display increased.

In several instances, color appeared to improve late in display – as indicated by a decrease in visual scores (see ground beef, strips loins and tenderloins at 43°F). These decreases were not a return of redness. Rather the apparent decrease resulted from removal of discolored packages the preceding period, leaving product with less overall discoloration remaining in the case.

In general, the color deterioration profiles followed an expected pattern. Namely, the freshest product (baseline packages) had the most stable, red color and the most days in display needed to reach borderline discoloration (Table 3 scores to 3.5) of all treatments. Exceptions occurred for the inside portion of the inside round and tenderloin products, where the product exposed to CO had slightly more stable color than the baseline product (Table 3). These two muscle areas are well known by retailers as having short color life. Thus, CO appeared to improve color life when the inherent muscle chemistry desired for color was limited.

For product from MAP, the longer the storage time, the faster the deterioration, especially at the higher storage temperature (Tables 2 and 3). For packages stored at 43°F, which was a mildly abusive temperature, color deterioration would be expected to accelerate. This phenomenon also is illustrated in Figures 1-10.

Changes in  $a^*$  values (and other instrumental measures of color not shown) followed the same pattern of color deterioration observed by the visual panelists. There was no evidence that color shelf life was unexpectedly lengthened by exposure of meat to CO in MAP. The question remaining is whether the color life of product in CO masked spoilage, ie, were microbial counts higher than expected based on the degree of discoloration?

#### **COLOR DETERIORATION AND MICROBIAL GROWTH:**

**Baseline Display Study:** Initial, pre-display microbiological data suggested that the raw materials were fresh and processed using good hygienic practices. For intact cuts, lactic acid bacteria, generic *E. coli*, and total coliform counts were below the detection limit of 1.76 CFU/in<sup>2</sup>. Initial, pre-display APC for intact muscles ranged from 1 to 1.63 log<sub>10</sub> CFU/in<sup>2</sup>. Post-display counts were higher ( $P < 0.05$ ) than pre-display APC which was an increase in bacterial proliferation and typical deterioration. However, all product had sufficient microbes to be susceptible to spoilage.

Baseline products were pulled from display when the visual panel scores reached  $\geq 3.5$ . However, the APC did not exceed 5 log<sub>10</sub> CFU/unit as shown in Figures 11-14 and lactic bacterial did not exceed 6 log<sub>10</sub> CFU/unit as shown in Figures 15-18. Furthermore, off-odor scores for product at end of display (Table 3) ranged from no to slight off odor. Thus, color life in this base population did not exceed microbial soundness.

**MAP Display Study:** Similar trends in microbial growth occurred in post-displayed samples stored in MAP compared to baseline products. Microbial patterns for product deterioration are shown in Table 4 and Figures 11-18. Products stored under MAP at a slightly abusive temperature showed, as expected, a more rapid increase ( $P < 0.05$ ) in microbial counts compared to samples stored at 35°F. For post-MAP (pre-display) and post-display samples, APC were higher at 43°F than 35°F (Table 4), and during the later days of storage at the higher temperature, differences were more obvious. Significant changes ( $P < 0.05$ ) occurred in all cuts and ground beef with the exception of SM. Counts for the SM muscle were lower than expected and no significant changes occurring until day 35 of MAP storage. This suggests that quality products that have been handled in a sanitary fashion can be stored in the MAP system up to 35 days without comprising microbial quality. The APCs for intact strip loin and tenderloin steaks stored at 35°F were lower ( $P < 0.05$ ) on all days of display on days 21 and 35 post-MAP than steaks stored at 43°F (Figures 12 and 14). Although products did not show a difference in APCs 7 days post-MAP, those products stored at the higher temperature (43°F) were more inferior 21 and 35 days post-MAP.

**Did Color Mask Spoilage?** Central to this research was to evaluate the idea that the color of CO treated meat might mask spoilage. Food scientists generally agree that meat

color is seriously discolored when microbial counts approach  $\log 10^6$ , and that off odors frequently appear at counts of  $10^7$  to  $10^8$ . Numerous studies of ground beef, frequently the product with the highest counts, show that consumer-purchased retail product often has counts of  $10^5$  to  $10^8$ .

Visual color scoring was considered as the "standard" for determining the time to remove products from display. Because the visual panel scores were the deciding factor for length of shelf life, the interdependence between visual color and APC, LAB, and odor were considered quite important.

Figures 19-21 show aerobic and lactic bacterial growth and odor scores at the end of display plotted against their corresponding visual color scores. All data observations were summed over storage temperature, storage time, and product type and plotted in one graph.

If color masked spoilage, then there should be multiple points in the upper left quadrant of the plot, the area represented by unacceptable microbial counts and off odors but with acceptable color (i.e., scores  $<3.5$ ). This did not occur with any frequency in any of the three plots. Thus, it does not appear that exposure of meat to CO during extended (up to 35 days at either 35° or 43°F) caused meat color to hide spoilage.

Table 1 - Carbon Dioxide (CO<sub>2</sub>) and Oxygen (O<sub>2</sub>) Levels in MAP Packages of ground beef (GB) and steaks from strip loins (LD), inside round (SM), and tenderloin (TL).

Meat Cut	Storage Temperature, °F	Storage Time, days	CO <sub>2</sub> , %	O <sub>2</sub> , %
GB	35	7	28.4	0
GB	43	7	28.7	0
GB	35	14	27.7	0
GB	43	14	28.3	0
GB	35	21	27.4	0
GB	43	21	28.0	0
LD	35	7	33.3	0
LD	43	7	34.2	0
LD	35	21	32.4	0
LD	43	21	31.8	0
LD	35	35	31.1	0
LD	43	35	28.5	0
SM	35	7	28.9	0
SM	43	7	29.7	0
SM	35	21	27.9	0
SM	43	21	27.3	0
SM	35	35	26.8	0
SM	43	35	24.6	0
TL	35	7	34.3	0
TL	43	7	34.8	0
TL	35	21	33.6	0
TL	43	21	32.3	0
TL	35	35	32.5	0
TL	43	35	29.2	0

Table 2 - Means for initial visual color and a\* values for beef cuts exposed to carbon monoxide during storage at 35° and 43°F in Active Tech MAP vs. baseline cuts exposed only to oxygen.

Trait	Product	Baseline cuts	Time <sup>d</sup> in Active Tech MAP, days at 35° F		
			7	14 / 21	21 / 35
Initial Visual Color	GB	1.3a	1.6b	1.7b	1.8b
	LD	2.2b	2.5b	1.8a	2.2b
	ISM	1.8ab	2.0b	1.7a	2.0b
	OSM	2.6b	2.6b	1.9a	2.5b
	TL	1.9a	2.0a	1.9a	2.1a
Initial a* Values (redness)	GB	23.4a	25.6b	25.9b	25.6b
	LD	25.8a	25.7a	27.1ab	28.1b
	ISM	28.5a	26.9a	30.0a	29.4a
	OSM	27.4a	27.7a	29.8a	29.5a
	TL	23.6a	27.5b	30.0c	29.3c
			Time <sup>d</sup> in Active Tech MAP, days at 43° F		
Initial Visual Color	GB	1.3a	1.7b	1.8b	2.5c
	LD	2.2a	2.3a	2.1a	2.0a
	ISM	1.8a	1.8a	1.7a	2.4b
	OSM	2.6b	2.2a	2.2a	2.0a
	TL	1.9a	2.0ab	1.8a	2.2b
Initial a* Values (redness)	GB	23.4a	25.7b	25.1b	25.5b
	LD	25.8a	25.5a	28.7b	27.5b
	ISM	28.5a	28.7a	28.6a	27.5a
	OSM	27.4a	27.7a	30.2b	29.4ab
	TL	23.6a	27.8b	28.7b	26.4b

a-c Means in the same row with a different letter differ (P<0.05).

d Ground beef stored 7, 14, and 21 days, other muscles 7, 21, and 35 days.

Table 3 - Means for days to visual unacceptable visual color (score of 3.5) and odor at end of display for beef cuts exposed to carbon monoxide during storage at 35° and 43°F in Active Tech MAP vs. baseline cuts exposed only to oxygen.

Trait	Product	Baseline cuts	Time <sup>e</sup> in Active Tech MAP, days at 35° F		
			7	14 / 21	21 / 35
Days in display to unacceptable color	GB	3.6c	3.0b	3.0b	2.3a
	LD	6.2c	5.0b	5.2b	3.8a
	ISM	3.2a	4.8c	4.0bc	3.5ab
	OSM	4.8c	3.5b	3.4b	2.6a
	TL	2.6a	3.0b	3.2b	2.8ab
			Time <sup>e</sup> in Active Tech MAP, days at 43° F		
Days in display to unacceptable color	GB	3.6d	3.0cd	2.3b	1.5a
	LD	6.2d	5.0c	3.3b	2.3a
	ISM	3.2b	4.0bc	3.1b	2.0a
	OSM	4.5d	3.0c	2.4b	1.6a
	TL	2.6ab	3.0b	2.3ab	1.7a
			Time <sup>e</sup> in Active Tech MAP, days at 35° F		
Off-odor score <sup>f</sup> at end of display	GB	1.5a	1.9a	2.8b	2.4ab
	LD	1.3a	1.3a	2.3b	2.3b
	SM	1.5a	2.2a	3.0b	3.0b
	TL	1.6a	1.2a	3.1b	3.3b
			Time <sup>e</sup> in Active Tech MAP, days at 43° F		
Off-odor score <sup>f</sup> at end of display	GB	1.5a	3.3a	3.6a	3.9a
	LD	1.3a	2.9a	3.3ab	3.6b
	SM	1.5a	2.2a	3.4b	4.0b
	TL	1.6a	2.7a	3.3b	3.8c

a-d Means in the same row with a different letter differ (P<0.05).

<sup>e</sup> Ground beef stored 7, 14, and 21 days, other muscles 7, 21, and 35 days.

<sup>f</sup> Off-odor scale: 1 = none, 2 = slight, 3 = Small, 4 = Moderate, 5 = Extreme.

Table 4 - Means for aerobic plate counts (APC) on beef cuts exposed to carbon monoxide during storage at 35° and 43°F in Active Tech MAP vs. baseline cuts exposed only to oxygen.

Trait	Product	Baseline cuts	Time <sup>e</sup> in Active Tech MAP, days at 35° F		
			7	14 / 21	21 / 35
End of MAP storage APCs, log 10 cfu	GB	2.7a	2.6a	4.7b	5.5b
	LD	.7ab	0.2a	1.4bc	1.7c
	SM	1.0b	0.3a	0.3a	0.3a
	TL	1.3b	0.2a	2.6bc	3.1c
End of display APCs, log 10 cfu	GB	4.3a	4.4ab	5.6b	5.5b
	LD	1.4ab	0.4a	2.9bc	3.4c
	SM	0.6a	0.1a	0.6a	2.0b
	TL	0.3a	1.3b	3.5c	3.4c
			Time <sup>e</sup> in Active Tech MAP, days at 43° F		
End of MAP storage APCs, log 10 cfu	GB	2.7a	4.6b	5.8c	6.0c
	LD	0.7a	1.3ab	3.2c	5.1d
	SM	1.0b	0.1a	0.1a	2.8c
	TL	1.3a	1.6a	3.7b	4.0b
End of display APCs, log 10 cfu	GB	4.3a	5.8b	5.9b	6.1b
	LD	1.4a	1.3a	2.8b	5.3c
	SM	0.6a	0.3a	0.7a	2.5b
	TL	0.3a	3.3b	4.2b	4.6b

a-d Means in the same row with a different letter differ (P<0.05).

<sup>e</sup> Ground beef stored 7, 14, and 21 days, other muscles 7, 21, and 35 days.

**Figure 6**  
**Ground Beef  $a^*$  Values (Redness) Deterioration**  
**During Display Following Storage**






Figure 7  
Strip Loin  $a^*$  Values (Redness) Deterioration  
During Display Following Storage






Figure 8  
 Inside Round (inside portion)  $a^*$  Values (Redness)  
 Deterioration During Display Following Storage



**Figure 9**  
**Inside Round (outside portion)  $a^*$  Values (Redness)**  
**Deterioration During Display Following Storage**



**Figure 10**  
**Tenderloin  $a^*$  Values (Redness) Deterioration**  
**During Display Following Storage**






Figure 11  
Ground Beef Total Aerobic Plate Counts  
During Display Following Storage



**Figure 12**  
**Strip Loin Total Aerobic Plate Counts**  
**During Display Following Storage**



**Figure 13**  
**Inside Round Total Aerobic Plate Counts**  
**During Display Following Storage**





Figure 14  
Tenderloin Total Aerobic Plate Counts  
During Display Following Storage





**Figure 15**  
**Ground Beef Lactic Acid Bacteria**  
**During Display Following Storage**





Figure 16  
Strip Loin Lactic Acid Bacteria  
During Display Following Storage



**Figure 17**  
**Inside Round Lactic Acid Bacteria**  
**During Display Following Storage**




Figure 18  
Tenderloin Lactic Acid Bacteria  
During Display Following Storage




Figure 19  
Aerobic Plate Count  $\text{Log}_{10}$  CFU vs Visual Color




Figure 20  
Lactic Acid Bacteria Count Log<sub>10</sub> CFU vs Visual Color





Figure 21  
Odor vs. Visual Color



**ATTACHMENT 5**





WELDING SUPPLY, INC.

SPECIALTY GASES DIVISION

## Certificate of Conformance

6000 COURT STREET ROAD  
SYRACUSE, NEW YORK 13206  
(315) 463-5241

6670 MARTIN STREET  
ROME, NEW YORK 13440  
(315) 329-3378

214 NORTH FOURTH STREET  
FULTON, NEW YORK 13069  
(315) 552-5012

23771 STATE ROUTE 12  
WATERTOWN, NEW YORK 13601  
(315) 785-5438

10149 STATE HIGHWAY 56  
MASSENA, NEW YORK 13662  
(315) 764-6728

311 CLARK STREET  
AUBURN, NEW YORK 13021  
(315) 252-5398

1 LEE STREET  
UTICA, NEW YORK 13502  
(315) 724-3382

417 COMMERCE ROAD  
VESTAL, NEW YORK 13850  
(607) 757-6348

1448 MILITARY TURNPIKE  
PLATTSBURGH, NEW YORK 12601  
(516) 553-1240

5 GREGORY DRIVE  
SOUTH BURLINGTON, VT 05403  
(802) 852-4572

48 DORAN AVENUE  
GENEVA, NEW YORK 14456  
(315) 781-6880

Customer: Pactiv Packaging

Material Submitted: Carbon Monoxide, Research Purity 99.99% min.

Date Reported: 05/08/01

Component	Specification
Carbon Monoxide	99.99% min.
Oxygen	< 0.5 PPM
Nitrogen	< 10 PPM
Carbon Dioxide	< 20 PPM
Methane	< 5 PPM
Ethane	< 1 PPM
Propane	< 1 PPM
Dimethyl Ether	< 1 PPM
Hydrogen	< 1 PPM
Moisture	< 1 PPM

Note: Analysis are conducted utilizing approved analytical method (s) and are correct to within the analytical accuracies of this (these) method (s).

Quality Control Approved

*Lonnie Nash*

05-08-01

CONSISTENTLY FULFILLING CUSTOMER'S EXPECTATIONS THROUGH INNOVATIVE SOLUTIONS AND TEAMWORK

000191

**ATTACHMENT 6**



## Standard Practice for Analysis of Reformed Gas by Gas Chromatography<sup>1</sup>

This standard is issued under the fixed designation D 1946; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon ( $\epsilon$ ) indicates an editorial change since the last revision or reapproval.

### 1. Scope

1.1 This practice covers the determination of the chemical composition of reformed gases and similar gaseous mixtures containing the following components: hydrogen, oxygen, nitrogen, carbon monoxide, carbon dioxide, methane, ethane, and ethylene.

1.2 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.*

### 2. Referenced Documents

#### 2.1 ASTM Standards:

E 260 Practice for Packed Column Gas Chromatography<sup>2</sup>

### 3. Summary of Practice

3.1 Components in a sample of reformed gas are physically separated by gas chromatography and compared to corresponding components of a reference standard separated under identical operating conditions, using a reference standard mixture of known composition. The composition of the reformed gas is calculated by comparison of either the peak height or area response of each component with the corresponding value of that component in the reference standard.

### 4. Significance and Use

4.1 The information about the chemical composition can be used to calculate physical properties of the gas, such as heating (calorific) value and relative density. Combustion characteristics, products of combustion, toxicity, and interchangeability with other fuel gases may also be inferred from the chemical composition.

### 5. Apparatus

5.1 **Detector**—The detector shall be a thermal conductivity type or its equivalent in stability and sensitivity. The thermal conductivity detector must be sufficiently sensitive to produce

a signal of at least 0.5 mV for 1 mol % methane in a 0.5-mL sample.

5.2 **Recording Instruments**—Either strip chart recorders or electronic integrators, or both, are used to display the separated components. Although a strip chart recorder is not required when using electronic integration, it is highly desirable for evaluation of instrument performance.

5.2.1 The recorder, when used, shall be a strip chart recorder with a full-range scale of 5 mV or less (1 mV preferred). The width of the chart shall be not less than 150 mm. A maximum pen response time of 2 s (1 s preferred) and a minimum chart speed of 10 mm/min shall be required. Faster speeds up to 100 mm/min are desirable if the chromatogram is to be interpreted using manual methods to obtain areas.

5.2.2 **Electronic or Computing Integrators**—Proof of separation and response equivalent to that for the recorder is required for displays other than by chart recorder.

5.3 **Attenuator**—If manual methods are used to interpret the chromatogram, an attenuator must be used with the detector output signal to keep the peak maxima within the range of the recorder chart. The attenuator must be accurate to within 0.5 % between the attenuator range steps.

#### 5.4 Sample Inlet System:

5.4.1 The sample inlet system must be constructed of materials that are inert and nonadsorptive with respect to the components in the sample. The preferred material of construction is stainless steel. Copper and copper-bearing alloys are unacceptable.

5.4.2 Provision must be made to introduce into the carrier gas ahead of the analyzing column a gas-phase sample that has been entrapped in either a fixed volume loop or tubular section. The injected volume must be reproducible such that successive runs of the same sample agree within the limits of repeatability for the concentration range as specified in 11.1.1.

5.4.3 If the instrument is calibrated with pure components, the inlet system shall be equipped to introduce a sample at less than atmospheric pressure. The pressure-sensing device must be accurate to 0.1 kPa (1 mm Hg).

#### 5.5 Column Temperature Control:

5.5.1 **Isothermal**—When isothermal operation is used, the analytical columns shall be maintained at a temperature constant to 0.3°C during the course of the sample run and the corresponding reference run.

<sup>1</sup> This practice is under the jurisdiction of ASTM Committee D-3 on Gaseous Fuels and is the direct responsibility of Subcommittee D03.07 on Analysis of Chemical Composition of Gaseous Fuels.

Current edition approved March 30, 1990. Published May 1990. Originally published as 1946 – 62 T. Last previous edition D 1946 – 82.

<sup>2</sup> *Annual Book of ASTM Standards*, Vol 14.02.



**5.5.2 Temperature Programming**—Temperature programming may be used, as feasible. The oven temperature shall not exceed the recommended temperature limit for the materials in the column.

**5.6 Detector Temperature Control**—The detector temperature shall be maintained at a temperature constant to  $0.3^{\circ}\text{C}$  during the course of the sample run and the corresponding reference run. The detector temperature shall be equal to, or greater than, the maximum column temperature.

**5.7 Carrier Gas**—The instrument shall be equipped with suitable facilities to provide flow of carrier gas through the analyzer and detector at a flow rate that is constant to 1 % throughout the analysis of the sample and the reference standard. The purity of the carrier gas may be improved by flowing the carrier gas through selective filters before its entry into the chromatograph.

#### 5.8 Columns:

**5.8.1** The columns shall be constructed of materials that are inert and nonadsorptive with respect to the components in the sample. The preferred material of construction is stainless steel. Copper and copper-bearing alloys are unacceptable.

**5.8.2** Either an adsorption-type column or a partition-type column, or both, may be used to make the analysis.

NOTE 1—See Practice E 260 for general gas chromatography procedures.

**5.8.2.1 Adsorption Column**—This column must completely separate hydrogen, oxygen, nitrogen, methane, and carbon monoxide. If a recorder is used, the recorder pen must return to the baseline between each successive peak. Equivalent proof of separation is required for displays other than by chart recorder. Fig. 1 is an example chromatogram obtained with an adsorp-

tion column.

(1) Because of similarities in thermal conductivities, helium should not be used as the carrier gas for hydrogen when hydrogen is less than 1 % of the sample. Either argon or nitrogen carrier gas is suitable for both percent and parts per million quantities of hydrogen.

(2) The use of a carrier gas mixture of 8.5 % hydrogen and 91.5 % helium will avoid the problem of reversing polarities of hydrogen responses as the concentration of hydrogen in the sample is increased.

(3) The precision of measurement of hydrogen can be increased by using a separate injection for hydrogen, using either argon or nitrogen for the carrier gas.


(4) Another technique for isolating the hydrogen in a sample is to use a palladium transfer tube at the end of the adsorption column; this will permit only hydrogen to be transferred to a stream of argon or nitrogen carrier gas for analysis in a second thermal conductivity detector.

**5.8.2.2 Partition Column**—This column must separate ethane, carbon dioxide, and ethylene. If a recorder is used, the recorder pen must return to the baseline between each successive peak. Equivalent proof of separation is required for displays other than by chart recorder. Fig. 2 is an example chromatogram obtained with a partition column.

**5.8.3 General**—Those column materials, operated either isothermally or with temperature programming, or both, may be used if they provide satisfactory separation of components.

#### 6. Reference Standards


**6.1** Moisture-free mixtures of known composition are required for comparison with the test sample. They must contain known percentages of the components, except oxygen (Note



Column: 2-m by 6-mm inside diameter Type 13 $\times$  molecular sieves, 14 to 30 mesh  
Temperature:  $35^{\circ}\text{C}$

Flow rate: 60-mL helium/min  
Sample size: 0.5 mL

FIG. 1 Chromatogram of Reformed Gas on Molecular Sieve Column



Column: 1.2 m by 6.35 mm  
 Porapak Q, 50 to 80 mesh  
 Current setting: 225 mA

Temperature: 40°C  
 Flow rate: 50-mL helium/min  
 Sample size: 0.5 mL

FIG. 2 Chromatogram of Reformed Gas on Porapak Q Column

2), that are to be determined in the unknown sample. All components in the reference standard must be homogeneous in the vapor state at the time of use. The fraction of a component in the reference standard should not be less than one half of, nor differ by more than 10 mol % from, the fraction of the corresponding component in the unknown. The composition of the reference standard must be known to within 0.01 mol % for any component.

NOTE 2—Unless the reference standard is stored in a container that has been tested and proved for inertness to oxygen, it is preferable to calibrate for oxygen by an alternative method.

6.2 Preparation—A reference standard may be prepared by blending pure components. Diluted dry air is a suitable standard for oxygen and nitrogen.

NOTE 3—A mixture containing approximately 1 % of oxygen can be prepared by pressurizing a container of dry air at atmospheric pressure to 20 atm (2.03 MPa) with pure helium. This pressure need not be measured

precisely, as the fraction of nitrogen in the mixture such prepared must be determined by comparison to nitrogen in the reference standard. The fraction of nitrogen is multiplied by 0.280 to obtain the fraction of oxygen plus argon. Argon elutes with oxygen in the molecular sieves column. Do not rely on oxygen standards that have been prepared for more than a few days. It is permissible to use a response factor for oxygen that is relative to a stable component.

## 7. Preparation of Apparatus

7.1 Column Preparation—Pack a 2- to 3-m column (6-mm inside diameter stainless steel tubing) with Type 13× molecular sieves, 14 to 30 mesh, that have been dried 12 h or more at 300 to 350°C. Pack a second column (1 m by 6 mm) with Porapak Q,<sup>3</sup> 50 to 80 mesh, that has been dried 12 h or more at about 150°C. Shape the columns to fit the configuration of the oven in the chromatograph.

<sup>3</sup> Available from Waters Associates, Inc., Framingham, MA 01701.

NOTE 4—Variations in column material, dimensions, and mesh sizes of packing are permissible if the columns produce separations equivalent to those shown in Fig. 1 and Fig. 2. Better performance may be obtained by using a 2.1-mm stainless steel tubing with corresponding smaller mesh packing materials and substituting Haysep Q for Porapak Q.

7.2 *Chromatograph*—Place the proper column and sample volume in operation for the desired run in accordance with 8.1 and 8.2. For isothermal operation, the column should be maintained at a temperature between 30 and 45°C. When appropriate, column temperatures may be increased. Adjust the operating conditions and allow the instrument to stabilize. Check the stability by making repeat runs on the reference standard to obtain reproducible peak heights as described in 5.4.2 for corresponding components.

## 8. Procedure

8.1 *Sample Volume*—The sample introduced into the chromatographic column should have a volume between 0.2 and 0.5 mL. Sufficient accuracy can be obtained for the determination of all but the very minor components with this sample size. When increased sensitivity is required for the determination of components present in low concentrations, a sample size of up to 5 mL is permissible. However, components whose concentrations are in excess of 5 % should not be analyzed by using sample volumes greater than 0.5 mL.

### 8.2 Chromatograms:

8.2.1 *Adsorption Column* (Fig. 1)—Obtain a steady baseline on the recorder with a constant carrier gas flowrate appropriate to the column diameter. Introduce a sample of the unknown mixture at atmospheric pressure into the chromatograph and obtain a response similar to that of Fig. 1 of the components hydrogen, oxygen, nitrogen, methane, and carbon monoxide, which elute in that order. Repeat with a sample of the reference standard. If oxygen is present in the mixture, run a sample of air, either at an accurately measured reduced pressure, or air freshly diluted with helium, so that the partial pressure of oxygen is approximately equal to that of the oxygen in the mixture being analyzed.

NOTE 5—The peak for carbon monoxide can appear between those of nitrogen and methane if the molecular sieves have become contaminated. If this occurs, replace or regenerate the column packing by heating in accordance with 7.1.

8.2.2 *Partition Column* (Fig. 2)—Establish a steady baseline with the helium carrier gas flowing through the Porapak Q column. Introduce a sample of the reference standard and then a sample of the unknown mixture. Obtain responses similar to that shown in Fig. 2 for carbon dioxide, ethane, and ethylene.

8.2.3 All chromatograms for manual measurement should be run at a sensitivity setting that permits maximum peak height to be recorded for each component.

8.2.4 Column isolation valves may be used to make the entire analysis with a single injection if the separations specified in 5.8.2.1 and 5.8.2.2 are produced.

## 9. Calculation

9.1 The number of significant digits retained for the quantitative value of each component shall be such that accuracy is neither sacrificed nor exaggerated. The expressed numerical

value of any component in the sample should not be presumed to be more accurate than the corresponding certified value of that component in the calibration standard.

9.2 *Manual Measurement*—Measure the response of each component, convert to the same sensitivity for corresponding components in the sample and reference standard, and calculate the mole percent of each component in the sample as follows:

$$C = (A/B)(S) \quad (1)$$

where:

$C$  = mole percent of the component in the sample,

$A$  = response of the component in the sample,

$B$  = response of the component in the standard at the same sensitivity as with  $A$ , and

$S$  = mole percent of the component in the reference standard.

9.3 If a helium-diluted air mixture was run for oxygen calibration, calculate the fraction of oxygen in the mixture from the fraction of the nitrogen and the composition of the diluted air. Calculate the fraction of nitrogen in the mixture in accordance with 9.1, using the nitrogen response of the reference standard for comparison. Air composition values of 78.1 % nitrogen and 21.9 % oxygen should be used, as argon (0.9 % in air) elutes with oxygen on the molecular sieves column.

9.4 If air has been analyzed at reduced pressure to calibrate for oxygen, correct the equation for pressure as follows:

$$C = (A/B)(S)(P_a/P_b) \quad (2)$$

where:

$P_a$  = absolute pressure at which air was analyzed and

$P_b$  = barometric pressure when sample was analyzed, with both pressures being expressed in the same units.

9.5 Normalize the mole percent values by multiplying each value by 100 and dividing by the sum of the original values. The sum of the original values should not differ from 100.0 % by more than 1.0 %.

## 10. Analysis of the Reference Standard

10.1 If the composition of the reference standard is not known to a sufficient degree of accuracy, analyze it by the use of pure components for calibration. Obtain chromatograms of the standard as described in 8.2, except measure the pressure of each sample introduced to 0.133 kPa (1 mm Hg). When each chromatogram is obtained, calibrate each component by introducing a sample of the pure component at a pressure that closely approximates its partial pressure in the blend (for example, a component whose concentration in the standard is 50 % is analyzed at 50 % of the pressure at which the standard was analyzed). Use a minimum pressure of 0.665 kPa (5 mm Hg) for minor components. Repeat the analysis with the reference standard. Corresponding peak heights should agree within 1 mm or 1 % (whichever is larger) when recorded on a sensitivity setting that allows maximum response on the recorder chart.

10.2 Calculate the composition of the reference standard by the adjustment of responses of like components to the same sensitivity and calculate the concentration of each component as follows:

$$C = \frac{(100)(R)(P_p)}{(P)(P_r)} \quad (3)$$

where:

- $C$  = component concentration, mole percent;
- $R$  = response of the component in the reference standard;
- $P$  = response of the pure component;
- $P_p$  = pressure at which the pure component was analyzed; and
- $P_r$  = pressure at which the reference standard was analyzed, with both pressures being expressed in the same absolute units.

10.2.1 Normalize all values as described in 9.4.

## 11. Precision

11.1 The following data should be used to judge the acceptability of the results:

11.1.1 *Repeatability*—Duplicate results by the same operator should not be considered suspect unless they differ by more

than the following amounts:

Component, mol %	Repeatability
0 to 1	0.05
1 to 5	0.1
5 to 25	0.3
Over 25	0.5

11.1.2 *Reproducibility*—Results submitted by different laboratories should not differ by more than the amounts given in 11.1.1 when the same reference standard is used for calibration and the same composition is used for calculations. If calibration is made with pure components or with different reference standards, results submitted by each of two laboratories should not be considered suspect unless the results differ by more than the following amounts:

Component, mol %	Reproducibility
0 to 1	0.1
1 to 5	0.2
5 to 25	0.5
Over 25	1.0

## 12. Keywords

12.1 gaseous fuels

*The American Society for Testing and Materials takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.*

*This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.*

*This standard is copyrighted by ASTM, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-632-9585 (phone), 610-632-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org).*

ATTACHMENT 7





DEPARTMENT OF NUTRITION AND FOOD SCIENCES  
College of Agriculture  
College of Family Life  
Logan, UT 84322-8700  
Telephone: (435) 797-2126  
FAX: (435) 797-2379

Division of GRAS Notice Review  
Office of Food Additive Safety  
Center for Food Safety and Applied Nutrition  
Food and Drug Administration  
200 C St, SW  
Washington, DC 20204

August 17, 2001

Dear FDA Personnel,

I am a meat scientist with experience in studying the effects on fresh meats of various modified atmospheres. Based on my review of the details of the ActiveTech 2001 modified atmosphere system employing 0.4% carbon monoxide gas in a mixture with 60 percent carbon dioxide and the remainder nitrogen, as well as the published literature and common knowledge in the field, I confirm that the use of modified atmospheres including 0.4% CO to package fresh meats as used in the ActiveTech 2001 system is both safe and generally recognized as safe. If I can provide any further information or clarification, please contact me.

Sincerely yours,

A handwritten signature in cursive script that reads "Daren Cornforth".

Daren Cornforth, Ph.D.  
Professor, Nutrition & Food Sciences  
Utah State University  
435-797-2114  
darenc@cc.usu.edu

000199

**Daren Cornforth, Ph.D.**

Birthdate: May 23, 1949

Birthplace: Fort Collins, Colorado

Current Address: Department of Nutrition and Food Sciences  
Utah State University  
Logan, Utah 84322-8700  
(435) 797-2114 fax (435) 797-2379  
e-mail darenc@cc.usu.edu

**Education**

<u>Institution</u>	<u>Degree</u>	<u>Year Conferred</u>	<u>Scientific Field</u>
Colorado State University	B.S.	1971	Animal Science
Colorado State University	M.S.	1973	Animal Science
Michigan State University	Ph.D.	1978	Food Science & Human Nutrition

**Research and Professional Experience**

<u>Dates</u>	<u>Position</u>	<u>Duties</u>
1978 to present	Asst., Assoc., Full Prof., NFS	Teaching/Research, Meat Processing
1974 to 1978	Graduate Research Assistant at Michigan State University East Lansing, Michigan	Dissertation on cold- induced shortening and toughening of beef muscle.
1971 to 1973	Graduate Research Assistant at Colorado State University Fort Collins, Colorado	Thesis on the effects of breed, sex, and diet on muscle fiber type and fat cell development in growing calves.

**Memberships**

Institute of Food Technologists, Amer. Assoc. for the Advancement of Science, Sigma Xi (Past President, USU Chapter), American Meat Science Association, Farm House Fraternity, Rocky Mountain Elk Foundation (Life Member), Pheasants Forever, Nature Conservancy, Cache Valley Wildlife Federation (and on Executive Board).

## Current Teaching Assignment

NFS 5560      Chemistry of Food Systems  
NFS 6450      Meat Science  
NFS 1000      World of Food and Nutrition

I also assist with the NFS Food Science Club College Bowl team, and coordinate (with Dr. Carpenter) the Utah Future Farmers of America (FFA) state contest in Food Science and Technology.

## Invited Speaker (1997-2001)

Cornforth, D. P. 1997. *Pigment concentration and pH effects on degree of doneness of beef patties*. FMC Corporation. Atlanta, Ga, January, 1997.

Cornforth, D. P. 1997. *Nitrogen dioxide causes surface pinking on meats cooked in gas ovens*. Meat Industry Meat Conf. Chicago, IL, Oct 28-29.

Cornforth, D. P., Jiang, C. and Mumford, B. 1998. *Effect of storage time, storage and cooking temperature, and holding time after cooking on pigment levels in beef patties*. Symposium, New Developments in Meat Processing, Amer. Meat Sci. Assn. National Meeting, Storrs, CT., June 28-July 2.

Cornforth, D. P. 2000. Keynote speaker. *Pinking in Cooked Poultry. Current situation and general theories*. Given at the Symposium on Pinking in Cooked Poultry, Center for Excellence in Poultry Science, University of Arkansas, Fayetteville, Oct 12-13, 1999. Co-sponsored by Tyson Foods, NewlyWed Foods, and KFC, Inc.

Cornforth, D. P. 2000. *Pinking in Cooked Poultry*. Michigan State University, E. Lansing, MI, Nov. 7.

Cornforth, D. P. 2000. *Pinking in Cooked Poultry*. Michigan Turkey Processors, Zeeland, MI, Nov. 8.

Cornforth, D. P. 2001. *Meat Color*. Dinner speaker for the joint meeting of the Southern Minnesota section of IFT and the Meat Processing Short Course sponsored by the Agricultural Utilization Research Institute, Southwest State University, Marshall, MN, April 25.

## **National Offices and Committees (1997-2001)**

**USDA** Advisory Committee for the Safe Preparation of Ground Beef - 1997-8.

### **AMSA (American Meat Science Association)**

Graduate Student Poster Competition committee, Reciprocal Meat Conference, 1998-2002.

Executive Board 2000-2002.

**IFT** Executive Committee, Muscle Foods Division, Institute of Food Technologists, 1998-99.

Chair-Elect, Muscle Foods Division, 2000-2001.

Chair, Muscle Foods Division, 2001-2002.

Executive Council, Bonneville Section, IFT (Councilor representing Bonneville Section at National Meetings), 1998-99.

**Journal of Muscle Foods** – Editorial Board, 1998-2002

## **University Committee Assignments**

Committee for Laboratory Safety. 1997-2001.

Faculty Senate, 2000 – 2003.

Faculty Senate Executive Committee 2000 – 2001.

## **Grants (1997-2001)**

Cornforth, D. P. 1997-98. Factors affecting hamburger degree of doneness. McDonald's Corp. Oak Brook, IL. \$8,480.

Cornforth, D. P. 1998. Evaluation of various dairy fractions as inhibitors of lipid oxidation in precooked meats. Western Dairy Foods Center, Logan, UT, \$9,560.

Cornforth, D. P. 1999. Verification of nutrient label claims on lowfat milk, lean hamburger, and breakfast cereals. Dr. Alan Luke (Alumni Donor), \$1,500.

Cornforth, D. P. and Carpenter, C. E. 1999-2000. Evaluation of carbon monoxide treatment in modified atmosphere or vacuum packaging to increase color stability of fresh beef. National Cattlemen's Beef Association, \$14,500.

Carpenter, C.E. and Cornforth, D. P. 1999-2000. Effect of consumer bias for beef color and packaging on eating satisfaction. National Cattlemen's Beef Association, \$15,000.

Cornforth, D. P. 1999-2002. Evaluation of antioxidant properties of milk powders in cooked meats. Glanbia Foods, Twin Falls, ID. \$30,000.

Cornforth, D. P. 2000-2002. New antioxidants in cooked meats. Utah Agric. Exp. Stat. \$15,000.

Bailey, D. and Dickinson, D. (Cornforth, D.P., co-investigator). 2001-2003. Traceability: A market opportunity or threat to the US meat industry. USDA-CREES. \$160,000.

Cornforth, D. P. 2001-2002. Dried whey minerals as an antioxidant in processed meats. Dairy Management Institute (DMI). \$97,500.

#### Graduate Committees Chaired (1997-2001)

Heaton, K. M. 1998. Minimum levels of nitrite, nitric oxide, and carbon monoxide causing pinking in cooked beef and turkey rolls. M. S. thesis, Utah State University, Logan, Utah. (Now Extension Agent, Garfield & Kane Counties, Utah).

Moiseev, I. V. 1998. Prevention of pink discoloration and microbial safety of rolls and patties made from dark-cutting beef, PhD dissertation, Utah State University, Logan, Utah. (Now lab chemist, Borden Foods, Columbus, OH).

Racz, J. M. 1998. An inoculated pack study using *Clostridium sporogenes* PA 3679 to determine the shelf stability of a vacuum packaged meat/vegetable stick. M. S. thesis, Utah State University, Logan, Utah. (Now lab chemist, Fresenius Medical, Ogden, UT).

Jayasingh, P. 2001. Dried milk minerals as an antioxidant in processed meats. Ph.D. dissertation, Utah State University, Logan, Utah (work in progress).

#### Product Development and Extension

Stew Sticks are currently in retail production by Utah Jerky, Inc., Ogden, UT.

Cornforth, D. P., Bailey, D., and McEvoy, R. 1997. Developed a video of ostrich slaughter and processing, used by the new plant in Filmore, UT for employee training, and also used by USU extension agents.

### Refereed Publications (1997-2001)

- Moiseev, I. V. and Cornforth, D. P. 1997. Sodium hydroxide and sodium tripolyphosphate effects on bind strength and sensory characteristics of restructured beef rolls. *Meat Sci.* 45:53-60.
- Quinton, R. D., Cornforth, D. P., Hendricks, D. G., Brennand, C. P. and Su, Y. K. 1997. Acceptability and composition of some acidified meat and vegetable stick products. *J. Food Sci.* 62:1250-1254.
- Cornforth, D. P., Rabovitser, J. K., Ahuja, S., Wagner, J. C., Hanson, R., Cummings, B. and Chudnovsky, Y. 1998. Carbon monoxide, nitric oxide, and nitrogen dioxide levels in gas ovens related to surface pinking of cooked beef and turkey. *J. Agric. Food Chem.* 46:255-61.
- Lee, B., Hendricks, D. G. and Cornforth, D. P. 1998. Antioxidant effects of carnosine and phytic acid in a model beef system. *J. Food Sci.* 63:394-398.
- Lee, B., Hendricks, D. G. and Cornforth, D. P. 1998. Effect of sodium phytate, sodium pyrophosphate, and sodium tripolyphosphate on physicochemical characteristics of restructured beef. *Meat Sci.* 50:273-283.
- Moiseev, I. V. and Cornforth, D. P. 1999. Treatments for prevention of persistent pinking in dark-cutting beef patties. *J. Food Sci.* 64:738-43.
- Lee, B., Hendricks, D. G. and Cornforth, D. P. 1999. A comparison of carnosine and ascorbic acid on color and lipid stability in a ground beef patty model system. *Meat Sci.* 51:245-253.
- Heaton, K. M., Cornforth, D. P., Moiseev, I. V., Egbert, W. R. and Carpenter, C. E. 2000. Minimum sodium nitrite levels for pinking of various cooked meats as related to use of direct or indirect-dried soy isolates in poultry rolls. *Meat Sci.* 55:321-329.
- Carpenter, C. E., Cornforth, D. P., Whittier, D. 2001. Consumer preferences for beef color and packaging did not affect eating satisfaction. *Meat Science* 57:359-363.
- Jayasingh, P., Cornforth, D. P., Carpenter, C. E., and Whittier, D. 2001. Evaluation of carbon monoxide treatment in modified atmosphere packaging or vacuum packaging to increase color stability of fresh beef. *Meat Science*, In press.

### Book Chapters

- Cornforth, D. P. 2000. Miscellaneous Colorants - Cured Meat. Unit F6.2 in *Current Protocols in Food Analytical Chemistry*. S. J. Schwartz, Ed. John Wiley & Sons, Inc. New York, NY.
- Cornforth, D. P. 2001. Potential use of phytate as an antioxidant in cooked meats. Ch. 11 in *Food Phytates*. R. Reddy and S. K. Sathe, Eds. Technomic Publ., Inc., Rowayton, CT.

ATTACHMENT 8

# ENVIRON

August 27, 2001

Eric Greenberg  
Ungaretti & Harris  
3500 Three First National Plaza  
Chicago, IL 60602-4283

Re: Generally recognized as safe ("GRAS") determination for carbon monoxide from Pactiv ActiveTech food-contact packaging.

Dear Eric:

This letter reports my assessment, based on my review of the available information, that the potential consumer exposure to carbon monoxide from the Pactiv ActiveTech food-contact packaging is safe, and is also generally recognized as safe ("GRAS"), under the Federal Food, Drug, and Cosmetic Act ("FDCA" or "the Act") when used under conditions defined in the attached GRAS determination document. First, the safety of carbon monoxide in this use is demonstrated by comparison of the U.S. Environmental Protection Agency ("USEPA") national ambient air quality standard ("NAAQS") for carbon monoxide with the estimated daily intake ("EDI") of carbon monoxide under the intended conditions of use of the product. Exposure to a substance generally is considered safe for its intended use if the EDI is a fraction of the allowable daily intake ("ADI"). Second, the GRAS status of carbon monoxide in this use is affirmed by demonstrating that the safety of carbon monoxide from Pactiv ActiveTech food-contact packaging material in its intended use is generally recognized.

My GRAS determination was based on scientific procedures as outlined in the U.S. Food and Drug Administration ("FDA") regulations (at §170.30(b)). This section requires that the same quantity and quality of scientific evidence is available and is reviewed as is required to obtain approval of the substance as a food additive. Moreover, in addition to requiring scientific evidence of safety (as with a food additive), a GRAS determination also requires that this scientific evidence of safety be generally known and accepted by experts qualified in the appropriate scientific and technical fields. This common knowledge requirement of a GRAS determination includes two elements: (1) the data and information relied upon to establish the scientific basis for safety must be generally available; and (2) there must be a basis to conclude that there is a consensus among qualified experts about the safety of the substance for its intended use.

Based on the scientific literature, studies, conclusions, and restrictions presented in the GRAS determination document, I regard the proposed uses of carbon monoxide from Pactiv ActiveTech packaging material as generally recognized as safe because the use will result in an exposure that is well below an acceptable exposure level for carbon monoxide.

---

ENVIRON International Corporation

4350 North Fairfax Drive, Suite 300 • Arlington, VA 22203 USA  
Tel: (+1) (703) 516 2300 • Fax: (+1) (703) 516 2393 • [www.environcorp.com](http://www.environcorp.com)

000006



No evidence exists in the available information on carbon monoxide that demonstrates, or suggests reasonable grounds to suspect, a hazard to the public health when carbon monoxide is used at levels that might reasonably be expected from its proposed use as an additive in the Pactiv ActiveTech food-contact packaging material as defined in the GRAS determination document.

It is my opinion that other qualified and competent scientists reviewing the same publicly available data would reach the same scientific conclusion. Therefore, carbon monoxide, as an additive in the Pactiv ActiveTech food-contact packaging material, is generally recognized as safe.



---

Vasilios H. Frankos, Ph.D.  
Principal, Health Sciences  
ENVIRON International Corporation  
Arlington, Virginia

## VASILIOS H. FRANKOS, Ph.D.

### EDUCATION

- 1977 Ph.D., Pharmacology and Toxicology, University of Maryland Pharmacy School
- 1973 M.S., Biology, University of Maryland
- 1970 B.A., Biology, University of Maryland

### EXPERIENCE

Dr. Frankos is a Principal at ENVIRON Corporation and has over 20 years of experience in the toxicological and pharmacological evaluation of data used to assess the risks posed by foods and food additives, drugs, medical devices, cosmetics, pesticides, and environmental and occupational exposures. He has also been involved in the development of exposure and risk assessment methodology. Since joining ENVIRON, Dr. Frankos has led, contributed to, or managed hundreds of projects in these areas.

#### **Foods and Food Additives:**

Dr. Frankos has worked on a wide variety of projects evaluating the safety of foods, direct and indirect food additives, and food contaminants. As part of these food-related safety evaluations, he has developed strategies for testing new direct and indirect additives, evaluated toxicity test data to support safety determinations, prepared Generally Recognized as Safe (GRAS) reviews, performed exposure and risk assessments, and developed regulatory strategies. Dr. Frankos has also presented safety evaluations to the FDA on behalf of clients. Some of his major projects in the area of foods and food additives include:

- Provided ongoing FDA-related scientific and technical support to the Coalition for Safe Ceramicware (CSC). Performed a safety assessment of ceramic pitchers with glazes containing lead using data collected on migration of lead into a food simulant and into real foods. This assessment was submitted to the FDA as part of the CSC's comments on the FDA's proposed rule changing the action level for lead from ceramic pitchers.
- Developed direct food additive petitions and GRAS self-affirmation documents for numerous food additives including, novel fibers sources, an anti-caking agent, enzymes, sugars, and a major new class of food additives.
- Conducted a GRAS self-affirmation review, including an evaluation of safety data, of the first bioengineered food approved by the FDA. Presented this GRAS review to the FDA on behalf of the client, a major biotechnology company.
- Developed a GRAS affirmation document for a cellulose product, manufactured by a novel bacterial fermentation process, with proposed food use as a suspending/thickening agent. Designed, placed, and monitored preclinical toxicity studies required for FDA approval.
- Estimated doses posing no significant risk for chemicals that could potentially leach from packaging into food. Assessed the potential human exposure to these chemicals from migration from packaging into food. Compared the potential ingested dose to the no significant risk dose.

## VASILIOS H. FRANKOS, Ph.D.

- Petitioned the FDA to sanction expanded use in foods of an approved food additive. Prepared a review and update of existing toxicological literature on the material and estimated the increase in exposure likely to result from proposed new uses.
- Evaluated the carcinogenic risk associated with exposure to acrylamide residues in food and methylene chloride in decaffeinated tea.
- Addressed issues relating to FDA's regulation of polychlorinated biphenyl (PCB) residues. Examined whether tolerances for PCBs in fish could be reinterpreted for less chlorinated PCBs that have lower or no carcinogenic potency. Determined necessary research to establish differences in potency between PCBs.
- Developed an innovative exposure and safety assessment for a novel single cell protein (mycoprotein) meat substitute that has been submitted to the FDA for approval.
- Conducted a simulated FDA review of a food additive petition on a new artificial sweetener submitted to the FDA by the client's competitor. Review included critical evaluation of product chemistry, efficacy, estimates of human exposure, animal and human toxicology data, pharmacokinetics and metabolism information, and the basis for determining the acceptable daily intake of the sweetener.
- Performed a detailed evaluation of toxicity and carcinogenicity studies sponsored by a major drug company and studies from the published literature for the company's non-nutritive sweetener and assessed the toxicological significance to humans. Assisted in submission of a food additive petition. Provided continued regulatory support during the FDA review process.

### Human and Veterinary Drugs, Medical Devices, and Cosmetics:

Dr. Frankos has provided scientific and regulatory guidance to clients in the human and veterinary drug, medical device, and cosmetic industries. He has been involved in identifying and assessing the risks to humans associated with exposure to constituents of these products. He has assisted clients in these industries in interacting with the FDA and has assisted them in complying with all aspects of FDA regulations. Some of his major projects in the areas of drugs, medical devices, and cosmetics include:

- Reviewed two large epidemiological studies on the differences in adverse drug reaction rates between two types of radiographic contrast media. Prepared a safety review document on animal and human literature on contrast media.
- Performed an independent evaluation of a New Drug Application (NDA) submission to the FDA, with emphasis on review of efficacy studies.
- Assisted the medical device manufacturer in complying with FDA's post-approval requirements for its device including compliance with the Medical Device Reporting (MDR) rule, submission of updates to the PMA application, and ensuring that all labeling and marketing materials are in compliance with FDA regulations. Designed a post-marketing clinical trial for the device to comply with FDA recommendations.
- Evaluated the potential carcinogenic risks to humans of an over-the-counter (OTC) medication that is applied to the skin. Prepared a report on these findings that was submitted to the FDA.

**VASILIOS H. FRANKOS, Ph.D.**

- Prepared and submitted to the FDA a New Drug Application (NDA) for a drug that holds promise for dramatically decreasing the high percentage of reocclusion that occurs in angioplasty patients.
- Assisted a major pharmaceutical manufacturer in assessing potential health risks associated with a specific ingredient of various over-the-counter (OTC) drugs.
- Critically evaluated both published and unpublished studies on a psychoactive drug and rendered an opinion to the client on potential health effects of the drug and whether a no-observed-effect level (NOEL) had been established.
- Provided guidelines for subchronic testing to evaluate the safety for human use of an allergen desensitizer that was produced by polymerizing the allergen through a glutaraldehyde treatment.
- Analyzed the potential risk to humans resulting from the use of Furazolidone as an animal drug. Determined an estimate of this risk and presented the estimate to the FDA at a public hearing.
- Assisted a major manufacturer of veterinary drug products in developing an approach to dealing with mouse liver tumors and their usefulness as evidence of carcinogenicity.
- Reviewed and evaluated a New Animal Drug Application for FDA submission. Advised on the necessity of future studies.
- Assembled an expert panel to address the safety of an antimicrobial agent, extracted from a plant source, for use in oral hygiene products (e.g. toothpaste and oral rinse). The evaluation included a review of the preclinical and clinical toxicologic database, analysis of exposure, and determination of margin of safety associated with the proposed oral uses.
- Critically evaluated the evidence cited by FDA as the basis for considering nitrofurantoin animal drugs to be carcinogenic under the meaning of the Delaney Clause of the Federal Food, Drug, and Cosmetic Act.
- Reviewed toxicity data to be submitted in support of an Investigational Device Exemption (IDE) for an implantable medical device. Recommended and monitored the performance of supplemental tests, performed an exposure assessment of substances leaching from the device into the systemic circulation, characterized the risk to health from such exposures, and assisted in the presentation of these findings to the FDA.
- Evaluated toxicity data on the materials used in a device intended to be implanted in the abdominal cavity. Examined the adequacy of the existing data on the device material, the safety of the material used, and the safety of the proposed replacement material. Recommended studies to improve the data for submission to the FDA.
- Conducted a quantitative risk assessment on numerous color additives used in dermally applied cosmetics including an evaluation of toxicity, an analysis of exposure, and a determination of quantitative risks.
- Performed a hypothetical risk assessment for two colors used in cosmetics, based on the assumption that, if tested, they would produce tumors in rats. Demonstrated that such an outcome would still allow continued safe use of these colors in cosmetics.

## VASILIOS H. FRANKOS, Ph.D.

### Pesticides:

Dr. Frankos has assisted U.S. and foreign manufacturers in obtaining EPA and California registration of agricultural, forestry, and homeowner use pesticide products. Some of his major projects in the area of pesticides include:

Reviewed EPA's assessment of dietary oncogenic risk of two fungicides and advised the manufacturers of additional data needed to perform a quantitative risk assessment.

- Designed and supervised a field study to estimate exposure to a pesticide during "worst case" application. The study monitored the application of the chemical, measured exposure of the user during various phases of application and determined the effect of protective clothing on exposure.
- Reviewed the results of an aquatic organism field monitoring study and its supporting laboratory data for a major manufacturer of agricultural chemicals.
- Evaluated toxicity and prepared a risk assessment for residues of a fungicide in imported wines. Counseled client on process necessary to receive an EPA import tolerance for the fungicide. Advised client on additional data needed to support a tolerance.
- Designed and monitored toxicology studies for a German firm required for registration of a plant growth promotor and assisted in submitting data to the EPA.
- For a West German pesticide manufacturing company wishing to purchase the patent rights to a new pesticide developed in the U.S., provided counsel on the acceptability of the available data to EPA and OECD and the further data needed to obtain a registration in the U.S.

### Environmental and Occupational Exposures:

Dr. Frankos has directed numerous exposure and risk assessments involving hundreds of chemicals that have been associated with industrial processes, toxic waste or municipal incinerators, and hazardous waste sites. These assessments have used computerized models and include all routes of exposure. Some of his major projects in the areas of environmental and occupational exposures include:

- Performed a safety assessment for the consumption of drinking water in contact with a piece of equipment that could potentially release lead, including an estimation of 1990 baseline blood lead levels for four subpopulations using the Integrated Uptake/Biokinetic Model and a determination of the maximum acceptable concentration of lead in drinking water that was potentially in contact with the equipment. Collected and reviewed information on factors that affect the leachability of lead into drinking water.
- Assisted manufacturers of the plasticizer di-(2-ethylhexyl) phthalate (DEHP) in developing a method for estimating exposure resulting from the chemical's presence in polyvinyl chloride (PVC) consumer products such as vinyl-covered furniture, vinyl wallpaper, flooring, and shower curtains.
- Critically reviewed acrylamide's carcinogenic activity in Fischer 344 rats and designed a new cancer bioassay in rats that will improve risk assessments for acrylamide.

## **VASILIOS H. FRANKOS, Ph.D.**

- Reviewed the scientific basis for establishing safety factors needed to protect workers from reproductive toxicity associated with glycol ether exposure.
- Reevaluated the data from a National Cancer Institute (NCI) bioassay on dibutyltin diacetate to resolve discrepant interpretations by NCI and FDA.
- Reviewed FDA and National Toxicology Program (NTP) evaluations of the bioassay on dimethylterephthalate and rendered an opinion on the adequacy of the NTP and FDA decisions.
- Supported the EPA Office of Solid Waste in developing measures of inherent toxicity which can be combined with exposure estimates to provide a relative ranking for scheduling hazardous solid waste.
- Provided scientific litigation support for a municipality in which its waste treatment plant and landfill were contaminated with PCBs.
- Participated in a number of projects examining the developmental, embryotoxic, and teratogenic effects of lead observed in animals, including the effects of lead on the reproductive systems of male and female rats. Recommended a no-observed-adverse-effect level (NOAEL) for lead.

### **Exposure and Risk Assessment Methodology Development:**

Dr. Frankos has participated in the development of new exposure and risk assessment methodologies for federal and state regulatory agencies. He has also been integral in the development of exposure and risk modeling software. Some of his major projects in these areas include:

- Developed a background document for the EPA on reproductive toxicity risk assessment for use in drafting interagency risk assessment guidelines.
- Directed development of a scheme for the EPA that allows severity of toxic effects to be incorporated into safety evaluations of EPA regulated products.
- Summarized and compiled comments received by EPA on their proposed guidelines (1985) for developmental toxicity assessment.
- Assisted in evaluating EPA's procedures for estimating safe short-term exposure limits and in developing an alternative method that could be uniformly used by the entire agency.
- Assisted in the development of criteria for listing chemicals as developmental toxicants under California's Proposition 65.
- Directed the development of a computer software system, ERMA (Exposure and Risk Modeling Assistant) that enables ENVIRON to provide high quality, scientifically defensible evaluations of potential exposures and resultant risk.

Before joining ENVIRON Corporation, Dr. Frankos held the following positions:

- Associate Director, Life Sciences Division, Clement Associates. In that position he had the following experience:
  - Supervised a staff of eight scientists who assessed the risk posed by environmental contaminants, occupational carcinogens, pesticides, drugs, commercial product

VASILIOS H. FRANKOS, Ph.D.

- constituents, and food additives. Many of these projects involved evaluating human and animal toxicology data for use in conducting human risk assessments. This position required management of time constraints, budgetary limitations, and personnel allocations in a manner that provided the client with a scientifically defensible document.
- Directed preparation of reports for industry clients that included "Safety Assessment Strategies for Feminine Hygiene Products"; "The Impact of a Proposed Salicylate Warning on the Risk Associated with Diseases and Conditions in Children"; and "A Proposed Mechanism of Action for a Carcinogenic Hair Dye Ingredient."
  - Directed and prepared a report to OSHA entitled "Formaldehyde Risk Assessment for Occupationally Exposed Workers" and assisted in developing guidelines for interspecies data extrapolation for use by OSHA in its revised cancer policy.
  - Provided litigation toxicity support to private and industry clients and assisted private concerns in the development of testing protocols for the purposes of fulfilling regulatory requirements.
  - Served as expert reproductive toxicity witness in the House of Representative's hearing on the "Relationships of Exposure to Toxic Chemicals and Reproductive Impairment."
- Staff Science Advisor, Office of Health Affairs, Commissioners Office, Food and Drug Administration.
    - Provided professional scientific expertise in pharmacology, toxicology, biochemistry, and biology requisite to the effective accomplishment of the Agency's scientific overview and leadership function. Served as technical expert, scientific advisor, and liaison of the Commissioner at the Bureau level on matters relevant to toxicologic and pharmacologic safety assessment of toxic substances to which humans and animals are exposed.
    - Responsibilities included performing risk assessments on compounds (such as PCB's, methapyraline, caffeine, etc.) that could be utilized by the Commissioner's office in choosing between various regulatory options. Routinely reviewed toxicity data on compounds as diverse as caffeine, methylsalicylate, xylitol, and sodium nitrite. Provided litigation support and testified as expert toxicology witness at the Administrative Law Judge hearing on the safety of cyclamate.
    - Participated in the National Toxicology Program (NTP). Duties included review of agency nominations for toxicity testing by NTP, research planning for the regulatory needs of FDA, preparation of the Annual Carcinogen Report, and participation in numerous NTP workgroups.
    - Interagency and international initiatives included chairing the Interagency Regulatory Liaison Group (IRLG) Workgroup on Reproductive Toxicity Risk Assessment. Planned, coordinated, and published a three-day symposium entitled, "The Effects of Foods and Drugs on the Development and Function of the Nervous System." Chaired and organized a national workshop on Reproductive Toxicity Risk Assessment.
    - Served as Project Director of a contract with the Environment Teratology Information Computer Division, Oak Ridge National Research Laboratory to develop computerized teratology literature data extraction, indexing, and collation. This data base system is being

**VASILIOS H. FRANKOS, Ph.D.**

integrated into the ETIC master computer file and will be manipulated through the INQUIRE data base management system.

- Senior Toxicologist, Division of Toxicology, Center for Food Safety and Applied Nutrition, Food and Drug Administration.
  - Responsible for the toxicologic evaluation of compounds of special interest to the agency. Required expert review of data on carcinogenicity, promotion, reproduction, teratology, and pharmacokinetics. Presented scientific evaluations to congressional, departmental, and agency directors. Represented the agency on sensitive scientific issues. Areas of emphasis included evaluation of toxicity data on artificial sweeteners (saccharin, cyclamate, xylitol, etc.), development of reproductive toxicity risk assessment criteria, pharmacokinetics evaluation of compounds (methylene chloride, sodium nitrite, saccharin, etc.), and risk assessment.
  - Directed the development of a PC computer network to be used by the Review Toxicologist of the FDA Bureau of Foods. This system allows direct access to toxicology data bases at NLM, Oak Ridge, Dialog, and extensive FDA internal data bases and full manipulation, storage, and collation of personalized literature data bases.
- Toxicologist, Division of Toxicology, Bureau of Foods, Food and Drug Administration.
  - Responsible for toxicologic review of substances used as direct and indirect food additives. Duties required discussions with manufacturers, formulators, toxicologists, universities, and other agencies on experimental procedures for showing safety and estimating risk.
  - Additional duties and accomplishments included primary involvement in planning, testing, and implementing new proposed regulations that direct the priority assessment review of all food additives. Duties included designing and implementing modern toxicity testing protocols, protocol quality parameters, criteria for utilizing toxicity data, and computer compatible toxicology test summarization forms. Co-led a 12 person toxicology cyclic review team. This endeavor was awarded a commendable service citation.
  - Served as the toxicology member on the Program Advisory Board for the FDA's effort to modernize the storage and retrieval of safety information in the Bureau of Foods. Over two million dollars was expended for the computerization and microfiling of all Bureau petition files. This system is now fully operational within the FDA and is called SIREN.
- Research Assistant, Department of Pharmacology and Drug Abuse, Maryland Psychiatric Research Center.
  - Laboratory responsibilities included the daily analysis of the catecholamines and their metabolites in the urine of patients under different drug therapies. Used column chromatography to separate the monoamines and their metabolites with subsequent fluorometric determinations of their amounts. Techniques involved using radioisotopic methods to determine monoamine oxidase kinetics, substrate km, and oxygen requirements and preparing mitochondrial isolates to study by oxygraphic assay and radioisotopic methods.
  - Clinical responsibilities included the collection, collation, and statistical analysis of data obtained by researchers in various disciplines, such as clinical and physiological psychology, biochemistry, and pharmacology. Assisted in the design, implementation,



## **VASILIOS H. FRANKOS, Ph.D.**

and statistical analysis of a study concerning the use of naloxone and cyclazocine as narcotic antagonists in a population of paroled drug addicts. Implementation of the study involved administering, scoring, and statistically analyzing psychodiagnostic and intelligence tests.

### **HONORS**

- 1980      FDA Commendable Service Award for Direct Food Additive Cyclic Review
- 1969-70   Baltimore University Club Scholarship
- 1966-67   American Hellenic Education Scholarship

### **PROFESSIONAL ACTIVITIES AND MEMBERSHIPS**

Invited presentation entitled "Developmental and Reproductive Toxicity Testing of Implantable Medical Devices." Presentation for: 1996 Summer Short Course Series on Medical Device Biocompatibility: From Material Screening to Final Product Testing. Sponsored by: Case Western Reserve University. July 18-19, 1996, Baltimore, Maryland.

Invited presentation entitled "Risk Assessment: What is it? How can I use it?" Regional Meeting of the Association of Official Analytical Chemists (AOAC), October 27, 1994, College Park, Maryland.

Invited presentation entitled "Testing Requirements for Medical Devices" Annual Genetic Toxicology Workshop, May 3-5, 1993, Rockville, Maryland.

Society of Regulatory Toxicology and Pharmacology 1990 -

Food and Drug Law Institute 1990 -

Guest Lecturer, University of Vermont Law School "Risk Assessment at the Law" June 4-14, 1990, Burlington, Vermont.

Invited presentation entitled "Review of Safety and Toxicity of Sanguinaria and Sanguinarine" Symposium on Sanguinaria, April 25, 1990, Toronto, Ontario, Canada.

Invited presentation entitled "Health Risks and Safety Precautions" PCB Compliance, Cleanup and Disposal, March 27-28, 1990, Toronto, Ontario, Canada.

Environmental Law Institute 1989 -

**VASILIOS H. FRANKOS, Ph.D.**

Invited panel member "Weight of Evidence Considerations in Identifying Reproductive and Developmental Toxicants" Risk Assessment Issues in Developmental and Reproductive Toxicology, Sept. 18-19, 1989, Berkeley, California.

Invited presentation entitled "Health Risk and Safety Precautions" Current Issues in PCB Compliance, May 24-25, 1989, Toronto, Canada.

Regulatory Affairs Professionals Society 1988 -

Invited presentation entitled "Risk Assessment for Effects Other than Cancer" 1986 Conference for Food Protection, Aug. 17-20, 1986, Ann Arbor, Michigan.

Invited work group member at the EPA sponsored "Consensus Workshop on the Evaluation of Maternal and Developmental Toxicity" May 12-14, 1986, Rockville, Maryland.

Co-author of an invited paper entitled "Acrylonitrile as a Carcinogen: Research Needs for Better Risk Assessment" presented at the International Conference on "Occupational and Environmental Significance of Industrial Carcinogens" October 7-9, 1985, Milan, Italy.

Invited presentation entitled "FDA Perspectives on the use of Teratology Data for Human Risk Assessment" Symposium on "Risk Assessment for Developmental Toxicity" Annual meeting of the Society of Toxicology, March 13, 1984, Atlanta, Georgia.

Guest Lecturer and Expert Consultant to the Environmental Teratology Information Computer Division, Oak Ridge National Research Laboratory. 1982.

Expert Witness before the U.S. House of Representatives Committee on Science and Technology hearing on the "Relationship of Exposure to Toxic Chemicals and Reproductive Impairment," 1982.

Guest speaker, FDA National Scientific Health Professionals Meeting; spoke on the National Toxicology Program. 1981.

Member, National Toxicology Program Chemical Evaluation Committee. 1979-1982.

Member, National Toxicology Program Annual Carcinogen Report Workgroup. 1979-1982.

Invited workshop panel member, Workshop on Biological and Statistical Implications of the ED Study on Related Data Bases, Mt. Sterling, Ohio; sponsored by the Society of Toxicology for the National Center for Toxicological Research. 1981.

Chairman - Interagency Regulatory Liaison Group - Reproductive Toxicity Risk Assessment Group. 1980-1981.

## VASILIOS H. FRANKOS, Ph.D.

Chairman and Organizer of the IRLG - Reproductive Toxicity Risk Assessment Workshop held at FDA, Rockville, MD. 1981.

Participated in preparation of the National Science Foundation Third Annual Science and Technology Report. 1980.

Participated in Preparation of the National Science Foundation Five-Year Science Outlook. 1980.

Participated in publication of the First Annual Report on Carcinogens (DHHS/NTP). 1980.

Organizer and lecturer for a State Department and DHHS/FDA sponsored course conducted for the National Organization for Drug Control and Research (Cairo, Egypt), entitled: "A Course in Chronic and Reproductive Toxicity Testing and Risk Assessment." 1980.

Guest lecturer at FDA Consumer Exchange Meeting: Use of Risk Assessment for Decision-Making. 1980.

Co-chairman and organizer of the Fifth FDA Science Symposium on Effects of Foods and Drugs on the Development and Function of the Nervous System: Methods for Predicting Toxicity. Delivered presentation entitled "Symposium Summary and Future Directions for Neurotoxicology Testing." 1979.

Expert witness FDA Administrative Law Judge hearing on the Safety of Cyclamate Used as an Artificial Sweetener. 1979.

## PUBLICATIONS

Kruger, C.L., M.H. Whittaker, and V.H. Frankos. 1999. Genotoxicity Tests on D-Tagatose. *Regulatory Toxicology and Pharmacology*. 29(2):S36-S42.

Kruger, C.L., M.H. Whittaker, and V.H. Frankos. 1999. Developmental Toxicity of D-Tagatose. *Regulatory Toxicology and Pharmacology*. 29(2):S29-S35.

Turnbull, D., M.H. Whittaker, V.H. Frankos, and D. Jonker. 1999. 90-Day Oral Toxicity Study of D-Tagatose in Rats. *Regulatory Toxicology and Pharmacology*. 29(2):S1-S10.

Trunbull, D., M.H. Whitaker, V.H. Frankos, and D. Jonker. 1999. 13-Week Oral Toxicity Study with Stanol Esters in Rats. *Regulatory Toxicology and Pharmacology*. 29(1):216-226.

Whittaker, M.H., V.H. Frankos, D.H. Waalkens-Berendsen, A.P.M. Wolterbeek. 1999. 2-Generation Reproductive Toxicity Study of Plant Stanol Esters in Rats. *Regulatory Toxicology and Pharmacology*. 29(1):196-204.

Hester, T.R., N.F. Ford, P.J. Gale, J.L. Hammett, R. Raymond, D. Turnbull, V.H. Frankos, and M.B. Cohen. 1997. Measurement of 2,4-toluenediamine in urine and serum samples from women with mème or replicon breast implants. *Plastic and Reconstructive Surgery*. 100(5):1296-1298.

VASILIOS H. FRANKOS, Ph.D.

- Silverstein, B., K.M. Witkin, V.H. Frankos, and A.I. Terr. 1997. Assessing the Role of the Biomaterial Aquavene in Patient Reactions to Landmark Midline Catheters. *Regulatory Toxicology and Pharmacology* 25(1).
- Rodricks, J.V., V.H. Frankos, and L.M. Plunkett. 1995. Food Additives. In: *Regulatory Toxicology*. C.P. Chengeliss, J.F. Holson and S.C. Gad (eds.) Raven Press, New York, New York, 51-82.
- Spiegel, J.E., R. Rose, P. Karabell, V.H. Frankos, and D.F. Schmitt. 1994. Safety and Benefits of Fructooligosaccharides as Food Ingredients. *Food Technology* 85-89.
- Redenbaugh, K., T. Berner, D. Emlay, V. Frankos, W. Hiatt, C. Houck, M. Kramer, L. Malyj, B. Martineau, N. Rachman, L. Rudenko, R. Sanders, R. Sheehy, and R. Wixtrom. 1993. Regulatory Issues for Commercialization of Tomatoes with an Antisense Polygalacturonase Gene. *In Vitro Cell. Dev. Biol.* 29:17-26.
- Frankos, V.H., D.F. Schmitt, L.C. Haws, A.J. McEvily, R. Iyengar, S.A. Miller, I.C. Munro, F.M. Clydesdale, A.L. Forbes, and R.M. Sauer. 1991. Generally Recognized as Safe (GRAS) Evaluation of 4-Hexylresorcinol for Use as a Processing Aid for Prevention of Melanosis in Shrimp. *Regulatory Toxicology and Pharmacology* 14:202-212.
- Wilcock, K.E., A.B. Santamaria, V.H. Frankos, H.W. Fischer, F. Laden, E.A. Platz, and B.A. Jackson. 1990. Perspectives on Adverse Reaction Rates Associated with the Use of High Osmolar Ionic and Low Osmolar Nonionic Contrast Media. *Journal of the American College of Toxicology* 9(6):563-607.
- Frankos, V.H., D.J. Brusick, E.M. Johnson, H.I. Maibach, I. Munro, R.A. Squire, and C.S. Weil. 1990. Safety of Sanguinaria Extract as Used in Commercial Toothpaste and Oral Rinse Products. *Journal of the Canadian Dental Association* 56(7 Suppl):41-47.
- Schmitt, D., V. Frankos, and D. Richardson. 1990. Toxicologic Evaluation of Sanguinaria Extract. Eleventh Annual Meeting of the American College of Toxicology. Program and Abstracts. Abstract.
- Schmitt, D., V. Frankos, J. Westland, and T. Zoetis. 1990. Toxicological Evaluation of Cellulon™ Fiber: Genotoxicity, Acute and Subchronic Toxicity. Eleventh Annual Meeting of the American College of Toxicology. Program and Abstracts. Abstract.
- Rudenko, L., J. Adgate, M. Aponte-Pons, T. Berner, V. Frankos, R. Gregory, C.K. Lintner, N. Rachman, W. Sherman, T. Winters, and R. Wixtrom. 1990. Application of Risk Assessment Methodology to Genetically Engineered Food Products: A Generic Approach. Society for Risk Assessment. Abstract.

VASILIOS H. FRANKOS, Ph.D.

- Frankos, V., M. Stedman, and M.A. Friedman. 1989. A lifetime oncogenicity study of acrylamide administered to F344 rats in the drinking water. Tenth Annual meeting of the American College of Toxicology. Program and Abstracts. p.26. Abstract.
- Hanson, C.F., V.H. Frankos, and W.O. Thompson. 1989. Bioavailability of oxalic acid from spinach, sugar beet fibre and a solution of sodium oxalate consumed by female volunteers. *Fd. Chem. Toxic.* 27,3:181-184.
- Frankos, V., L.H. Dulak, M.A. Fiedman. 1989. Use of risk assessment in the statistical design of a carcinogenicity bioassay of acrylamide. *The Toxicologist* 9:179. Abstract.
- Strother, D.E., R.W. Mast, R.C. Kraska, and V. Frankos. 1988. Acrylonitrile as a carcinogen. Research needs for a better risk assessment. *Annals of the New York Academy of Sciences* 534:169-178.
- Hanson, C.F., V.H. Frankos, and W.O. Thompson. 1988. Low dietary availability of oxalic acid present in refined sugar beet pulp compared to spinach and sodium oxalate. *The Toxicologist* 8:88. Abstract.
- Strother, D.E., R.W. Mast, R.C. Krashka, and V. Frankos. 1988. Acrylonitrile as a Carcinogen: Research Needs for Better Risk Assessment, *Annals of the New York Academy of Sciences* 534:169-178.
- Schwetz, A. Bernard, R.W. Tyl et al. 1987. Consensus workshop on the evaluation of maternal and developmental toxicity work group III report: Low dose extrapolation and other considerations for risk assessment - models and applications. *Teratogenesis, Carcinogenesis, and Mutagenesis* 7:321-327.
- Frankos, V.H. and S.H. Rieth. 1987. Safety factors applied to various FDA pregnancy class drugs. *The Toxicologist* 7. Abstract.
- Kimmel, C.A., G.L. Kimmel, and V. Frankos. 1986. Editors IRLG Workshop on Reproductive Toxicity Risk Assessment. *Environmental Health Perspectives*, Vol 66, 193-221.
- Rodricks, V. Joseph, V. Frankos, D. Turnbull, R.G. Tardiff. 1986. Risk assessment for effects other than cancer. In *Food Protection Technology. Proceedings of the 1986 Conference for Food Protection.* Lewis Publishers, Inc.
- Frankos, V.H. 1985. FDA perspectives on the use of teratology data for human risk assessment. *Fundamental and Applied Toxicology*. Vol. 5, 615-625.
- Flamm, W.G. and V. Frankos. 1985. Formaldehyde: Laboratory Evidence. In "Interpretation of Negative Epidemiological Evidence for Carcinogenicity." IARC Scientific Publication 65.

VASILIOS H. FRANKOS, Ph.D.

- Siegel, D.M., V.H. Frankos, and M.A. Schneiderman. 1983. Formaldehyde risk assessment for occupationally exposed workers. *Regulatory Tox. and Pharm.* 3, 355-371.
- Frankos, V.H. 1982. Relationship of exposure to toxic chemicals and reproductive impairment. Expert testimony for Committee on Science and Technology, U.S. House of Representatives July 27, 1982. United State Government Printing Office.
- Frankos, V.H. 1982. Qualitative comparison of chemical teratogenesis in humans and animal species (Abstract). Third Annual Meeting of the American College of Toxicology, Washington, D.C.
- Siegel, D.M., V.H. Frankos, and M.A. Schneiderman. 1982. Formaldehyde risk assessment for occupationally exposed workers (Abstract). Third Annual Meeting of the American College of Toxicology, Washington, D.C.
- Frankos, V.H. and J. Wassom. 1982. Computerized teratology literature data extraction, indexing, and collation (Abstract). *Teratology*. 25:2 80A.
- Frankos, V.H. Symposium Summary and Suggested Future Directions for Detection of Neurotoxicity. 1980. In *The Effects of Foods and Drugs on the Development and Function of the Nervous System: Methods for Predicting Toxicity*. Edited by Gryder, R.M. and Frankos, V.H. U.S. Department of Health and Human Services, Food and Drug Administration, Publication No. DHHS/FDA 80-1076.
- Frankos, V.H. 1980. Reproductive toxicity risk assessment task group: Outline of work plan and request for comments. *Federal Register* 45:63553-63554.
- Gryder, R.M. and V.H. Frankos (eds.). 1980. *The Effects of Foods and Drugs on the Development and Function of the Nervous System: Methods for Predicting Toxicity*. U.S. Department of Health and Human Services, Food and Drug Administration.
- Frankos, V.H. and R. Platt. 1976. Glycerol accumulation and water content in larvae of *Limenitis archippus*: Their importance to winter and survival. *J. Insect Physiol.* 22:623-623.
- Frankos, V.H. and G. Butterbaugh. 1976. Characterization of norepinephrine metabolism following simultaneous intraventricular injection of H<sup>3</sup>-L-tyrosine and C<sup>14</sup>-DL-norepinephrine. *Pharmacologist* 18:135.
- Messiha, K.F., E. Bakutis, and V.H. Frankos. 1973. Simultaneous separation of acid metabolites of catecholamines: Application to urine and tissue. *Clin. Chim. Acta.* 45:159-164.

ATTACHMENT 9



Animal Sciences and Industry  
K-State Research and Extension  
232 Weber Hall  
Manhattan, KS 66506-0201  
785-532-6533  
Fax: 785-532-7059

August 8, 2001

Eric F. Greenberg  
Of Counsel  
Ungaretti & Harris  
3500 Three First National Plaza  
Chicago, IL 60602-4283

Dear Mr. Greenberg:

The purpose of this letter is to confirm that I believe the use of a small quantity of carbon monoxide in the modified packaging system known as ActiveTech (by PACTIV) is safe and should qualify as GRAS. I have been a research meat scientist for nearly 30 years and have focused most of that time on factors affecting meat color and shelf life, including packaging systems. Thus, I am familiar with most of the world literature on such systems.

Based on my review of the details of the ActiveTech 2001 modified atmosphere system employing 0.4% carbon monoxide gas in a mixture with 60 percent carbon dioxide and the remainder nitrogen, as well as the published literature and common knowledge in the field, I am confident that the use of modified atmosphere including small quantities of carbon monoxide (0.4%) to package fresh meats as used in ActiveTech 2001 system is both safe and generally recognized as safe.

Sincerely,

Melvin C. Hunt  
Professor

Kansas State University  
Agricultural Experiment  
Station and Cooperative  
Extension Service

"Knowledge  
life"



## MELVIN C. HUNT

Department of Animal Sciences and Industry  
Weber Hall  
Kansas State University  
Manhattan, KS 66506-0201

Ph: 913-532-1232  
Fax: 913-532-7059  
[Hhunt@oznet.ksu.edu](mailto:Hhunt@oznet.ksu.edu)

### PERSONAL DATA:

Born: February 10, 1942

Married: Rae Jean Opie, August 20, 1965; Daughters: Paige and Holly

### EDUCATION:

B.S. 1965 Animal Husbandry, Kansas State University, Manhattan, KS

M.S. 1970 Animal Science, Kansas State University, Manhattan, KS

Ph.D. 1973 Food Science, University of Missouri, Columbia, MO

### PROFESSIONAL EXPERIENCE:

- 1991- Chair, Undergraduate Food Science Program
- 1984- Professor, Kansas State University: 50% Teaching - 50% Research
- 1978-84 Associate Professor, Kansas State University
- 1975-78 Assistant Professor, Kansas State University
- 1973-75 Research Chemist, Tennessee Eastman Company
- 1968-73 Grad Research Assistant, Kansas State and University of Missouri
- 1966-68 Taught high school chemistry and biology, Kinsley, KS

### PROFESSIONAL AFFILIATIONS:

#### American Meat Science Association:

- President, 1995-96; Past-President, 1996-97
- Director and Executive Board, 1989-91
- Chair 1991 Reciprocal Meat Conference
- Parliamentarian
- Chair or member of numerous committees including:
  - Meat Color Guidelines, AMSA Teaching Award, Undergraduate Travel Award, Grad Student Poster Competition, Teaching Display, Resolutions, Meat Tenderness, Biochemistry-Biophysics, Packaging, Meat Color, Growth and Development, Reciprocation, Long Range Planning, Sustaining Membership, Endowment, and Research Priorities.

#### American Society of Animal Science:

- Chair and Chair-elect, Meat Science-Muscle Biology Section of National ASAS Meeting
- Chair, Midwestern ASAS Meat Science Section
- Editorial Board Journal Animal Science
- Teaching Award Committee, Midwestern ASAS Section

#### Institute of Food Technologists:

- Chair and Chair-elect of Muscle Foods Division, 1992-94
- Director of Muscle Foods Division
- Chair of Muscle Foods Nominating Committee
- Committee for two National Muscle Foods Symposia
- Journal of Food Science, Manuscript Review

CAST: Contributing member

Journal of Muscle Foods: Editorial Board

Meat Science: Manuscript Review

### **HONORARY AFFILIATIONS:**

Phi Kappa Phi, Sigma Xi, Phi Tau Sigma, Gamma Sigma Delta, Alpha Zeta

### **HONORS:**

- College of Agriculture Outstanding Faculty Award 1979
- College of Agriculture Outstanding Faculty Award 1982
- College of Agriculture Outstanding Faculty Award 1988
- College of Agriculture Outstanding Faculty Award 1998
- College of Agriculture Outstanding Academic Advisor 1983
- University Selection for Parents' Day Lecture 1979
- Outstanding Lecturer Award, ITAL, Campinas, Brazil 1981
- Honorary State Farmer Degree 1985
- Distinguished Teaching Award, Gamma Sigma Delta 1989
- Selected Instructor, National Food Science Satellite Program 1990
- Certificate of Meritorious Service, Kansas Ag Teachers Association 1992
- CASE Professor of the year, Kansas winner of national competition 1992
- Outstanding Advising Award, Gamma Sigma Delta 1994
- Distinguished Teaching Award, American Meat Science Association 1994
- Outstanding Food Scientist, Phi Tau Sigma 1996
- Outstanding KSU Instructor & Advisor Award, Mortar Board 1997
- Signal Service Award, American Meat Science Association 1997
- USDA Food & Agriculture Science Excellence in Teaching Award, 2000

### **DEPARTMENT, COLLEGE OF AG, AND UNIVERSITY ACTIVITIES:**

- Faculty Advisor: Block and Bridle, 6 years
- Faculty Advisor: Food Science Club, 3 years
- Faculty Advisor: Animal Science Grad Student Association, 16 years
- Faculty Advisor: Ag Student Council, elected for 2 terms (4 years)
- Chair, Weber Hall Building/Renovation Project
- Chair, KSU Meat Science Faculty
- Coordinator of KSU Meat Research Labs
- ASI Graduate Student Selection Committee
- ASI Undergraduate Career Development Committee
- ASI Library Committee
- ASI Scholarship, Loans and Honors Committee
- Department Representative for Gamma Sigma Delta, 10 years
- Student Team Coordinator, ASI Quadrathlon Teams
- Agriculture Student of the Month Selection Committee
- Agriculture Faculty of the Semester Selection Committee
- College of Agriculture Course and Curriculum Committee, chair and member
- College of Agriculture Academic Standards Committee, chair and member
- College of Agriculture Commencement Committee
- University Faculty Senator, College of Agriculture, two terms (6 years)
- University Academic Affairs Committee
- University Coordinating Committee for United Way
- KAES NCR-121 Chair and Secretary: Food & Feed Safety in Animal Production
- Food Science Undergraduate and Graduate Steering Committees
- Chair, Non-Traditional Studies Advisory Committee
- Elected by peers to ASI Teaching Advisory Committee
- Chair, KSU Undergraduate Food Science Program: Coordinate all course & curriculum and policy matters, scholarship, internships, recruitment, and record keeping

### **INDUSTRY-EXTENSION ACTIVITIES:**

- Numerous presentations at: MidWest Meat Processors Seminars, Kansas-Nebraska Curing and Sausage Short Courses, KSU Cattlemen's Day, KSU Swine Day
- Technical Assistance for: Tennessee Eastman Company, Ross Industries, Giant Food

Stores, Excel Corporation, IBP, Dorskocil Companies, Tenneco Packaging, Farmland, National Beef, Cryovac, Buckhead Beef, Dupont, Kalsec, Wendy's, Greater Omaha Beef, Hormel

- State FFA Livestock Awards Selection Committee
- State FFA Star Farmer Selection Committee
- State FFA Public Speaking Contest Judge
- Kansas Jr. Livestock Carcass Contest Judge
- Kansas Meat Processor Cured Meat Show Judge
- Missouri Meat Processor Cured Meat Show Judge

#### **TEACHING RESPONSIBILITIES:**

##### **Current Courses - KSU Campus:**

- ASI 350 Meat Science. 3hr. Lecture-lab introductory meat science  
Enrollment: Since 1979, 2031 students; currently running at maximum seating of 72
- ASI 610 Processed Meat Operations. 2hr. 50% responsibility, value-added processing  
Enrollment: 6 to 12 undergraduate and graduate students; since 1988, 35 students
- ASI 930 Advanced Meat Science. 3hr. Team-taught, highest level meats course  
Enrollment: Varies from 6 to 15 graduate students
- GENAG 500 Food Science Seminar. 1hr. Seminar for graduating seniors  
Enrollment: Varies from 6 to 15 students

##### **Current Courses - KSU Distance Learning Program:**

- ASI 340 Principles of Meat Science. 2hr. Web-based course for Continuing Education  
Enrollment: Since 1987, over 680 students
- GENAG 500 Food Science Seminar. 1hr. Seminar series for Distance Learning majors  
Enrollment: 3 to 15 undergraduate students per year, Continuing Education
- GENAG 630 Food Science Problems. 1hr. Detailed written investigation of current topics  
Enrollment: 2 to 8 students per year through Continuing Education

##### **Previously Taught Courses:**

- Topics in Meat Science and Muscle Biology
- Meats Judging Team (at University of Missouri)
- Meat Processing
- Livestock and Meat Evaluation
- Animal Agriculture and Consumers

#### **INTERNATIONAL COURSE ACTIVITIES:**

- Meat Science and Technology Short Course for Latin America, Institute for Food Technology, Campinas, Brazil, 6 weeks, one of two international lecturers
- Meat Science Facilities, University of Monterrey, Monterrey, Mexico
- Lecturer for five KSU International Meat Science Courses, International Meat and Livestock Program, Kansas State University
- Sabbatical leave, fall 1992, visiting scientist to Norwegian Food Research Institute, As
- Have attended 8 International Congresses of Meat Science and Technology

#### **ADVISING RESPONSIBILITIES:**

- Undergraduate Advisees: average of 26 for the last 10 years
- Graduate Students Supervised:      Graduate Student Committees:
  - 12 Masters Students                      - 43 Masters
  - 6 PhD Students                              - 20 PhD
- Coordinate student-company relations for employment and internships for FSI

#### **RESEARCH INTERESTS:**

- Myoglobin chemistry and meat color, Methods of color measurement, Cooked meat color and food safety, Postmortem factors affecting meat quality, Collagen chemistry, Low-fat ground beef and processed meats; Six major company packaging projects funded since 1994 dealing with shelf life, color life, cold chain management, product palatability, and microbiology

#### **PUBLIC AND COMMUNITY ACTIVITIES:**

- Manhattan Optimist Club: committees for many youth activities
- Coach, Girls (16-18) ASA fast pitch softball traveling team
- Executive Committee, Riley County Extension Council
- Asst. Superintendent, sheep division, Riley County Fair
- Judge at Manhattan High School oratorical contest
- FarmHouse Fraternity, alumni board and committee work
- Snyder Award for Alumni Service, FarmHouse Fraternity
- Activities of First Presbyterian Church

Melvin C. Hunt  
Professor  
Department of Animal Sciences and Industry  
Kansas State University

Refereed Journal Articles

- Hunt, M.C., R.A. Smith, D.H. Kropf and H.J. Tuma. 1975. Factors affecting showcase color stability of frozen lamb in transparent film. *J. Food Sci.* 40:637.
- Hunt, M.C. and H.B. Hedrick. 1977. Profile of fiber types and related properties of five bovine Muscles. *J. Food Sci.* 42:513.
- Hunt, M.C. and H.B. Hedrick. 1977. Histochemical and histological characteristics of bovine muscle from four quality groups. *J. Food Sci.* 42:578.
- Hunt, M.C. and H.B. Hedrick. 1977. Chemical, physical and sensory characteristics of bovine muscle from four quality groups. *J. Food Sci.* 42:716.
- Thomas, J.D., D.M. Allen, M.C. Hunt and C.L. Kastner. 1977. Nutritional regimen, post-slaughter conditioning temperature, and vacuum packaging effects on beef carcass and retail cut bacteriology. *J. Food Prot.* 40:678.
- Harrison, A.R., M.E. Smith, D.M. Allen, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1978. Nutritional regimen effects on quality and yield characteristics of beef. *J. Anim. Sci.* 47:383.
- Loveday, H.D., M.E. Dikeman, M.C. Hunt and A.D. Dayton. 1978. Adipose tissue water related to bovine carcass composition. *J. Anim. Sci.* 47:606.
- Smith, M.E., C.L. Kastner, M.C. Hunt, D.H. Kropf and D.M. Allen. 1979. Elevated conditioning temperature effects on carcasses from four nutritional regimens. *J. Food Sci.* 44:158.
- Gutowski, G.H., M.C. Hunt, C.L. Kastner, D.H. Kropf and D.M. Allen. 1979. Vacuum aging, display, and level of nutrition effects on beef quality. *J. Food Sci.* 44:140.
- Fung, D.Y.C., C.L. Kastner, M.C. Hunt, M.E. Dikeman and D.H. Kropf. 1980. Mesophilic and psychrotrophic populations on hot-boned and conventionally processed beef. *J. Food Prot.* 43:547.
- Hayward, L.H., M.C. Hunt, C.L. Kastner and D.H. Kropf. 1980. Blade tenderization effects on beef longissimus sensory and Instron textural measurements. *J. Food Sci.* 45:925.
- Harrison, A.R., D.H. Kropf, D.M. Allen, M.C. Hunt and C.L. Kastner. 1980. Relationships of spectrophotometric reflectance measurements to beef muscle visual color. *J. Food Sci.* 45:1052.
- Burson, D.E., M.C. Hunt, D.M. Allen, C.L. Kastner and D.H. Kropf. 1980. Ration energy density and time on feed effects on beef longissimus palatability. *J. Anim. Sci.* 51:875.
- Fung, D.Y.C., C.L. Kastner, C-Y. Lee, M.C. Hunt, M.E. Dikeman and D.H. Kropf. 1981. Initial chilling rate effects on bacterial growth of hot-boned beef. *J. Food Prot.* 44:539.
- Wu, J.J., D.M. Allen, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1981. Nutritional effects on beef collagen characteristics and palatability. *J. Anim. Sci.* 53:1256.
- Hall, J.B. and M.C. Hunt. 1982. Collagen solubility of A-maturity bovine longissimus muscle as affected by nutritional regimen. *J. Anim. Sci.* 55:321.
- Sleper, P.S., M.C. Hunt, D.H. Kropf, C.L. Kastner and M.E. Dikeman. 1983. Electrical stimulation effects on myoglobin properties of bovine longissimus muscle. *J. Food Sci.* 48:479.

- Axe, J.E., Bowles, C.L., Kastner, M.E., Dikeman, M.C., Hunt, D.H., Kropf and G.A. Milliken. 1983. Effects of beef carcass electrical stimulation, hot boning, and aging on unfrozen and frozen longissimus dorsi and semimembranosus steaks. *J. Food Sci.* 48:332.
- Lyon, M., C.L. Kastner, M.E. Dikeman, M.C. Hunt, D.H. Kropf and J.R. Schwenke. 1983. Effects of electrical stimulation, aging, and blade tenderization hot-boned beef psoas major and triceps brachii muscles. *J. Food Sci.* 48:131.
- Greathouse, J.R., M.C. Hunt, M.E. Dikeman, L.R. Corah, C.L. Kastner and D.H. Kropf. 1983. Ralgro implanted bulls: performance, carcass characteristics, longissimus palatability and carcass electrical stimulation. *J. Anim. Sci.* 57:355.
- Burson, D.B., M.C. Hunt, D.E. Schafer, D. Beckwith and J.R. Garrison. 1983. Effects of stunning method and time interval from stunning to exsanguination on blood splashing in pork. *J. Anim. Sci.* 57:918.
- Shivas, S.D., D.H. Kropf, M.C. Hunt, C.L. Kastner, J.L.A. Kendall and A.D. Dayton. 1984. Effects of ascorbic acid on the display life of ground beef. *J. Food Prot.* 47:11.
- Choi, Y.I., C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1984. Effects of electrical stimulation and hot boning on functional characteristics of preblended beef muscle in model systems. *J. Food Sci.* 49:867.
- Claus, J.R., D.H. Kropf, M.C. Hunt, C.L. Kastner and M.E. Dikeman. 1984. Effects of beef carcass electrical stimulation and hot boning on muscle display color of polyvinylchloride packaged steaks. *J. Food Sci.* 49:1021.
- Kropf, D.H., M.E. Dikeman, M.C. Hunt and H.R. Cross. 1984. Lighting type and intensity effects on beef carcass grade factors. *J. Anim. Sci.* 59:105.
- Shivas, S.D., C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1985. Effects of electrical stimulation, hot boning, and chilling on bull semimembranosus muscle. *J. Food Sci.* 50:36.
- Claus, J.R., D.H. Kropf, M.C. Hunt, C.L. Kastner and M.E. Dikeman. 1985. Effects of beef carcass electrical stimulation and hot boning on muscle display color of unfrozen vacuum packaged steaks. *J. Food Sci.* 50:881.
- Dikeman, M.E., A.D. Dayton, M.C. Hunt, C.L. Kastner, J.B. Axe and H.J. Ilg. 1985. Conventional versus accelerated beef production with carcass electrical stimulation. *J. Anim. Sci.* 61:573.
- Burson, D.E. and M.C. Hunt. 1986. Proportion of collagen types I and III in four bovine muscles differing in tenderness. *J. Food Sci.* 51:51.
- Burson, D.E. and M.C. Hunt. 1986. Heat-induced changes in the proportion of types I and III collagen in bovine longissimus. *Meat Sci.* 17:153.
- Burson, D.E., M.C. Hunt, J.A. Unruh and M.E. Dikeman. 1986. Proportion of types I and III collagen in longissimus collagen from bulls and steers. *J. Anim. Sci.* 63:453.
- Flores, H.A., C.L. Kastner, D.H. Kropf and M.C. Hunt. 1986. Effects of blade tenderization and trimming of connective tissue on hot-boned, restructured, pre-cooked roast from cows. *J. Food Sci.* 51:1176.
- Unruh, J.A., C.L. Kastner, D.H. Kropf, M.E. Dikeman and M.C. Hunt. 1986. Effects of low-voltage electrical stimulation during exsanguination on meat quality and display colour stability. *Meat Sci.* 18:281.
- Allen, D.M., M.C. Hunt, A. Luchiari Filho, R.J. Danler and S.J. Goll. 1987. Effects of spray-chilling and carcass spacing on beef carcass cooler shrink and grade factors. *J. Anim. Sci.* 64:165.
- McCormick, R.J., D.H. Kropf, G.R. Reeck, M.C. Hunt and C.L. Kastner. 1987. Effect of heating temperature and muscle type on porcine muscle extracts as determined by reverse phase high performance liquid chromatography. *J. Food Sci.* 52:1481.

- Kluber, E.F. III, J.E. Minton, J.S. Stevenson, M.C. Hunt, D.L. Davis, T.A. Hoagland and J.L. Nelssen. 1988. Growth, carcass traits, boar odor and testicular and endocrine functions of male pigs fed a progesterogen, Altrenogest. *J. Anim. Sci.* 66:470.
- Johnson, R.D., M.C. Hunt, D.M. Allen, C.L. Kastner, R.J. Danler and C.C. Schrock. 1988. Moisture uptake during washing and spray chilling of holstein and beef-type steer carcasses. *J. Animal. Sci.* 66:2180.
- Claus, J.R., M.C. Hunt and C.L. Kastner. 1989. Effects of substituting added water for fat on the textural, sensory, and processing characteristics of bologna. *J. Muscle Foods.* 1:1.
- Claus, J.R., M.C. Hunt, C.L. Kastner and D.H. Kropf. 1990. Low-fat, high-added water bologna: Effects of massaging, preblending, and time of addition of water and fat on the physical and sensory characteristics. *J. Food Sci.* 55:338.
- Kenney, P.B. and M.C. Hunt. 1990. Effect of water and salt content on protein solubility and water retention of meat preblends. *Meat Sci.* 27:173.
- Troyer, D.L., R.O. Oyster and M.C. Hunt. 1991. A combination histochemical stain for equine muscle. *Anat. Histol. Embryol.* 20:44.
- Whipple, G., M. Koohmaraie, M.E. Dikeman, J.D. Crouse, M.C. Hunt and R.D. Klemm. 1990. Evaluation of attributes that affect longissimus muscle tenderness in *Bos Taurus* and *Bos Indicus* cattle. *J. Anim. Sci.* 68:2716.
- Claus, J.R. and M.C. Hunt. 1991. Characteristics of low-fat, high-added water bologna formulated with textural modifying ingredients. *J. Food Sci.* 56:643.
- Whipple, G., M.C. Hunt, R.D. Klemm, D.H. Kropf, R.D. Goodband, J.L. Nelssen, R.H. Hines and B.R. Schrick. 1992. Effects of porcine somatotropin and supplemental lysine on porcine muscle histochemistry. *J. Muscle Foods.* 3:217.
- Troutt, E.S., M.C. Hunt, D.E. Johnson, J.R. Claus, C.L. Kastner, D.H. Kropf and S. Stroda. 1992. Chemical, physical, and sensory characterization of ground beef containing 5 to 30 percent fat. *J. Food Sci.* 57:25.
- Troutt, E.S., M.C. Hunt, D.E. Johnson, C.L. Kastner and D.H. Kropf. 1992. Characteristics of low-fat ground beef containing texture-modifying ingredients. *J. Food Sci.* 57:19.
- Goll, S.J., C.L. Kastner, M.C. Hunt and D.H. Kropf. 1992. Effects of glucose and internal cooking temperature on the characteristics of low fat, pre- and post-rigor restructured beef roasts. *J. Food. Sci.* 57:834.
- Nold, R.A., J.A. Unruh, M.C. Hunt and C.W. Spaeth. 1992. Effects of implanting ram and wether lambs with zeranol at birth and weaning on palatability and muscle collagen characteristics. *J. Anim. Sci.* 70:2752.
- Warren, K.E., M.C. Hunt, C.L. Marksberry, O. Sørheim, D.H. Kropf and M.J. Windisch. 1992. Modified-atmosphere packaging with carbon dioxide for bone-in pork loins. *J. Muscle Foods.* 3:283.
- Nold, R.A., J.A. Unruh, C.W. Spaeth and M.C. Hunt. 1992. Effects of implanting ram and wether lambs with zeranol on pelt characteristics and removal. *Sheep Res. J.* 8:81.
- Brester, G.W., P. Lhermite, B.K. Goodwin and M.C. Hunt. 1993. Quantifying the effects of new product development: The case of low-fat ground beef. *J. Agric. & Resource Econ.* 18:239.
- Garcia Zepeda, C.M., C.L. Kastner, D.H. Kropf, M.C. Hunt, P.B. Kenney, J.R. Schwenke and D.S. Schleusener. 1993. Utilization of surimi-like products from pork with sex-odor in restructured, precooked pork roasts. *J. Food Sci.* 58:53.
- Hague, M.A., K.E. Warren, M.C. Hunt, D.H. Kropf, C.L. Kastner, S.L. Stroda, and D.E. Johnson. 1994. Endpoint temperature, internal cooked color, and expressible juice color relationships in ground beef patties. *J. Food Sci.* 59:465-470.
- Letelier, V., C.L. Kastner, P.B. Kenny, D.H. Kropf, M.C. Hunt, and C.M. Garcia Zepeda. 1995. Flaked sinew addition to low-fat cooked salami. *J. Food Sci.* 60:245-249.

- Lavelle, C., M.C. Hunt, and D.H. Kropf. 1995. Display Life and Internal Cooked Color of Ground Beef from Vitamin E-Supplemented Steers. *J. Food Sci.* 60:1-4,6.
- Sørheim, O., D.H. Kropf, M.C. Hunt, M.T. Karwoski, and K.E. Warren. 1996. Effects of modified gas atmosphere packaging on pork loin colour, display life and drip loss. *Meat Sci.* 43:203-212.
- Campbell, R.E. and M.C. Hunt. 1996. Yields, chemical composition, and value of beef shank tissues obtained using Baader processing. *J. Anim. Sci.* 74:786-789.
- Warren, K.E., M.C. Hunt, D.H. Kropf, M.A. Hague, C.L. Waldner, S.L. Stroda, and C.L. Kastner. 1996. Chemical properties of ground beef patties exhibiting normal and premature brown internal cooked color. *J. Muscle Foods.* 7:303-314.
- Warren, K.E., M.C. Hunt, and D.H. Kropf. 1996. Myoglobin oxidative state affects internal cooked color development of ground beef patties. *J. Food Sci.* 61:513-515,519.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, M.C. Hunt, J.L. Marsden, E.J. Rubio Canas, C.L. Kastner, W.G. Kuecker, and T. Mata. 1996. Color and oxidative rancidity of gamma and electron beam-irradiated boneless pork chops. *J. Food Sci.* 61:1000-1005,1093.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, E. Chambers IV, M.E. Hollingsworth, M.C. Hunt, J.L. Marsden, C.L. Kastner, and W.G. Kuecker. 1996. Sensory analysis and consumer acceptance of irradiated boneless pork chops. *J. Food Sci.* 61:1261-1266.
- Campbell, R.E., M.C. Hunt, D.H. Kropf, and C.L. Kastner. 1996. Low-fat ground beef from desinewed shanks with reincorporation of processed sinew. *J. Food Sci.* 61:1285-1288.
- Luchsinger, S.E., D.H. Kropf, E. Chambers IV, C.M. Garcia Zepeda, M.C. Hunt, S.L. Stroda, M.E. Hollingsworth, J.L. Marsden, and C.L. Kastner. 1997. Sensory analysis of irradiated ground beef patties and whole muscle beef. *J. Sensory Studies.* 12:105-126.
- Garcia Zepeda, C.M., C.L. Kastner, J.R. Wolf, J.E. Boyer, D.H. Kropf, M. C. Hunt and C.S. Setser. 1997. Extrusion and low-dose irradiation effects on destruction of *clostridium sporogenes* spores in a beef-based product. *J. Food Prot.* 60:777-785.
- Carmack, C.F, C.L. Kastner, M.C. Hunt, D.H. Kropf, C.M. Garcia Zepeda, and J.R. Schwenke. 1997. Sensory, chemical, and physical evaluation of reduced fat ground beef patties with "natural" flavorings. *J. Muscle Foods.* 8:199-212.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, M.C. Hunt, S.L. Stroda, J.L. Marsden, and C.L. Kastner. 1997. Color and oxidative properties of irradiated whole muscle beef. *J. Muscle Foods.* 8:427-443.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, M.C. Hunt, S.L. Stroda, J.L. Marsden, and C.L. Kastner. 1997. Color and oxidative properties of irradiated ground beef patties. *J. Muscle Foods.* 8:445-464.
- Hunt, M.C., O. Sørheim, and E. Slinde. 1999. Color and heat denaturation of myoglobin forms in ground beef. *J. Food Sci.* 64:847-851.
- Powell, T.H., M.C. Hunt, and M.E. Dikeman. 1999. Enzymatic assay to determine collagen thermal denaturation and solubilization. *Meat Sci.* 54:307-311
- Powell, T.H., M.E. Dikeman, and M.C. Hunt. 2000. Tenderness and collagen composition of beef semitendinosus roasts cooked by conventional convective cooking and modeled, multi-stage, convective cooking. *Meat Sci.* 55:421-425.
- Killinger, K.M., M.C. Hunt, and R.E. Campbell. 2000. Factors affecting premature browning during cooking of ground beef purchased at retail. *J. Food Sci.* 65:585-587.



- Campbell, P.C., M.C. Hunt, P. Levis, and E. Chambers IV. 2000. Dry-aging effects on palatability of beef strip loins. *J. Food Sci.* Accepted.
- Campbell, R.G., M.C. Hunt, P. Levis, and E. Chambers IV. 2000. Dry-aging effects on palatability of beef strip loins. *J. Food Sci.* In Press.
- Sammel, L.M., M.C. Hunt, D.H. Kropf, K.A. Hachmeister, C.L. Kastner, and D.E. Johnson. 2001. Influence of chemical characteristics of beef inside and outside semimembranosus on color traits. *J. Food Sci.* In Press.
- Sammel, L.M., M.C. Hunt, D.H. Kropf, K.A. Hachmeister, C.L. Kastner, and D.E. Johnson. 2001. Comparison of assays for metmyoglobin reducing ability in beef inside and outside semimembranosus. *J. food Sci.* In Review.
- Yancey, E.J., M.C. Hunt, M.E. Dikeman, P.B. Addis, and E. Katsanidis. 2001. Effects of postexsanguination vascular infusion of cattle with a solution of saccharides, sodium chloride, phosphates, and vitamins C, E, or C+E on meat display-color stability. *J. Anim. Sci.* In Press.
- Lien, R., M.C. Hunt, S. Anderson, D.H. Kropf, T.M. Loughin, M.E. Dikeman, and J. Velazco. 2001. Effects of endpoint temperature on the internal color of pork patties of different myoglobin form, initial cooking state and quality. *J. Food Sci.* In Review.
- Lien, R., M.C. Hunt, S. Anderson, D.H. Kropf, T.M. Loughin, M.E. Dikeman, and J. Velazco. 2001. Effects of endpoint temperature on the internal color of pork loin chops of different quality. *J. Food Sci.* In Review.
- Lawrence, T. M.C. Hunt, and D.H. Kropf. 2001. Surface roughening of pre-cooked, cured beef round muscles reduces iridescence. *J. Muscle Foods.* In Review.

#### Abstracts / Posters

- Hunt, M.C., R.A. Smith, D.H. Kropf and H.J. Tuma. 1969. Factors affecting showcase color stability of frozen lamb in transparent film. *J. Anim. Sci.* 29:123. (Abstr.).
- Kropf, D.H., M.C. Hunt and A.D. Dayton. 1970. Reflectance spectrophotometry for following frozen beef and lamb color changes. *J. Anim. Sci.* 31:185. (Abstr.).
- Kropf, D.H., H.J. Tuma, J.A. Santamaria, M.C. Hunt and D.E. Schafer. 1971. Display light intensity effects on frozen beef color stability. *J. Anim. Sci.* 33:219. (Abstr.).
- Hunt, M.C. and H.B. Hedrick. 1973. Properties of bovine longissimus muscle from four quality groups. *J. Anim. Sci.* 37:263. (Abstr.).
- Allen, D.M., M.C. Hunt, C.L. Kastner, D.H. Kropf, V. Chen, A. Harrison and M.E. Smith. 1976. Improving grass-fed beef quality. *J. Anim. Sci.* 43, 236 (Abstr.).
- Kuntapanit, C., B.E. Brent, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1978. Fluorometric TBA analysis for muscle food autoxidation. *J. Anim. Sci. Abstr.* 13.
- Harrison, R.A., M.E. Smith, D.M. Allen, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1978. Nutritional regimen effects on quality and yield characteristics of beef. *Ann. Meeting Amer. Soc. Anim. Sci. Abstr.* 172.
- Harrison, A.R., D.H. Kropf, D.M. Allen, M.C. Hunt and C.L. Kastner. 1978. Nutritional regimen, vacuum packaging and display length effects on color of four beef muscles. *Ann. Meeting Amer. Soc. Anim. Sci. Abstr.* 173.
- Corte, O., D.M. Allen, B.E. Brent, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1978. Longissimus and biceps femoris collagen characteristics of beef finished on four nutritional regimens. *Ann. Meeting Amer. Soc. Anim. Sci. Abstr.* 163.

- Fung, D.Y.C., C.L. Kastner, M.C. Hunt, M.E. Dikeman and D.H. Kropf. 1979. Mesophilic and psychrotrophic populations of hot-boned and conventionally processed beef. 39th Ann. Meeting of IFT, St. Louis, MO. (Abstr.).
- Harrison, A.R., D.H. Kropf, D.M. Allen, M.C. Hunt and C.L. Kastner. 1979. Nutritional regimen, vacuum packaging and display length effects on color of four beef muscles. 39th Ann. Meeting of IFT, St. Louis, MO. (Abstr.).
- Harrison, A.R., D.H. Kropf, D.M. Allen, M.C. Hunt and C.L. Kastner. 1979. Relationship of various spectrophotometric reflectance measurements to beef muscle visual color. 39th Ann. Meeting of IFT, St. Louis, MO. (Abstr.).
- Hayward, L.H., M.C. Hunt, D.M. Allen, C.L. Kastner and D.H. Kropf. 1979. Blade tenderization effects on beef longissimus sensory, and Instron textural measurements. 39th Ann. Meeting of IFT, St. Louis, MO. (Abstr.).
- Francis, S., D. Allen and M.C. Hunt. 1980. Animal agriculture and consumers. J. Anim. Sci. Midwest ASAS (Abstr.).
- Obanion, D.S., R.R. Schalles and M.C. Hunt. 1980. Relationship among performance of yearling bulls and longissimus amino acids and histochemical characteristics. J. Anim. Sci. Midwest ASAS (Abstr.).
- Hunt, M.C., J.L.A. Kendall, M.E. Dikeman, C.L. Kastner and D.H. Kropf. 1980. Ground beef from electrically stimulated and hot-boned carcasses. 13th Ann. Meeting Midwestern Sec. of Amer. Soc. Anim. Sci. Abstr. 30.
- Wu, J.J., D.M. Allen, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1980. Nutritional effects on beef palatability and collagen characteristics. 13th Ann. Meeting Midwestern Sec. of Amer. Soc. Anim. Sci. Abstr. 33.
- Kastner, C.L., M.E. Dikeman, K.N. Nagele, M. Lyon, M.C. Hunt and D.H. Kropf. 1980. Effects of carcass electrical stimulation and hot boning on selected beef muscles. Proc. European Meeting of Meat Res. Workers. Abstr. 26:I-2, 40.
- Lyon, M., C.L. Kastner, M.E. Dikeman, M.C. Hunt, D.H. Kropf and J.R. Schwenke. 1980. Beef carcass electrical stimulation and hot boning effects on psoas major and triceps brachii muscles. Proc. Recip. Meat Conf., 33:85. (Abstr.).
- Hale, D.S., M.C. Hunt and C.L. Kastner. 1981. Fiber orientation effects on Warner-Bratzler shear values. 14th Ann. Meeting Midwestern Sec. of Amer. Soc. of Anim. Sci. (Abstr.).
- Lyon, M., C.L. Kastner, M.E. Dikeman, M.C. Hunt, D.H. Kropf and J.R. Schwenke. 1981. Beef carcass electrical stimulation and hot boning effects on psoas major and triceps brachii muscles. 14th Ann. Meeting Midwestern Sec. of Amer. Soc. of Anim. Sci. (Abstr.).
- Claus, J.R., D.H. Kropf, M.C. Hunt, C.L. Kastner and M.E. Dikeman. 1981. Muscle color display life as influenced by electrical stimulation and hot boning. J. Anim. Sci. 53 (Suppl. 1):210.
- Dikeman, M.E., S.C. Olsen, C.L. Kastner and M.C. Hunt. 1981b. Conventional versus accelerated beef production and processing. II. J. Anim. Sci. 53 (Suppl. 1):212.
- Greathouse, J.R., M.C. Hunt, M.E. Dikeman and L.R. Corah. (1982). Effects of zeranol implants on performance and carcass characteristics of bulls implanted from birth to slaughter. J. Anim. Sci. 55 (Suppl. 1):240.
- Dikeman, M.E., J.R. Greathouse, M.C. Hunt, C.L. Kastner, P.S. Sleper, D.H. Kropf and L.R. Corah. 1982. Effects of zeranol implants and electrical stimulation on palatability of bulls implanted from birth to slaughter. II. J. Anim. Sci. 55 (Suppl. 1):238.
- Hunt, M.C., P.S. Sleper, D.H. Kropf, M.E. Dikeman and C.L. Kastner. 1982. Myoglobin properties of electrically stimulated bull longissimus muscle. J. Anim. Sci. 55 (Suppl. 1):242.

- Rao, B.R., D.H. Kropf, C.L. Kastner, M.E. Dikeman and M.C. Hunt. 1982. Influence of freezing temperatures and storage time on aroma and flavor of ground beef. *J. Anim. Sci.* 55 (Suppl. 1):255.
- Burson, D.E., M.C. Hunt, D.E. Schafer and D. Beckwith. (1982). Stunning method and time interval to sticking effects on blood splashing in pork. *J. Anim. Sci.* 55 (Suppl. 1):233.
- Shivas, S.D., D.H. Kropf, C.L. Kastner, J.L.A. Kendall, M.C. Hunt and A.D. Dayton. 1982. Effects of ascorbic acid on display life of ground beef. *Internatl. Symp. Meat Science and Technol.*, Lincoln, Nebraska, Nov. 1982.
- Burson, D.E., M.C. Hunt, D.E. Schafer, D. Beckwith and J.R. Garrison. 1982. Stunning method and time interval from stunning to exsanguination effects on blood splashing in pork. *Internatl. Symp. Meat Science and Technol.*, Lincoln, Nebraska, Nov. 1982.
- Greathouse, J.R., M.C. Hunt, M.E. Dikeman, L.R. Corah, C.L. Kastner and D.H. Kropf. 1982. Ralgro implanted bulls: performance, carcass, characteristics and longissimus palatability. *Internatl. Symp. Meat Science and Technol.*, Lincoln, Nebraska, Nov. 1982.
- Hunt, M.C., P.S. Sleper, D.H. Kropf, C.L. Kastner and M.E. Dikeman. 1982. Electrical stimulation effects on myoglobin characteristics. *Internatl. Symp. Meat Science and Technol.*, Lincoln, Nebraska, Nov. 1982.
- Fung, D.Y.C., Fung, C-Y Lee, C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1982. Microbiology of hot-boned beef. *Internatl. Symp. Meat Science and Technol.*, Lincoln, Nebraska, Nov. 1982.
- Shivas, S.D., C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1983. Effects of electrical stimulation, hot boning and chilling rate on bull semimembranosus muscle. *J. Anim. Sci.* 57 (Suppl. 1):231.
- Choi, Y.I., C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1983. Effects of electrical stimulation and hot boning on functional characteristics of preblended beef muscle in model system. *Proc. Recip. Meat Conf.* 36:190.
- Unruh, J.A., C.L. Kastner, D.H. Kropf, M.E. Dikeman and M.C. Hunt. 1984. Effects of low voltage electrical stimulation during exsanguination on characteristics of beef longissimus and semimembranosus muscles. *J. Anim. Sci.* 59 (Suppl. 1):90.
- Kropf, D.H., C.M. Jaunsolo, M.E. Dikeman, M.C. Hunt and C.L. Kastner. 1984. Display color stability of vacuum packaged rib steaks from bulls and zeranol implanted bulls and steers. *J. Anim. Sci.* 59 (Suppl. 1):234.
- Hunt, M.C., C.M. Jaunsolo, D.H. Kropf, C.L. Kastner and M.E. Dikeman. 1984. Correlation of reflectance measurements to visual color for frozen and unfrozen skin-tight packaged beef longissimus. *J. Anim. Sci.* 59 (Suppl. 1):234.
- Unruh, J.A., C.L. Kastner, D.H. Kropf, M.E. Dikeman and M.C. Hunt. 1984. Effect of low voltage electrical stimulation during exsanguination on characteristics of beef longissimus and semimembranosus muscles. *Proc. Recip. Meat Conf.* 37:180.
- Burson, D.E. and M.C. Hunt. 1985. Heat-induced changes on the proportion of types I and III collagens in bovine longissimus. *J. Anim. Sci.* 61 (Suppl. 1):94.
- Hunt, M.C. and D.E. Burson. 1985. Proportion of types I and III collagens in four bovine muscles. *Ann. Meeting of IFT, Atlanta, GA (Abstr.)* #340.
- Burson, D.E., M.C. Hunt, J.A. Unruh and M.E. Dikeman. 1985. Proportions of types I and III collagen in intramuscular collagen of bulls and steers. *J. Anim. Sci.* 61 (Suppl. 1):94.
- Dikeman, M.E., T. Timmis, M.C. Hunt, R.H. Hines, G. Highfill and J.S. Stevenson. 1985. Effects of Compudose 2000 implants on performance, carcass, meat quality traits and serum testosterone in young boars. *J. Anim. Sci.* 61 (Suppl. 1):279.

- Hunt, M.C., M.E. Dikeman, G.A. Highfill, R.H. Hines and T. Timmis. 1985. Effects of sex and Compudose 2000 implantation on porcine longissimus and semimembranosus histochemistry. *J. Anim. Sci.* 61 (Suppl. 1):282.
- Kluber, III, E.F., J.S. Stevenson, D.L. Davis, M.C. Hunt, J.E. Minton and J.L. Nelssen. 1986. Growth, carcass, sexual and boar odor traits of growing male pigs fed altrenogest. *J. Anim. Sci.* 63(Suppl. 1):85.
- Allen, D.M., A. Luchiari Filho and M.C. Hunt. 1986. Effects of spray versus conventional chilling of beef carcasses and cut yields and USDA grade factors. *J. Anim. Sci.* 63(Suppl. 1):254.
- Claus, J.R., S.J. Goll, M.C. Hunt, G.A. Whipple and E.S. Wright. 1987. Meat Science at Kansas State University. *Proc. Recip. Meat Conf. Teaching Display.*
- Johnson, R.D., M.C. Hunt, D.M. Allen, C.L. Kastner and C.C. Schrock. 1988. Effects of carcass washing and length of spray cycle during chilling on surface fat and lean moisture contents on weight yields of holstein and beef steer carcasses. *J. Anim. Sci.* 66 (Suppl. 1):124.
- Goll, S.J., C.L. Kastner, M.C. Hunt and D.H. Kropf. 1988. Effects of glucose and internal cooking temperature on the characteristics of low fat, pre- and post-rigor, restructured beef roasts. *J. Anim. Sci.* 66 (Suppl. 1):126.
- Claus, J.R. and M.C. Hunt. 1988. Processed meats with reduced caloric density. *Kansas State University Cattlemen's Day.*
- Claus, J.R. and M.C. Hunt. 1988. Low-fat processed meats. *Kansas Agric. Exp. Conference.*
- Hunt, M.C. 1988. Spray chilling of beef carcasses. *Kansas Agric. Exp. Conference.*
- Goll, S.J., C.L. Kastner, M.C. Hunt and D.H. Kropf. 1988. Value added, restructured beef products. *Kansas Agric. Exp. Conference.*
- Whipple, G., M.C. Hunt, R.D. Klemm, D.H. Kropf, R.D. Goodband, J.L. Nelssen, R.H. Hines and B.R. Schricker. 1989. Effects of porcine somatotropin and supplemental lysine on porcine muscle histochemistry. *J. Anim. Sci.* 67(Suppl. 1):161.
- Whipple G., M. Koohmaraie, M.E. Dikeman, J.D. Crouse, M.C. Hunt and R.D. Klemm. 1989. Evaluation of attributes that affect longissimus muscle tenderness in *Bos taurus* and *Bos indicus* cattle. *Proc. Recip. Meat Conf.* 42:
- Nold, R.A., J.A. Unruh, C.W. Spaeth, J.E. Minton and M.C. Hunt. 1990. Effect of implanting zeranol at birth and weaning on performance, pelting, carcass and subprimal yield characteristics of ram and wether lambs. *J. Anim. Sci.* 68(Suppl. 1):93.
- Kenney, P.B. and M.C. Hunt. 1990. Effect of water and salt content on protein solubility and water retention of meat preblends. *J. Anim. Sci.* 68(Suppl. 1):61.
- Hunt, M.C., E.S. Troutt, D.E. Johnson, J.R. Claus, C.L. Kastner, D.H. Kropf and S. Stroda. 1990. Low-fat ground beef: physical, chemical and sensory characteristics. *J. Anim. Sci.* 68(Suppl. 1):327.
- Hunt, M.C. and J.R. Claus. 1990. Effect of texture modifying ingredients on characteristics of low-fat, high-added water bologna. *J. Anim. Sci.* 68(Suppl. 1):350
- Warren, K.E., M.C. Hunt, C.L. Marksberry, O. Sorheim and D.H. Kropf. 1990. Modified atmosphere packaging with 100% carbon dioxide for bone-in pork loins. *J. Anim. Sci.* 68(Suppl. 1):351
- Wang, H., D.H. Kropf, M.C. Hunt, and C.L. Kastner. 1990. Causes of iridescence in precooked beef. 49th Ann. Meeting of IFT, Dallas, TX (Abstr.)
- Nold, R.A., J.A. Unruh, C.W. Spaeth, J.E. Minton and M.C. Hunt. 1991. Effect of implanting ram and wether lambs with zeranol at birth and weaning on palatability and muscle collagen characteristics. *J. Anim. Sci.* 69(Suppl. 1):63.

- Zepeda, Garcia C.M., C.L. Kastner, D.H. Kropf, M.C. Hunt, P.B. Kenney, J.R. Schwenke, and D.S. Schluesener. 1991. Utilization of surimi-like products from pork with sex odor in restructured, precooked pork roasts. *J. Anim. Sci.* 69(Suppl. 1):97.
- Kropf, D.H., O. Sorheim, M.C. Hunt, K.E. Warren and M. Menninen. 1991. Modified atmosphere packaging and carcass chill effects on pork loins. *J. Anim. Sci.* 69(Suppl. 1):337.
- Brester, G.W., P. Lhermite, B.K. Goodwin and M.C. Hunt. 1992. Quantifying the effects of new product development: The case of low-fat ground beef. *Amer. Agric. Econ. Assoc. Annual Meeting*.(Abstr).
- Payne, C.A., M.C. Hunt, K.E. Warren, J.M. Hayden, J.E. Williams and H.B. Hedrick. 1992. Effects of nutrition on histochemical characteristics of four bovine muscles differing in growth impetus. *J. Anim. Sci.* 70(Suppl. 1):221.
- Warren, K.E., C.A. Payne, M.C. Hunt, G. Whipple, M. Koohmaraie, J. Hayden, J. Williams and H.B. Hedrick. 1992. Carcass composition, protease activity, shear force, and myofibrillar fragmentation index of longissimus thoracis muscle from crossbred steers as influenced by dry-matter intake. *J. Anim. Sci.* 70(Suppl. 1): 221.
- Hague, M.A., K.E. Warren, M.C. Hunt, D.H. Kropf and S.L. Stroda. 1992. Relationship between endpoint temperature, internal cooked color, and expressible juice color of ground beef patties made from imported trimmings and A- and D/E-maturity lean. *J. Anim. Sci.* 70(Suppl. 1):225.
- Payne, C.A., M.C. Hunt, K.E. Warren, J.M. Hayden, J.E. Williams and H.B. Hedrick. 1992. Histochemical properties of four bovine muscles as influenced by compensatory gain and growth impetus. *Proc. Internatl. Congress Meat Sci. and Technol.* 38:121.
- Kropf, D., H. Wang, M. Hunt and C. Kastner. 1992. Causes and solutions of iridescence in precooked meat. *Proc.*
- Hunt, M.C. and R.E. Campbell. 1992. Lean, fat, and connective tissue from beef shanks processed with a Baader *with a Baader*
- Warren, K.E., M.C. Hunt, M.A. Hague, D.H. Kropf, S.L. Stroda and D.E. Johnson. 1993. Inconsistencies in expressible juice and internal cooked color of ground beef patties. *J. Anim. Sci.* 71(Suppl. 1):50.
- Warren, K.E., M.C. Hunt, C.A. Payne, E.S. Troutt, A.E. Boyle, D.H. Kropf and D.E. Johnson. 1993. Investigation of the color stability of commercial pizza pepperoni during lighted display. *J. Anim. Sci.* 71(Suppl. 1):50.
- Campbell, R.E. and M.C. Hunt. 1993. Value-added processing of beef shanks by Baader desinewing. *J. Anim. Sci.* 71(Suppl. 1):51.
- Payne, C.A., M.C. Hunt and P.A. Seib. 1993. Effects of native and phosphate-modified starch on the yield and texture *yield and tex*
- Warren, K.E., M.C. Hunt, D.H. Kropf, S.J. Smith, and S.L. Stroda. 1993. Chemical Characterization of ground beef *of ground be*
- Osburn, W.N., J.T. Keeton and M.C. Hunt. 1993. Utilization of konjac flour in low-fat, precooked lamb sausages. *J. Anim. Sci.* 71(Suppl. 1):51.
- Esquivel, O., D.H. Kropf, M.C. Hunt and J.R. Schwenke. 1993. Effect of texturizers on low-fat fermented hard beef salami. *Ann. Meeting Inst. Food Tech.*
- Marksberry, C.L., D.H. Kropf, M.C. Hunt and C.L. Kastner. 1994. The effect of fat level, pH, and carcass maturity on the cooked internal color of ground beef patties at five end-point temperatures. *Ann. Meeting Inst. Food Tech.*
- Powell, T.H. and M.C. Hunt. 1994. Enzymatic collagen analysis determines heat-altered and solubilized collagen in cooked meat. *Ann. Meeting Inst. Food Tech.*
- Warren, K.E., M.C. Hunt, D.H. Kropf and C.L. Kastner. 1994. Chemical properties of ground beef patties exhibiting normal and premature brown internal cooked color. *Ann. Meeting Inst. Food Tech.*

- Warren, K.E., M.C. Hunt and D.H. Kropf. 1994. Myoglobin oxidative state affects internal cooked color development of ground beef patties. Ann. Meeting Inst. Food Tech.
- Payne, C.A., M.C. Hunt, P.A. Seib, D.H. Kropf and C.L. Kastner. 1994. Evaluation of commercial starches in a model meat batter system containing low-fat/high added water. Ann Meeting Inst. Food Tech.
- Marksberry, C.L., D.H. Kropf, M.C. Hunt and C.L. Kastner. 1994. The effect of fat level, pH, and carcass maturity of the cooked internal color of ground beef patties at five end-point temperatures. Ann. Meeting Inst. Food Tech.
- Payne, C.A., M.C. Hunt, P.A. Seib, D.H. Kropf and C.L. Kastner. 1994. Evaluation of commercial starches in a
- Warren, K.E., M.C. Hunt, D.H. Kropf and C.L. Kastner. 1994. Chemical properties of ground beef patties exhib
- Warren, K.E., M.C. Hunt and D.H. Kropf. 1994. Myoglobin oxidative state affects internal cooked color development of ground beef patties. Ann. Meeting Inst. Food Tech.
- Hunt, M.C., C.A. Payne and P.A. Seib. 1994. Properties of low-fat, high added-water beef sausages containing ph
- Hunt, M.C., K.E. Warren, D.H. Kropf, M.A. Hague, C.L. Waldner, S.L. Stroda, and C.L. Kastner. 1994. Factors affecting premature browning in cooked ground beef. Internatl. Cong. Meat Sci. & Technol. Netherlands.
- McCart, V.E., R.K. Phebus, D.Y.C. Fung, and M.C. Hunt. 1995. Effects of fat level and fat replacement in ground beef on the survival of *Listeria monocytogenes* and *Escherichia coli* O157:H7 during storage and cooking. Ann. Meeting Inst. Food Tech.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M. Hollingsworth, and C.L. Kastner. 1995. Palatability, color and product life of low-dose irradiated precooked ground beef. Proc. Internatl. Cong. Meat Sci. & Technol. 41:224-225.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M. Hollingsworth, and C.L. Kastner. 1995. Palatability, color and product life of low-dose irradiated beef steaks. Proc. Internatl. Cong. Meat Sci. & Technol. 41:272-273.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M. Hollingsworth, and C.L. Kastner. 1995. Palatability, color and product life of low-dose irradiated raw ground beef. Proc. Internatl. Cong. Meat Sci. & Technol. 41:278-279.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, E.J. Rubio Canas, M.C. Hunt, E. Chambers IV, M. Hollingsworth, and C.L. Kastner. 1995. Sensory analysis and consumer acceptance of low-dose irradiated boneless pork chops. Proc. Internatl. Cong. Meat Sci. & Technol. 41:416-417.
- Hunt, M.C., O. Sørheim, and E. Slinde. 1995. Effects of myoglobin form on internal cooked color development in ground beef. Proc. Internatl. Cong. Meat Sci. & Technol. 41:394-395.
- Lavelle, C.L., M.C. Hunt, and D.H. Kropf. 1995. Expressible juice and internal cooked color of ground beef patties from vitamin E-supplemented steers. Proc. Internatl. Cong. Meat Sci. & Technol. 41:396-397.
- Lavelle, C.L., M.C. Hunt and D.H. Kropf. 1995. Display life and internal cooked color of ground beef from vitamin E supplemented cattle. Ann. Meeting Inst. Food Tech.
- Hunt, M.C. and T.H. Powell. 1995. Physical properties of cooked meat--color and texture. Ann. Meeting Inst. Food Tech.
- Kropf, D.H., M.C. Hunt, O. Sorheim, K.E. Warren, C.L. Waldner, and F.W. Pohlman. 1995. Modified atmosphere packaging on the color of fresh and frozen muscle foods. Ann. Meeting Inst. Food Tech.
- Kropf, D.H., M.C. Hunt, S.E. Luchsinger, C.M. Garcia Zepeda, E. Chambers IV, J.L. Marsden, M.E. Hollingsworth, S.L. Stroda, E.J. Rubio Canas, and C.L. Kastner. 1995. Sensory analysis, color, and product

life of low-dose irradiated beef and pork. In Activities Report of the Research and Development Associates for Military Food and Packaging Systems. 48 (1): 327-345.

Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, E. Chambers IV, M.E. Hollingsworth, M.C. Hunt, J.L. Marsden, S.L. Stroda, E.J. Rubio Canas, C.L. Kastner, W.G. Kuecker, and T. Mata. 1996. Consumer acceptance of irradiated boneless pork chops. Abst. of Institute of Food Technologists Annual Meeting.

Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, E. Chambers IV, M.E. Hollingsworth, M.C. Hunt, J.L. Marsden, S.L. Stroda, E.J. Rubio Canas, C.L. Kastner, W.G. Kuecker, and T. Mata. 1996. Sensory analysis, color, and product life of irradiated boneless pork chops. Abst. of Institute of Food Technologists Annual Meeting.

Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M.E. Hollingsworth, and C.L. Kastner. 1996. Sensory analysis, color, and product life of irradiated frozen raw ground beef patties. Abst. of Institute of Food Technologists Annual Meeting.

Luchsinger, S.L., D.H. Kropf, and M.C. Hunt. 1996. Irradiation studies on meat. Proc. Res. on Salmonellosis in the Food Safety Consortium. U.S. Animal Health Assoc., Little Rock, AR.

Venkat, V., M.C. Hunt, and D.H. Kropf. 1997. Photographic guidelines for pork display color. Proc. Reciprocal Meat Conf. 50:167.

Henderson, L. and M.C. Hunt. 1997. Multimedia distance learning. Proc. Reciprocal Meat Conf. 50.

Kropf, D.H. and M.C. Hunt. 1998. Endpoint cooking temperature and meat color. Proc. Reciprocal Meat Conf. 51:144-148.

Clark, T.J., M.C. Hunt, D.H. Kropf, and G.R. DelDuca. 1998. Rebloom and display color stability of beef packaged in and ultra-low oxygen modified atmosphere system. Proc. Reciprocal Meat Conf. 51:185.

Schoenbeck, J.J., M.C. Hunt, T.E. Dobbels, M.E. Dikeman, and S.L. Stroda. 1998. Display color stability of steaks and ground beef from carcasses cardiovascularly infused immediately after exsanguination. Proc. Reciprocal Meat Conf. 51:186.

Killinger, K.M., M.C. Hunt, and R.E. Campbell. 1998. Incidence of premature browning in ground beef patties purchased at retail. Proc. Reciprocal Meat Conf. 51:185.

Schoenbeck, M.K., D.H. Kropf, M.C. Hunt, J.S. Pontius, S.L. Stroda, and S. Hawthorne. 1998. The effects of pH, myoglobin form and endpoint temperature on cooked ground beef color. Proc. Reciprocal Meat Conf. 51:186.

Venkat, V.R., M.C. Hunt, and D.H. Kropf. 1998. Photographic guidelines for pork display color stability. Ann. Meeting Inst. Food Tech.

Campbell, R.E. and M.C. Hunt. 1998. Sensory and physical characteristics of dry-aged beef strip steaks. Ann. Meeting Inst. Food Tech.

Hawthorne, S.A., R. Lien, M.C. Hunt, D.H. Kropf, and M. Hardin. 1999. Effects of endpoint temperature on internal cooked color development in ground pork and loin chops. Ann. Meeting Inst. Food Tech.

Hunt, M.C., S.A. Hawthorne, R. Lien, D.H. Kropf, and M. Hardin. 1999. Factors affecting the denaturation of myoglobin in ground pork patties. Proc. 45<sup>th</sup> Internatl. Congress Meat Sci. and Technol.

Dikeman, M.E., T.E. Dobbels, J.J. Schoenbeck, and M.C. Hunt. 1999. Effects of vascular infusion of cattle after exsanguination with a solution of saccharides, sodium chloride, and phosphates or with calcium chloride on carcass traits and meat palatability. J. Anim. Sci. 77 (Suppl. 1):77:23.

Yancey, E.J., M.C. Hunt, M.E. Dikeman, T.E. Dobbels, and P.B. Addis. 1999. Effects of vascular infusion of cattle after exsanguination with saccharides, sodium chloride, and phosphates plus vitamin C, E, or C+E on meat display color stability. J. Anim. Sci. 77 (Suppl. 1):77:172.

- Hachmeister, K.A., D.H. Kropf, J.L. Marsden, V.S. Gill, C.L. Kastner, and M.C. Hunt. 1999. Sensory analysis of electron beam pasteurized ground beef patties. Proc. New Mexico Environmental Health Conf. Albuquerque.
- Dikeman, M.E., T.E. Dobbels, M.C. Hunt, and J.J. Schoenbeck. 1999. Effects of post-bleeding vascular injection of cattle with a solution of sugars, sodium chloride, and phosphates or calcium chloride on carcass traits and meat palatability. Kansas Agr. Ep. Sta. Rep. of Prog. No 831:17-20.
- Hachmeister, K.A., D.H. Kropf, J.L. Marsden, V.S. Gill, C.L. Kastner, M.C. Hunt, and R.J. Kaye. 1999. Thiamin and riboflavin retention of electron beam pasteurized ground beef patties. Proc. Ann. Meeting Food Safety Consortium. p. 167.
- Hachmeister, K.A., D.H. Kropf, J.L. Marsden, V.S. Gill, R.J. Kaye, C.L. Kastner, and M.C. Hunt. 1999. Effects of repetitive high energy pulsed power (RHEPP) irradiation on sensory attributes, color, and shelf life of ground beef. Kansas Agr. Ep. Sta. Rep. of Prog. No 831:1-3.
- Hawthorne, S.A., R. Lein, M.C. Hunt, and D.H. Kropf. 1999. Ground pork cooked color guide. Kan. Ag. Exp Stat.
- Hawthorne, S.A., R. Lein, M.C. Hunt, and D.H. Kropf. 1999. Pork chop cooked color guide. Kan. Ag. Exp Stat.
- Hawthorne, S.A., M.C. Hunt, D.H. Kropf, and R. Lien. 1999. Effects of endpoint temperature on internal cooked color development in ground pork and loin chops. Proc. Ann Meeting Food Safety Consortium. p. 187.
- Slinde, Erik, Melvin Hunt and Oddvin Sorheim. 2000. Effects of myoglobin from on colour and heat denaturation of ground beef. Proc. 35<sup>th</sup> Norwegian Biochemical Contact Meeting. Tromso Norway. Abstract no. 4.
- Ceylan, E., D.Y.C. Fung, M.C. Hunt and C.L. Kastner. 2000. Synergistic effects of garlic and heating in controlling *Escherichia coli* 0157:H7 in ground beef patties. Ann. Meeting Inst. Food Tech.
- Ceylan, E., D.Y.C. Fung, M.C. Hunt and C.L. Kastner. 2000. Synergistic effects of garlic and heating in controlling *Escherichia coli* 0157:H7 in ground beef patties. Proc. Ann. Meeting Food Safety Consortium. p.
- Kropf, D.H. and M.C. Hunt. 2000 Shelf-life attributes of retail meat and poultry. Symposium: Factors influencing the Successful Implementation of Centralized Packaging of Meat and Poultry. Inst. Food Technologist Annual Meeting, Dallas TX, June 10-14.
- Lehmuller, Peter and Melvin Hunt. 2000 Perception of degrees of doneness of beef by professional chefs. Proc Reciprocal Meat Conf. 53:133.



Conference Proceedings, Symposia, Books, etc.

- Kropf, D.H., H.J. Tuma, M.C. Hunt, R.A. Smith, M.L. Sandberg, J.A. Santamaria, J.A. Fry and D.E. Schafer. 1971. Retail display case lighting. Proc. 12th Annual Meeting of Food Distribution Research Society.
- Kropf, D.H., H.J. Tuma, C.C. Allen, M.C. Hunt, M.L. Sandberg and D.E. Schafer. November, 1974. Evaluation of color and other properties of frozen beef. Proc. of Symposium on Objective Methods for Food Evaluation (Sponsored by U.S. Army Natick Labs and National Research Council).
- Kropf, D.H., M.C. Hunt, C.C. Allen and H.J. Tuma. 1976. Color measurement of red meat. IFT Symposium on Color measurement in foods.
- Hunt, M.C. 1980. Meat color measurement. Proc. Recip. Meat Conf. 33:41.
- Kastner, C.L., M.E. Dikeman, K.N. Nagele, M. Lyon, M.C. Hunt and D.H. Kropf. 1980. Effects of carcass electrical stimulation and hot boning on selected beef muscles. Proc. European Meeting of Meat Research Workers. Vol. 2. 26:40.
- Hunt, Melvin C. 1981. Conversion of muscle to meat. Proc. 2nd Internatl. Course of Meat Technol. Inst. de Tecnologia de Alimentos, Campinas, SP-Brazil. p. 14-1.
- Hunt, Melvin C. 1981. Meat emulsions. Proc. 2nd. Internatl. Course of Meat Technol. Inst. de Tecnologia de Alimentos, Campinas, SP-Brazil. p. 23-1.
- Hunt, Melvin C. 1981. Nitrite and chemistry of cured color. Proc. 2nd Internatl. Course of Meat Technol. Inst. de Tecnologia de Alimentos, Campinas, SP-Brazil. p. 24-1.
- Hunt, Melvin C. 1981. Tumbling, massaging and restructured meat. Proc. 2nd Internatl. Course of Meat Technol. Inst. de Tecnologia de Alimentos, Campinas, SP-Brazil. p. 26-1.
- Kropf, D.H., C. Kuntapanit, M.C. Hunt, C.L. Kastner and B.E. Brent. 1984. Effect of vacuum aging and display on lipid oxidation of subcutaneous fat and three layers of beef longissimus. Proc. European Meat Res. Workers. 30:221.
- Kropf, D.H. and M.C. Hunt. 1984. Effect of display conditions on meat products. Proc. Meat Ind. Res. Conf. p. 153.
- Hunt, M.C. and D.H. Kropf. 1985. Fresh and cured meat color analyses. IFT Muscle Foods Symposium.
- Kropf, D.H., M.C. Hunt and Dorly Piske. 1985. Color formation and retention in fresh meat. Proc. Meat Ind. Res. Conf. p. 62.
- Hunt, M.C. and D.H. Kropf. 1987. Color and Appearance. In: Restructured Meat and Poultry Products. Adv. in Meat Research. Vol. 3, p. 125. AVI Publishing Co., Inc., Westport, CT.
- Bjerklie, S. and M.C. Hunt. 1988. Practical meat photography. Proc. Recip. Meat Conf. 41:
- Hunt, M.C., G.A. Highfill, M.E. Dikeman and R.H. Hines. 1988. Sex and compudose® implantation effects on porcine longissimus and semimembranous fiber types. Proc. Internatl. Congress Meat Sci. and Technology. 34:62.
- Hunt, M.C. 1989. Career opportunities in meat/food science. Southern Univ. Shreveport, MS. Live, nationwide satellite tele-video presentation. Co-sponsored by Campbells Soup Co.
- Hunt, M.C., J.C. Acton, R.C. Benedict, C.R. Calkins, D.P. Cornforth, L.E. Jeremiah, D.G. Olson, C.P. Salm, J.W. Savell and S.D. Shivas. 1991. Guidelines for Meat Color Evaluation. Proc. Recip. Meat Conf. 44:232.
- Payne, C.A., M.C. Hunt, K.E. Warren, J.M. Hayden, J.E. Williams and H.B. Hedrick. 1992. Histochemical properties of four bovine muscles as influenced by compensatory gain and growth impetus. Proc. Internatl. Congress Meat Sci. and Technol. 38:121.

- Kropf, D., H. Wang, M. Hunt and C. Kastner. 1992. Causes and solutions of iridescence in precooked meat. *Proc. Internatl. Congress Meat Sci. and Technol.* 38: 515.
- Hunt, M.C. and R.E. Campbell. 1992. Lean, fat, and connective tissue from beef shanks processed with a Baader desinewer. *Proc. Internatl. Congress Meat Sci. and Technol.* 38:1219.
- Payne, T. and M. Hunt. 1993. Starch use in meat products. *Proc. Midwest Meat Proc. Seminar.* 22:I-1.
- Hunt, M.C. and D.H. Kropf. 1993. Color measurement of meat and meat products. *Proc. Recip. Meat Conf.* 46:59.
- Hunt, M.C. and R. E. Campbell. 1993. Spray chilling of beef and pork carcasses for weight conservation. *Proc. Meat Ind. Res. Conf.*
- Hunt, M.C. and T.H. Powell. 1995. Physical properties of cooked meat - color and texture. *Muscle Foods Symposium. Ann. Meeting Inst Food Tech.*
- Kropf, D.H., M.C. Hunt, K.E. Warren, M.A. Hague, C.L. Waldner, S.L. Stroda, and C.L. Kastner. 1995. Cooked ground beef color is unreliable indicator of maximum internal temperature. *Amer. Chem Society Ann. Meeting, Anaheim, CA.*
- Lavelle, C., M.C. Hunt, and D.H. Kropf. 1995. Cooked ground beef color and safety. *Proc. Meat Industry Res Conf. Amer. Meat Inst.* p. 167.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, E. Chambers IV, M.E. Hollingsworth, M.C. Hunt, J.L. Marsden, S.L. Stroda, E.J. Rubio Canas, C.L. Kastner, W.G. Kuecker, and T. Mata. 1995. Consumer acceptance of low-dose irradiated boneless pork chops. In *Annual Report, Swine Day, Kansas State University*, p. 129-130.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, E. Chambers IV, M.E. Hollingsworth, M.C. Hunt, J.L. Marsden, S.L. Stroda, E.J. Rubio Canas, C.L. Kastner, W.G. Kuecker, and T. Mata. 1995. Flavor and aroma of low-dose irradiated boneless pork chops. In *Annual Report, Swine Day, Kansas State University*, p. 131-134.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, E. Chambers IV, M.E. Hollingsworth, M.C. Hunt, J.L. Marsden, S.L. Stroda, E.J. Rubio Canas, C.L. Kastner, W.G. Kuecker, and T. Mata. 1995. Display life and related traits of low-dose irradiated boneless pork chops. In *Annual Report, Swine Day, Kansas State University*, p. 135-138.
- Luchsinger, S.E., D.H. Kropf, and M.C. Hunt. 1996. Irradiation studies on meat. *Proc. Symp. Studies on Food Borne Salmonellosis. Food Safety Consortium and U.S. Public Health Assoc.* Little Rock, AK.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M.E. Hollingsworth, and C.L. Kastner. 1996. Sensory traits, color, and shelf life of irradiated beef steaks. In *Annual Report, Cattleman's Day, Kansas State University*, p. 4-7.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M.E. Hollingsworth, and C.L. Kastner. 1996. Sensory traits, color, and shelf life of irradiated raw ground beef patties. In *Annual Report, Cattleman's Day, Kansas State University*, p. 8-10.
- Luchsinger, S.E., D.H. Kropf, C.M. Garcia Zepeda, J.L. Marsden, S.L. Stroda, M.C. Hunt, E. Chambers IV, M.E. Hollingsworth, and C.L. Kastner. 1996. Sensory traits, color, and shelf life of irradiated precooked ground beef patties. In *Annual Report, Cattleman's Day, Kansas State University*, p. 11-13.
- Kropf, D. H., S. E. Luchsinger, C.M. Garcia-Zepeda, E. Chambers, IV., and M.C. Hunt. 1996. Irradiation: A path to a safer food supply. *Proc. Meat Industry Res. Conf. Amer. Meat Inst.*
- Venkat, V., M.C. Hunt, and D.H. Kropf. 1997. Photographic guidelines for pork display color. *Proc. Reciprocal Meat Conf.* 50:

Luchsinger, J.E., M.C. Hunt, and D.H. Kropf. 1997. High carbon dioxide modified-atmosphere packaging (MAP) for beef steaks. Kans. Ag. Exp. Stat. Rpt. of Progress 783.

Powell, T.H., M.E. Dikeman, and M.C. Hunt. 1999. Modeled, multi-stage convection cooking of beef semitendinosus roasts to denature collagen and to optimize tenderness. Kansas Agr. Expt. Sta. Rpt. Prog. No. 831:34-36.

Hunt, M.C., J.J. Schoenbeck, M.E. Dikeman, T.E. Dobbels, and S.L. Stroda. 1999. Color stability of steaks from carcasses vascularly infused immediately after exsanguination. Kansas Agr. Expt. Sta. Rep. of Prog. No. 831:21-25.

Hunt, M.C., K.M. Killinger, and R.E. Campbell. 1999. Incidence of premature browning during cooking in ground beef purchased at retail. Kansas Agr. Expt. Sta. Rpt. of Prog No. 831:11-13.

Hachmeister, K.A. D.H. Kropf, J.L. Marsden, V.S. Gill, C.L. Kastner, M.C. Hunt, and R.J. Kaye. 2000. Thiamin and riboflavin retention in ground beef patties pasteurized by electron beam. KAES Rpt. of Progress 850. Kansas State University, Manhattan.

Schoenbeck, M.K., D.H. Kropf, M.C. Hunt, S.L. Stroda, and S. Hawthorne. 2000. Effects of pH, myoglobin form, and endpoint temperature on cooked ground beef color. KAES Rpt of Progress 850. Kansas State University, Manhattan.

#### Bulletins, Seminars, Short Courses, Other Presentations

Kropf, D., H.J. Tuma, M. Sandberg, J. Santamaria, D. Schafer, M. Hunt and R. Smith. November, 1970. Effects of film permeability, freezing system, packaging time and display case temperature on frozen beef steaks. Study II. Effect of display light intensity on frozen beef steaks. Report to: E.I. DuPont De Nemours & Co.

Allen, D.M., M.C. Hunt, C.L. Kastner and D.H. Kropf. 1976. Kansas State University ResearchUp-Date. Improving systems of marketing red meat. Proc. Midwest Meat Processor's Seminar, Kansas State University. p. 3.

Allen, D.M., M.C. Hunt, C.L. Kastner, D.H. Kropf, V. Chen, A. Harrison, O. Corte, C. Kuntapanit, M.E. Smith and J. Thomas. 1976. Method of improving quality of grass-fed beef. Kansas State Univ. Cattlemen's Day Report, No. 262. p. 79.

Allen, D.M., M.C. Hunt, C.L. Kastner, D.H. Kropf, G.H. Gutowski, A.R. Harrison and M.E. Smith. 1977. Characteristics of beef finished on selected feeding regimes. Kansas State Univ. Cattlemen's Day Rept. p. 105.

Hunt, M.C. 1978. Effect of feeding regime and time on feed on beef quality and yield. Proc. Midwest Meat Processors Seminar. p. E-1.

Allen, D.M., M.C. Hunt, C.L. Kastner, D.H. Kropf, A.R. Harrison, B.E. Brent and J. Riley. 1978. Effects on carcass traits of beef ration, energy level, and length of feeding. Kansas State Univ. Cattlemen's Day Report 320. p. 86.

Allen, D.M., M.C. Hunt, C.L. Kastner, D.H. Kropf, V. Chen, O. Corte, G.H. Gutowski, A.R. Harrison, C. Kuntapanit, J.D. Thomas and M.E. (Smith) McCurry. 1978. Carcass characteristics, palatability, and shelf life of beef finished on selected feeding regimens. Kansas State Univ. Cattlemen's Day Report 320. p. 79.

Hunt, M.C. 1979. Mechanical Blade Tenderization. Proc. MidWest Meat Processors Seminar. p. H-1.

Burson, D.E., M.C. Hunt, L.H. Hayward, C.L. Kastner, D.H. Kropf and D.M. Allen. 1979. Nutritional effects on beef palatability. Kansas State Univ. Cattlemen's Day Report 350. p. 65.

Burson, D.E., L.H. Hayward, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1979. Mechanical tenderization of meat. Kansas State Univ. Cattlemen's Day Report 350. p. 68.

- Fung, D.Y.C., C. Kastner, M. Hunt, M. Dikeman and D. Kropf. 1979. Mesophilic and psychrotrophic populations of hot-boned and conventionally beef. Presented at Institute of Food Technologists Annual Meeting.
- Harrison, A.R., D.H. Kropf, D.M. Allen, M.C. Hunt and C.L. Kastner. 1979. Relationship of various spectrophotometric reflectance measurements to beef muscle visual color. Ann. Meeting IFT.
- Erickson, D.B., J.H. McCoy, J.B. Riley, D.S. Chung, P.G. Nason, D.M. Allen, M.E. Dikeman, D.Y.C. Fung, M.C. Hunt, C.L. Kastner and D.H. Kropf. 1980. Hot processing: economic feasibility of hot processing beef carcasses. Bulletin 639. Kansas Agricultural Experiment Station, Manhattan.
- Hunt, M.C., J.L.A. Kendall, M.E. Dikeman, C.L. Kastner and D.H. Kropf. 1980. Ground beef from electrically stimulated and pre-rigor processed carcasses. Kansas State Univ. Cattlemen's Day Report, No. 377. p. 12.
- McCoy, J., P. Nason, D. Chung, C. Kastner, A. Lawrence, M. Dikeman, M. Hunt and D. Kropf. 1980. Hot processing--potential for application in the beef processing industry. Study I: Economic feasibility of hot processing beef carcasses. Kansas State Univ. Cattlemen's Day Report, No. 377. p. 3.
- Nagele, K., M.E. Dikeman, M.C. Hunt, C.L. Kastner, D.H. Kropf and M. Lyon. 1980. Hot processing--potential for application in the beef processing industry. Study II. Electrically stimulated and hot-processed beef--color and eating qualities. Kansas State Univ. Cattlemen's Day Report, No. 377. p. 5.
- Fung, D.Y.C., C-Y. Lee, C. Kastner, M. Dikeman, M. Hunt, D. Kropf and M. Lyon. 1980. Hot processing--potential for application in the beef processing industry. Study III: Hot processed beef--microbiological characteristics. Kansas State Univ. Cattlemen's Day Report, No. 377. p. 7.
- Fung, D.Y.C., C.L. Kastner, M.C. Hunt, M.E. Dikeman and D.H. Kropf. 1980. Recent developments in microbiological studies on hot-boned beef. Proc. Internatl. Symp. on Recent Advances in Food Sci. and Technol., Jan. 9-11, Taipei, Rep. of China. Vol. II. p. 341.
- Hunt, M.C. 1980. Renovation of KSU meat laboratory. Proc. Midwest Meat Processors Seminar.
- Hunt, M.C. 1980. Energy use in the meat industry. KSU Energy Symposium.
- Wu, J.J., C.L. Kastner, M.C. Hunt, D.H. Kropf and D.M. Allen. 1981. Nutritional effects of beef connective tissue characteristics and eating qualities. Kansas State Univ. Cattlemen's Day Report, No. 394. p. 19.
- Hunt, M.C. 1981. Potential beef carcass grade changes. Presentation at Cattlemen's Day.
- Hunt, M.C. 1981. Tumbling and massaging. Presentation at KSU Curing School.
- Hunt, M.C. 1981. Packaging of cured meat. Presentation at KSU Curing School.
- Bowles, J.E., C.L. Kastner, M.E. Dikeman, M.C. Hunt, J.L.A. Kendall and M. Lyon. 1981. Continuous versus intermittent electrical stimulation of beef carcasses and their effect on hot-boned muscle pH decline. Kansas State Univ. Cattlemen's Day Report No. 394, p. 25.
- Greathouse, J.R., M.C. Hunt, D.E. Schafer and D.H. Kropf. 1981. Vacuum packaging of frozen bacon. Proc. Midwest Meat Processors Seminar. p. G-1.
- Bowles, J.E., C.L. Kastner, M.E. Dikeman, M.C. Hunt, J.L.A. Kendall and M. Lyon. 1981. Continuous versus intermittent electrical stimulation of beef carcasses and their effect on hot-boned muscle pH decline. Kansas State Univ. Cattlemen's Day Report, No. 394. p. 25.
- Greathouse, J.R., M.C. Hunt, M.E. Dikeman, L.R. Corah, C.L. Kastner and R. J. Pruitt. 1982. Effects of ralgro implants on growth, sexual development, carcass characteristics and eating quality of bulls implanted from birth to slaughter. Kansas State Univ. Cattlemen's Day Report No. 413. p. 60.
- Hunt, M.C. 1982. Vacuum packaged ground beef and other fresh retail cuts. Proc. Midwest Meat Processors Seminar. p. F-1.

- Burson, D.E., M.C. Hunt, D.E. Schafer, D. Beckwith and J.R. Garrison. 1982. Stunning method and time interval from stunning to bleeding effects on blood splashing in pork. Kansas State Univ. Swine Day Report No. 422, p. 122.
- Greathouse, J.R., M.C. Hunt, D.E. Schafer, D.H. Kropf and J.L. A. Kendall. 1982. Packaging effects on storage of frozen bacon. Meat Plant Magazine. December p. 18.
- Shivas, S.D., C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1983. Eating and cooking loss characteristics of electrically stimulated and hot boned bull inside round muscle chilled at different rates. Kansas State Univ. Cattlemen's Day Report No. 427, p. 69.
- Choi, Y.I., C.L. Kastner, M.E. Dikeman, M.C. Hunt and D.H. Kropf. 1983. Effects of electrical stimulation and hot boning on the functional characteristics of presalted beef muscle used in sausage manufacturing. Kansas State Univ. Cattlemen's Day Report No. 427, p. 74.
- Shivas, S.D., D.H. Kropf, C.L. Kastner, M.C. Hunt, J.L.A. Kendall and A.D. Dayton. 1983. Ascorbic acid and ground beef display life. Kansas State Univ. Cattlemen's Day Report No. 427, p. 74.
- Hunt, M.C. 1983. Factors affecting meat tenderness. Proc. Midwest Meat Processors Seminar. B-1.
- Kropf, D.H., M.E. Dikeman, M.C. Hunt and H.R. Cross. 1983. Lighting effects on beef carcass grade factors. Kansas State Univ. Cattlemen's Day Report No. 427, p. 76.
- Timmis, T., G. Highfill, M.E. Dikeman, M.C. Hunt and R.H. Hines. 1983. Effect of Compudose on elimination of boar odor. Kansas State Univ. Swine Day Report No. 442, p. 115.
- Hunt, M.C. 1984. Making sausage meats work for you. Proc. Kansas-Nebraska Sausage School.
- Axe, J.B., C.L. Kastner, M.E. Dikeman, M.E. Hunt, D.H. Kropf and D.G. Gray. 1984. Effects of low voltage electrical stimulation during bleeding and hot boning on beef loin eye and top round muscles. Kansas State Univ. Cattlemen's Day Report No. 448, p. 1.
- Unruh, J.A., C.L. Kastner, D.H. Kropf, M.E. Dikeman and M.C. Hunt. 1984. Effects of low voltage electrical stimulation during bleeding on characteristics of beef loin eye and top round muscles. Kansas State Univ. Cattlemen's Day Report No. 448, p. 6.
- Timmis, T., M.E. Dikeman, M.C. Hunt, R.H. Hines, G. Highfill and J.S. Stevenson. 1984. Effects of compudose implants on performance, carcass, meat quality traits and serum testosterone in young boars. Kansas State Univ. Swine Day Report No. 461, p. 92.
- Kropf, D.H. and M.C. Hunt. 1985. Packaging and Meat Color. MSU Packaging Symposium.
- Kluber, III, E.F., D.S. Pollmann, D.L. Davis, J.S. Stevenson, M.C. Hunt, J.E. Minton and J.L. Nelssen. 1985. Body growth and testicular characteristics of boars fed a synthetic progestogen, altrenogest. Kansas State Univ. Swine Day Report No. 486, p. 112.
- Flores, H.A., C.L. Kastner, D.H. Kropf and M.C. Hunt. 1986. Methods of tenderization for value-added, hot-boned, restructured, pre-cooked roasts from cows. Kansas State Univ. Cattlemen's Day Report No. 494, p. 88.
- Hunt, M.C. and D.E. Schafer. 1986. Diet, health, nutrition - some issues involving meat. Proc. Midwest Meat Processors Seminar. H-1.
- Kropf, D.H., M.C. Hunt and Dorly Piske. 1987. Color formation and retention in fresh beef. Kansas State Univ. Cattlemen's Day Report No. 514, p. 44.
- Todd, S.L., J.R. Claus, F.E. Cunningham and M.C. Hunt. 1987. Pork nuggets formulated with dietary fiber. Kansas State Univ. Swine Day Report No. 528, p. 114.

- Kropf, D.H., S. Hung and M.C. Hunt. 1987. Effect of display lighting on fresh pork longissimus packaged in oxygen-permeable and oxygen-barrier films. Kansas State Univ. Swine Day Report No. 528, p. 101.
- Goll, S.J., C.L. Kastner, M.C. Hunt and D.H. Kropf. 1988. Effects of sugar, internal cooking temperature, and hot-boning on the characteristics of low-fat, restructured, value-added beef roasts. Kansas State Univ. Cattlemen's Day Report No. 539, p. 81.
- Hunt, M.C. 1988. Cholesterol in meat. Kansas State Univ. Cattlemen's Day Report, No. 539.
- Claus, J.R., M.C. Hunt and C.L. Kastner. 1988. Effects of substituting added water for fat on the textural, sensory and processing characteristics of bologna. Proc. Midwest Meat Processors Seminar, p. C-1.
- Hunt, M.C., G. Whipple-Van Patter, D.H. Kropf, R.D. Klemm, R.D. Goodband, J.L. Nelssen, R.H. Hines and B.R. Schricker. 1989. Will porcine somatotropin (pST) lower pork quality? Kansas State Univ. Swine Day Rept. of Progress 581, p. 171.
- Kenney, P.B., C.L. Kastner, D.H. Kropf, M.C. Hunt and C. Garcia. 1989. Refining beef raw materials for further processing. Kansas State University Cattlemen's Day.
- Wang, H., D.H. Kropf, M.C. Hunt and C.L. Kastner. 1989. Iridescence in cooked beef. Kansas State University Cattlemen's Day.
- Dikeman, M.E., G. Whipple and M.C. Hunt. 1989. Characteristics responsible for tenderness differences in Bos taurus and Bos indicus cattle. Kansas Agric. Exp. Conference.
- Hunt, M.C., K.E. Warren and D. H. Kropf. 1990. Packaging of pork loins. Kansas State Univ. Swine Day.
- Hunt, M.C. 1990. Precooked deli meats. Proc. Missouri Meat Processors Assoc.
- Wang, H., D.H. Kropf, M.C. Hunt and C.L. Kastner. 1990. Effects of processing variables on iridescence in precooked beef. Kansas State Univ. Cattlemen's Day Rept. of Progress 592, p. 48.
- Whipple, G., M. Koohmaraie, M.E. Dikeman, J.D. Crouse, M.C. Hunt and R.D. Klemm. 1990. Evaluation of attributes affecting tenderness differences between Bos taurus and Bos indicus cattle. Kansas State Univ. Cattlemen's Day Rept. of Progress 592, p. 38.
- Hunt, M.C. 1990. Developing low-fat beef products. Cattlemen's Day Presentation.
- Warren, K.E., M.C. Hunt, C.L. Marksberry, O. Sørheim and D.H. Kropf. 1990. Bone-in pork loins: Modified atmosphere packaging to extend shelf life. Kansas State Univ. Swine Day Rept. of Progress 610, p. 108.
- Kropf, D.H., O. Sørheim, M.C. Hunt, M. Menninen and K.E. Warren. 1990. Effects of Modified atmosphere packaging and carcass chill rate on pork loins. Kansas State Univ. Swine Day Rept. of Progress 610, p. 112.
- Garcia Zepeda, C.M., C.L. Kastner, D.H. Kropf, M.C. Hunt, P.B. Kenney, J.R. Schwenke and D.S. Schleusener. 1990. Utilization of surimi-like products from pork with sex-odor in restructured, precooked pork roasts. Kansas State Univ. Swine Day Rept. of Progress 610, p. 115.
- Cunningham, F., D. Fung, M. Hunt, C. Kastner, D. Kropf, B. Larson, D. Schafer, D. Simms, S. Smith, and M. Vanik
- Nold, R.A., J.A. Unruh, C.W. Spaeth, and M.C. Hunt. 1991. The effects of implanting ram and wether lambs with zeranol at birth and weaning on pelt removal and pelt characteristics. Kansas State Univ. Sheep Day Rept. of Progress 624, p. 23.
- Nold, R.A., J.A. Unruh, C.W. Spaeth, and M.C. Hunt. 1991. The effects of implanting ram and wether lambs with zeranol at birth and weaning on palatability characteristics. Kansas State Univ. Sheep Day Rept. of Progress 624, p. 26.
- Hunt, M.C. 1991. Low-fat ground beef: Industry applications. Cattlemen's Day Presentation.

- Hunt, M.C. 1991. Complex carbohydrates, starches and dietary fiber in low-fat ground beef. National Low-fat Ground Beef Symposium--Baton Rouge, LA. National Live Stock and Meat Board.
- Hunt, M.C. 1991. Low-fat ground beef: Technology Perspectives. National Press Conference, Waldorf Astoria, New York, NY. Sponsored by Ketchum Public Relations and the National Live Stock and Meat Board.
- Hunt, M.C. 1991. Methods for production of low-fat ground beef. National Grocers Association Annual Meeting. Kansas City, MO.
- Hunt, M.C. 1991. Low-fat ground beef--Applications and practice. Meat Operations Conference. National-American Wholesale Grocers' Association. Kansas City, MO.
- Hunt, M.C. 1991. Development and manufacture of low-fat ground beef. Minnesota Grocers Association Annual Meeting. Minneapolis, MN.
- Hunt, M.C. 1991. Low-fat regional press conference for food writers. Sponsored by the Minnesota Beef Council. Minneapolis, MN.
- Hunt, M.C. 1992. Applications of low-fat meat technology to ground products. Proc. Kansas Midwest Meat Processors Conference.
- Hunt, M.C., E.S. Troutt, C.L. Kastner, D.H. Kropf and S. Stroda. 1991. Low-fat ground beef. Kansas Agric. Exp. Conference.
- Payne, C.A., K.E. Warren and M.C. Hunt. 1992. Nutritional effects on fiber characteristics of four bovine muscles varying in growth impetus. Kansas Agric. Exp. Conference.
- Warren, K.E., C.A. Payne and M.C. Hunt. 1992. Influence of dry matter intake on animal growth, carcass composition and longissimus muscle tenderness in crossbred steers. Kansas Agric. Exp. Conference.
- Campbell, R.E. and M.C. Hunt. 1992. Recovery of high quality lean and connective tissue from beef shanks. Kansas Agric. Exp. Conference.
- Hunt, Melvin. 1992. Developing meat products -- an activity for high school students. Kansas Ag. Educ. Mid-winter Conference.
- Hunt, M.C. and D.H. Kropf. 1992. Fresh and processed meat color chemistry and color assessment. W.R. Grace Conference.
- Hunt, Melvin. 1993. Relationship between meat packaging and meat color. Kansas Ag. Educ. Mid-winter Conference.
- Hunt, Melvin. 1993. New developments in beef processing. Cattlemen's Day Presentation.
- Hunt, Melvin. 1993. Advanced technologies for the meat industry. Food and Agribus. Industry of Kansas City.
- Carmack, C.F., C.L. Kastner, M.C. Hunt, D.H. Kropf and J.R. Schwenke. 1993. Can "natural" flavorings enhance the flavor of low-fat ground beef? Kansas State Univ. Cattlemen's Day Rpt. of Progress.
- Marksberry, C.L., D.H. Kropf, M.C. Hunt, M.A. Hague, and K.E. Warren. 1993. Ground beef patty cooked color guide. Kan. Ag. Exp. Stat.
- McCauley, W.H., D.H. Kropf, and M.C. Hunt. 1996. Cured meat color guide. Kan. Ag. Exp. Stat.
- Hunt, M.C. 1997. Food safety and cooked color of ground beef patties. USDA Workshop. Washington, D.C.
- Hawthorne, S.A., R. Lein, M.C. Hunt, and D.H. Kropf. 1999. Ground pork cooked color guide. Kan. Ag. Exp. Stat.
- Hawthorne, S.A., R. Lein, M.C. Hunt, and D.H. Kropf. 1999. Pork chop cooked color guide. Kan. Ag. Exp. Stat.
- Hunt, M.C. 1999. Distance learning program for Food Science at KSU. Provost Series on Education at KSU.

Hunt, M.C. 1999. Distance learning methods. KSU College of Agriculture Focus Series.

### Educational Materials

#### Course Syllabi:

- Distance Learning - Complete course, *PRINCIPLES OF MEAT SCIENCE* in "any-time or real-time" distance learning format via audiotapes, teleconferencing, on-line discussions involving collaborative learning, problem solving and critical thinking via the Internet.
- In-Class - Laboratory Materials for *MEAT SCIENCE*, a series of 14 study guides for lab exercises  
- Laboratory Materials for *PROCESSED MEAT OPERATIONS*

#### Other Materials and Activities:

USDA Grant: - Expanding Undergraduate Education for Food Industry Personnel via Technology.  
1994-96 USDA Challenge Grant Program, \$79,479

Web-based Course - Principles of Meat Science, KSU Division of Continuing Education

- Color Guides
- Ground Beef Patty Cooked Color Guide
  - Cured Meat Color Guide
  - Cooked Pork Chop Color Guide
  - Ground Pork Patty Cooked Color Guide

- Science Series
- Lesson Plans for: Promoting Ag Science for Secondary Schools
  - Developing New Meat Products
  - Color Chemistry in Meat Products
  - Meat Packaging Exercises for High School Students

- Slides Series:
- Unraveling the Mystery of Premature Browning in Cooked Ground Beef Patties
  - Doneness of Cooked Ground Beef
  - Dynamics of Conversion of Myoglobin Forms
  - Role of Pigment Layers in Influencing Surface Meat Color
  - Spray Chilling of Carcasses
  - Don't be Broken-Hearted because of High-fat in Ground Beef
  - Commercial Sausage, Ham and Bacon Production
  - Food Science at KSU
  - ASI Quadrathlon - why I should participate
  - Updated: Muscle-Bone Anatomy; Beef-Pork-Lamb Cut Identification

- Video Tapes:
- Beef Carcass Electrical Stimulation and Hot Boning  
(Edited with M. E. Dikeman)

- Store Survey:
- Out-of-class assignment for Analysis of Retail Meat Section of Grocery Stores

- Diet Survey:
- Out-of-class assignment for computerized class project of Nutritional Value of Muscle Foods in the student's diet

- Current topic:  
Survey
- Out-of-class assignment for critically analyzing printed literature on a variety of livestock and meat industry topics

- Web Sites:
- Out-of-class assignment for evaluation and collection of scientific facts about muscle biology and meat science

### Theses and Dissertations

Gutowski, G.H. 1977. Effect of vacuum aging, display and level of nutrition on beef quality. M.S. Thesis, Kansas State University.

Hayward, L.H. 1978. Blade tenderization effects on subjective and instron objective textural measurements of longissimus steaks from cattle fed various nutritional regimes. M.S. Thesis, Kansas State University.



- Burson, D.E. 1979. Ration energy density and time on feed effects on beef longissimus palatability. M.S. Thesis, Kansas State University.
- Hall, J.B. 1981. Collagen solubility of A-maturity bovine longissimus muscle as affected by nutritional regimen. M.S. Thesis, Kansas State University.
- Greathouse, J.R. 1982. Ralgro implanted bulls: Performance, carcass characteristics, longissimus palatability and carcass electrical stimulation. M.S. Thesis, Kansas State University.
- Sleper, P.S. 1982. Myoglobin properties of electrically stimulated bovine longissimus muscle. M.S. Thesis, Kansas State University.
- Highfill, G.A. 1984. Effects of sex and compudose implantation on porcine muscle histochemistry. M.S. Thesis, Kansas State University.
- Burson, D.E. 1985. Effects of muscle, heat and sex on the proportions of types I and III bovine intramuscular collagen. Ph.D. Dissertation. Kansas State University.
- Claus, J.R. 1989. Characterization of low-fat processed meat containing dietary fiber. Ph.D. Dissertation, Kansas State University.
- Troutt, S.E. 1990. A chemical, physical and sensory characterization of low-fat ground beef. M.S. Thesis, Kansas State University.
- Warren, K.E. 1990. Modified atmosphere packaging with 100% carbon dioxide for bone-in pork loins. M.S. Thesis, Kansas State University.
- Hague, M.A. 1992. Internal and expressible juice color of cooked ground beef patties. M.S. Thesis, Kansas State University.
- Londono Villegas, J.F. 1993. Effects of realimentation and trenbolone acetate implantation of cull cows on tenderness and cooked color. M.S. Thesis, Kansas State University.
- Payne, A.C. 1993. Starch functionality and modification for batter sausages. PhD. Dissertation, Kansas State University.
- Warren, K.E. 1994. Factors affecting premature browning of heated myoglobin. PhD. Dissertation, Kansas State University.
- Conner, J.G. 1995. Fat trimming and vitamin E effects on subprimal and retail cut yields and shelf life. M.S. Thesis, Kansas State University.
- Schoenbeck, J.J. 1997. Effects of carcass infusion on color and display life of several bovine muscles. M.S. Thesis, Kansas State University.
- Clark, T.J. 1998. Rebloom and display color stability of beef and pork packaged in an ultra-low oxygen modified atmosphere active packaging system. M.S. Thesis, Kansas State University.
- Campbell, R.E. 2000. Effects of internal structure, patty shrinkage, and desinewing on palatability attributes of ground beef. PhD dissertation.
- Sammel, L.M. 2000. Chemical characterization, color stability, and comparison of assays for metmyoglobin reducing ability in beef inside and outside semimembranosus. MS thesis.
- Lien, R. 2001. Effects of endpoint temperature, pork quality, and cooking factors on internal cooked color of pork chops and patties. MS thesis.
- Wendelburg, J. 2001. Physical, chemical and microbial qualities of blade tenderized prime rib. MS Thesis.
- Ballard, C. 2002. Carbon monoxide in modified atmosphere packaging. MS Thesis.

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

# CURRICULUM VITAE

Navn/Name: Sørheim, Oddvin  
Privatadresse/Private address: Lyngveien 24A, 1430 Ås  
E-mail: oddvin.sorheim@matforsk.no  
Telefon/Telephone: +47-64970100  
Arbetssted/Place of employment: MATFORSK – Norwegian Food Research Institute  
Osloveien 1, 1430 Ås  
Stilling/Present position: Sjefingeniør / Senior Research Technologist  
Dr.agric. / Ph.D.

## Utdanning/Education:

2000, Dr. agric./Ph.D., Norges landbrukshøgskole/Agricultural University of Norway  
1982, Cand. agric./M.Sc., Næringsmiddelfag, Norges landbrukshøgskole/Agricultural University of Norway

## Erfaring/Experience:

01.05. 1985 til nå ved/ to present at MATFORSK, arbeidet med forskning og industrioppdrag på kjøtt i hovedsak innen gasspakking, farge, mørhet, salting og funksjonelle egenskaper/ mainly working in research and industry consulting on packaging, color, tenderness, salting and functional properties of meat.  
01.08. 1989 til 01.06. 1990, forskningsopphold/visiting scientist, Kansas State University, Department of Animal Sciences and Industry, Manhattan, KS, USA  
01.06. 1982 til 30.04. 1985, produksjonskonsulent/consultant, Slakterienes Salgssentral/Norwegian Meat Cooperative, Oslo

## Vitenskapelige originalarbeider/Scientific publications:

- Sørheim, O., Idland, J., Halvorsen, E.C., Frøystein, T., Lea, P., Hildrum, K.I. 2001. Influence of beef carcass stretching and chilling rate on the tenderness of *M. longissimus dorsi*. Meat Science, Vol 57, pp 79-85.
- Kjos, N.P., Øverland, M., Bryhni, E.A., Sørheim, O. 2000. Food waste products in diets for growing-finishing pigs: effect on growth performance, carcass characteristics and meat quality. Acta Agric. Scand., Sect. A, Animal Sci., Vol 50, pp 193-204.
- Hunt, M.C., Sørheim, O., Slinde, E. 1999. Color and heat denaturation of myoglobin forms in ground beef. Journal of Food Science, Vol 64, pp 847-851.
- Sørheim, O., Nissen, H., Nesbakken, T. 1999. The storage life of beef and pork packaged in an atmosphere with low carbon monoxide and high carbon dioxide. Meat Science, Vol 52, pp 157-164.
- Sørheim, O., Aune, T., Nesbakken, T. 1997. Technological, hygienic and toxicological aspects of carbon monoxide used in modified-atmosphere packaging of meat. Trends in Food Science & Technology, Vol 8, pp 307-312.
- Sørheim, O., Erlandsen, T., Nissen, H., Lea, P., Høyem, T. 1997. Effects of modified atmosphere storage on colour and microbiological shelf life of normal and pale, soft and exudative pork. Meat Science, Vol 47, pp 147-155.
- Sørheim, O., Kropf, D.H., Hunt, M.C., Karwoski, M.T., Warren, K.E. 1996. Effects of modified gas atmosphere packaging on pork loin colour, display life and drip loss. Meat Science, Vol 43, pp 203-212.
- Nissen, H., Sørheim, O., Dainty, R.H. 1996. Effects of vacuum, modified atmospheres and storage temperature on the microbial flora of packaged beef. Food Microbiology, Vol 13, pp 183-191.
- Sørheim, O., Grini, J.Aa., Nissen, H., Andersen, H.J., Lea, P. 1995. Pork loins stored in carbon dioxide - Colour and microbiological shelf life. Fleischwirtschaft, 75, 679-681.

Vitenskapelige publikasjoner og presentasjoner/Other publications and presentations:

- Sørheim, O. 2000. Effects of modified atmosphere packaging on colour and microbiological shelf life of red meats. (Dr.agric./Ph.D.-thesis, Agricultural University of Norway).
- Sørheim, O., Nissen, H. 2000. Current technology for meat MAP. Int. Food Marketing & Technology, Vol 14, No 4, pp 39-42.
- Sørheim, O., Nissen, H. 1996. Modified atmosphere packaging of red meats. The European Food & Drink Review, No 4, pp 77-80.

Bok/book

- Hildrum, K.I., Dainty, R.H., Egelanddal, B., Larssen, E.G., Mielnik, J., Sørheim, O. 1996. Meat for the Consumer. MATFORSK Ås, 593 pp, ISBN 82-90394-58-6.

Annet/others

- Risvik, E., Hildrum, K.I., Russwurm jr, H., Dainty, R.H., Egelanddal, B., Green, A., Larssen, E.G., Merok, K.J., Mielnik, J., Norang, R.E., Næs, H., Olsen, M.Ø., Roskifte, P., Slinde, E., Sørheim, O. 1996. The 42nd ICoMST: "Meat for the Consumer", Norway 1-6 September 1996 (organising committee).
- Sørheim, O. 2001 - . Editorial board, Journal of Muscle Foods.

Proceedings:

- Sørheim, O., Nissen, H., Aune, T., Nesbakken, T. 2001. Use of carbon monoxide for retail meat packaging. Proc. International Animal and Agriculture Food Science Conference, Indianapolis, USA, in press.
- Sørheim, O., Lea, P., Nissen, H., Nesbakken, T. 2001. Effects of a high CO<sub>2</sub> /low CO atmosphere on colour and yield of cooked ground beef patties. Proc. 47th ICoMST, Krakow, Poland, 4 p., in press.
- Sørheim, O., Idland, J., Frøystein, T., Lea, P., Hildrum, K.I. 2000. Combined effects of aitch bone suspension and chilling rate on the tenderness of beef muscles. Proc. 46th ICoMST, 27 August - 1 September 2000, Buenos Aires, Argentina, Vol 1, pp 402-403.
- Bryhni, E.A., Kjos, N.P., Øverland, M., Sørheim, O. 1999. Food waste products in diets for growing-finishing pigs. Effect on growth performance, carcass characteristics and meat quality. Proc. 45th ICoMST, 1-6 August 1999, (Eds. Japan Society for Meat Science and Technology) Yokohama, Japan, pp 82-83.
- Sørheim, O., Nissen, H., Nesbakken, T. 1998. Color stabilization of pork chops packaged with a low level of carbon monoxide. Proc. 51st Annual Reciprocal Meat Conference: Current Advances in Meat and Poultry Color, 28 June - 1 July 1998, (American Meat Science Association) Storrs, Connecticut, USA, Vol 51, pp 187-188.
- Sørheim, O., Nissen, H., Nesbakken, T. 1997. Shelf life and colour stability of beef loin steaks packaged in a modified atmosphere with carbon monoxide., Proc. 43rd ICoMST, 27 July - 1 August 1997, (Ed. J. Bass) Auckland, New Zealand, Vol 43, pp 694-695. ISBN 0-473-04549-4.
- Sørheim, O., Lea, P., Gilde, M., Nissen, H. 1996. Effects of packaging gases on the colour of beef . Proc. 42nd ICoMST (Ed. K.I. Hildrum), MATFORSK, Lillehammer, Norway, 1-6 September 1996, pp 110-111, ISBN 82-90394-58-6.
- Hunt, M.C., Sørheim, O., Slinde, E. 1995. Effects of myoglobin form on internal cooked color development in ground beef. Proc. 41st ICoMST, American Meat Science Association, San Antonio, Texas, USA, 20-25 August 1995, Vol II, pp 394-395.



## DEPARTMENT OF HEALTH &amp; HUMAN SERVICES

Public Health Service

Food and Drug Administration  
Washington, DC 20204

Eric Greenberg  
Ungaretti and Harris  
3500 Three First National Plaza  
Chicago, IL, 60602-4405

Re: GRAS Notice No. GRN 000083

Dear Mr. Greenberg:

The Food and Drug Administration (FDA) is responding to the notice, dated August 29, 2001, that Ungaretti and Harris submitted on behalf of Pactiv Corporation (Pactiv) in accordance with the agency's proposed regulation, proposed 21 CFR 170.36 (62 FR 18938; April 17, 1997; Substances Generally Recognized as Safe (GRAS)). FDA received the notice on September 4, 2001, and designated it as GRAS Notice No. GRN 000083.

The subject of the notice is carbon monoxide (CO). The notice informs FDA of the view of Pactiv Corporation (Pactiv) that CO is GRAS, through scientific procedures, for use as a component of a gas mixture in a modified atmosphere packaging (MAP) system. The level of CO in this MAP system is 0.4 percent. The other components of the MAP system are carbon dioxide (30 percent) and nitrogen (69.6 percent). The MAP system would be used for packaging fresh cuts of case ready muscle meat and ground case ready meat to maintain wholesomeness, provide flexibility in distribution, and reduce shrinkage of the meat. The case ready meats would be removed from the MAP system prior to retail display.

As part of its notice, Pactiv includes letters from a panel of individuals (Pactiv's GRAS panel) who evaluated the data and information that are the basis for Pactiv's GRAS determination. Pactiv considers the members of its GRAS panel to be qualified by scientific training and experience to evaluate the safety of substances added to food. Pactiv's GRAS panel evaluated information and data on the chemical identity, manufacture and processing, conditions of proposed use, and estimated daily intakes of CO used in a MAP system for meat. Pactiv's GRAS panel also evaluated studies (published and unpublished) of the effects of CO used in a MAP system for meat. Members of the GRAS panel reviewed and evaluated the publicly available information summarized in the GRAS notice. Based on the data and information reviewed, Pactiv's GRAS panel concludes that CO, when produced in accordance with current good manufacturing practice and meeting appropriate food grade specifications, is GRAS, through scientific procedures under the conditions of its intended use.

The notice describes publicly available information pertaining to the identity and characteristic properties of CO. Carbon monoxide (Chemical Abstracts Service Registry Number 630-08-0) is a colorless, odorless, gas. The notice includes a list of properties of CO and identifies the

Page 2 - Mr. Greenberg

manufacturer who currently supplies CO to Pactiv. Pactiv intends to use CO at a minimum purity of 99.99 percent ("commercial grade"). Pactiv includes a list of specifications for CO with limits on the levels of other gases and considers CO of this purity to be "food grade."

The notice describes information about existing regulations and notices regarding food substances that contain CO as a significant component:

- Wood smoke, which includes CO as a component, is permitted by regulation as an ingredient in meat and poultry products under regulations issued by the U.S. Department of Agriculture (9 CFR 318.7(c)(4), 381.147(c)(4) and 424.21(c)).
- Combustion product gas, which includes CO as a component at a maximum level of 4.5 percent by volume, is approved for use in the production of beverages and other foods (except fresh meat) under FDA's regulations (21 CFR 173.350).
- Tasteless smoke, which includes CO as a primary component, is the subject of GRN 000015 for use on raw tuna, before it is frozen, to preserve its taste, aroma, texture, and color. In response to GRN 000015, FDA had no questions regarding the notifier's conclusion that tasteless smoke is GRAS under the intended conditions of use.

The notice describes the estimated consumption of CO per meal as a consequence of its intended use as a component in a MAP system for storing meat. Assuming that 30 percent of the CO present in the MAP is absorbed into the meat and that there is an 85 percent reduction of CO due to cooking the meat, Pactiv calculates a realistic intake estimate to be 0.084 milligrams (mg) CO per meal. Pactiv also calculates a worst case intake estimate to be 1.88 mg CO per meal, assuming that 100 percent of the CO present in the MAP is absorbed into the meat and that there is no reduction in CO during cooking. Pactiv cites published articles to support the assumptions used in the realistic exposure estimate and to support the conclusion that exposure to CO is safe at this level.

The notice describes published reports of studies demonstrating the technical effect and safety of using CO as a component of a MAP system (similar to the MAP system that is the subject of GRN 000083) for storing meat. These reports include published data (microbial growth profiles and odor and color data) from meat stored in MAP containing CO, CO<sub>2</sub>, and N<sub>2</sub>, and meat stored in MAP containing only CO<sub>2</sub> and N<sub>2</sub>. Pactiv concludes that the presence of CO in MAP systems allows the meat to maintain a desirable red color during storage. In addition, CO neither affects the ability of the MAP system to slow the growth of a variety of microorganisms, nor affects the characteristic odor of meat spoilage.

The notice describes an unpublished study using the MAP system that is the subject of GRN 000083. The study examined the effects of the system on initial meat color, stability of color during display, and the relationship between color deterioration and microbial growth. The notice also includes unpublished pictures that compare the ageing (color deterioration) of meats stored for 20 days in an environment of CO, CO<sub>2</sub>, and N<sub>2</sub>, to the ageing of fresh cut meat and the ageing of meat stored in a high oxygen environment. From these data, Pactiv concludes that once meat is removed from a MAP system containing CO, its color deteriorates at a similar rate to

Page 3 - Mr. Greenberg


that of meat that has not been exposed to CO. Pactiv also concludes that the use of CO in a MAP system does not result in red color life extension that could mask microbial spoilage of the meat.

Based on the information provided by Pactiv, as well as other information available to FDA, the agency has no questions at this time regarding Pactiv's conclusion that CO is GRAS under the intended conditions of use. The agency has not, however, made its own determination regarding the GRAS status of the subject use of CO. As always, it is the continuing responsibility of Pactiv to ensure that food ingredients that the firm markets are safe, and are otherwise in compliance with all applicable legal and regulatory requirements.

During its evaluation of GRN 000083, OFAS consulted with the Labeling and Consumer Protection Staff of the Food Safety and Inspection Service (FSIS) of the United States Department of Agriculture regarding the use of CO in meat products. Based on the information submitted by Pactiv, FSIS has concluded that the MAP system (ActiveTech™ 2001) as described in Pactiv's notice, and used under the conditions stated in Pactiv's notice, would be acceptable for packaging red meat cuts and ground meat. In FSIS' view, Pactiv has demonstrated that this MAP system complies with FDA's definition of a processing aid that appears in labeling regulations (21 CFR 101.100(a)(3)). There is no lasting functional effect in the food and there is an insignificant amount of carbon monoxide present in the finished product under the proposed conditions of use. As such, similar to uses of other MAP gases (e.g., nitrogen), there are no labeling issues in regard to meat cuts and ground meat packaged using this MAP. Additionally, when considering the use of a food ingredient or additive in a meat product, FSIS historically has treated each livestock species separately. However, in this case, the data submitted by Pactiv can be extrapolated to all species of livestock. If you have any additional questions, you should direct your inquiry to Dr. Robert Post, Director, Labeling and Consumer Protection Staff, Office of Policy, Program Development and Evaluation, Food Safety and Inspection Service, 300 12th Street, SW, Room 602, Washington, DC 20250-3700. The telephone number of his office is (202) 205-0279 and the FAX number is (202) 205-3625.

In accordance with proposed 21 CFR 170.36(f), a copy of the text of this letter, as well as a copy of the information in your notice that conforms to the information in proposed 21 CFR 170.36(c)(1), is available for public review and copying on the homepage of the Office of Food Additive Safety (on the Internet at <http://www.cfsan.fda.gov/~lrd/foodadd.html>).

Sincerely,



Alan M. Rulis, Ph.D.

Director

Office of Food Additive Safety

Center for Food Safety

and Applied Nutrition